



# Particle collisions near static spherically symmetric black holes

Eva Hackmann<sup>a</sup>, Hemwati Nandan<sup>b,c,\*</sup>, Pankaj Sheoran<sup>d</sup>

<sup>a</sup> ZARM, University of Bremen, Am Fallturm 2, 28359 Bremen, Germany

<sup>b</sup> Department of Physics, Gurukul Kangri Vishwavidyalaya, Haridwar-249 407, India

<sup>c</sup> Center for Space Research, North-West University, Mafikeng 2745, South Africa

<sup>d</sup> Instituto de Física y Matemáticas, Universidad Michoacana de San Nicolás de Hidalgo, Edificio C-3, 58040 Morelia, Michoacán, Mexico



## ARTICLE INFO

### Article history:

Received 15 July 2020

Received in revised form 26 September 2020

Accepted 4 October 2020

Available online 7 October 2020

Editor: A. Volovich

### Keywords:

Particle collision

Center of mass

Black holes

## ABSTRACT

It has been shown by Bañados, Silk and West (BSW) that the center of mass energy ( $E_{\text{cm}}$ ) of test particles starting from rest at infinity and colliding near the horizon of a Schwarzschild black hole is always finite. In this short note, we extend the BSW scenario and study two particles with different energies colliding near the horizon of a static spherically symmetric black hole. Interestingly, we find that even for the static spherically symmetric (i.e., Schwarzschild like) black holes it is possible to obtain an arbitrarily high  $E_{\text{cm}}$  from the two test particles colliding near but outside of the horizon of a black hole, if one fine-tunes the parameters of geodesic motion.

© 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

In 2009, Bañados, Silk and West (BSW) first showed [1] that an extremal axially symmetric rotating black hole (BH) can act as a particle accelerator: an extremely high amount of center of mass energy ( $E_{\text{cm}}$ ) can be produced from the collision of two test particles starting from rest at infinity. This is in contrast to static spherical symmetric BHs where  $E_{\text{cm}}$  is always finite. They pointed out that the extremal rotating BHs may therefore be used as an important probe of high energy scale physics. Since then collisions of geodesic (and charged) particles near the horizon of BHs [2–15] have been analysed. The study of BSW effect is not limited to geodesic particles only but extended to spinning, accelerated particles as well [16–18]. However, all these different cases of collisions of particles (i.e., geodesic, charged, spinning, and accelerated) in the vicinity of a BH are accompanied by some serious limitations. For the case of geodesic particles, extremely high  $E_{\text{cm}}$  is obtained if the collision occurs near the horizon of an extremal rotating BH and the energy  $E$  and the angular momentum  $L$  of one of the colliding particles meet a critical condition [1]. On the other hand, when the collision of spinning particles is considered far away from the event horizon of a BH, the main difficulty is the unavoidable occurrence of a superluminal region (i.e. the region where the four velocity of the spinning particle changes from time-


like to spacelike) [17]. Finally, one can obtain the arbitrarily high  $E_{\text{cm}}$  even for the static spherically symmetric case, but it requires that one of the colliding particles must be accelerated, which in turn requires the existence of some external source (i.e. which act as an engine to accelerate the particle) [18]. With these limitations on the BSW effect, it is natural to investigate whether a BH in its simplest form (i.e. static spherically symmetric BH) can act as an accelerator for colliding geodesic particles.

In this paper, we study a collision of two particles near the horizon of a static spherically symmetric BH. We assume that one of the colliding particles originates near the horizon of the BH and possesses very small radial motion with energy such that  $E_1 = \epsilon \ll 1$ . On the contrary, the second particle is generic, for instance falling from rest at infinity with energy  $E_2$  as shown in Fig. 1. This situation for the colliding particles was mentioned in passing in [19,20] for Schwarzschild spacetime but has not been studied in detail yet. However, it is worth mentioning that a similar type of scenario, when one of the colliding particles possesses the energy  $E = 0$ , was also discussed in [15], but only for a special 2 + 1 dimensional Bañados-Teitelboim-Zanelli (BTZ) BH spacetime. Here we elaborate this scenario for general spherically symmetric spacetimes, and some novel features of the particle acceleration process in the vicinity of a BH are reported.

The paper is organized as follows. In Sec. 2, we briefly describe the metric of the static spherically symmetric BH, the equation of motion of the geodesic particles, and the expression for the center of mass energy  $E_{\text{cm}}$ . In Sec. 3, we explain how a static spherically symmetric BH can act as a particle accelerator for two geodesic par-

\* Corresponding author.

E-mail addresses: [eva.hackmann@zarm.uni-bremen.de](mailto:eva.hackmann@zarm.uni-bremen.de) (E. Hackmann), [hnandan@associate.iucaa.in](mailto:hnandan@associate.iucaa.in) (H. Nandan), [hukmipankaj@gmail.com](mailto:hukmipankaj@gmail.com) (P. Sheoran).



**Fig. 1.** Schematic view of two geodesic particles with different energies ( $E_1 \neq E_2$ ) colliding near the horizon of static and spherically symmetric BH.

ticles colliding in its vicinity. Finally, in Sec. 4, we summarize our main results with a concluding remark. Throughout this paper, we choose the  $(-, +, +, +)$  signature for the metric tensor, Greek indices run from 0 to 3, and both the fundamental constants  $G$  and  $c$  are set equal to unity.

## 2. Collision of geodesic particles

A static and spherically symmetric spacetime is given by,

$$g = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (1)$$

We assume that we have a BH solution with an event horizon located at  $r = r_H$ . Dependent on the choice of the metric function  $f(r)$  we may have additional horizons, for instance a cosmological or a Cauchy horizon. The two Killing vectors  $u_t = -E$  and  $u_\phi = L$  thus give rise to two constants of motion as below,

$$-E = g_{tt}\dot{t}, \quad (2)$$

$$L = g_{\phi\phi}\dot{\phi}, \quad (3)$$

where a dot denotes a derivative with respect to the proper time of the particle. Due to spherical symmetry, without loss of generality, we entirely restrict to the plane  $\theta = \pi/2$ . From the normalization condition  $g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu = -1$ , one can then obtain the following equation of geodesic motion,

$$\dot{r}^2 = E^2 - f(r) \left(1 + \frac{L^2}{r^2}\right) =: R(r). \quad (4)$$

Let us now consider two particles of masses  $m_1$  and  $m_2$  moving on geodesics in a BH spacetime. Their energy in the center of mass frame is given by,

$$E_{\text{cm}}^2 = -g_{\mu\nu}(p_1^\mu + p_2^\mu)(p_1^\nu + p_2^\nu) \quad (5)$$

$$= m_1^2 + m_2^2 - 2g_{\mu\nu}p_1^\mu p_2^\nu, \quad (6)$$

where  $p_i^\mu$  denotes the four momentum of the two particles. If we assume  $m_1 = m_2 = m$ , one can easily obtain,

$$\frac{E_{\text{cm}}^2}{2m^2} = 1 - \frac{L_1 L_2}{r^2} + \frac{E_1 E_2 - \sqrt{R_1(r)R_2(r)}}{f(r)}, \quad (7)$$

where the indices refer to the two different particles. Regardless of the apparent singularity at the horizon, where  $f(r_H) = 0$ , for generic particles the center of mass energy remains finite as the nominator also vanishes at  $r = r_H$ . The limit is given by,

$$\begin{aligned} \lim_{r \rightarrow r_H} \frac{E_{\text{cm}}^2}{2m^2} &= 1 - \frac{L_1 L_2}{r_H^2} + \frac{E_1}{2E_2} \left(1 + \frac{L_2^2}{r_H^2}\right) \\ &+ \frac{E_2}{2E_1} \left(1 + \frac{L_1^2}{r_H^2}\right). \end{aligned} \quad (8)$$

In the original paper by BSW, it was assumed from the beginning that the two particles start from rest at infinity, implying  $E_1 = E_2 = 1$ . In this case, we see that the center of mass energy is always finite. Later, it was pointed out by Zaslavskii that for charged particles in a Reissner-Nordström spacetime, or geodesic particles in a stationary (but non-static) and axially symmetric spacetime the parameters can be fine-tuned such that the expression Eq. (8) becomes arbitrarily large. For the Schwarzschild spacetime, Grib et al. [19,20] show that, in the limit  $r \rightarrow r_H$  considered above, if the point of collision coincides with the radial turning point  $\dot{r} = 0$  of one of the particles than the energy may also grow without bound. We generalise here the argument from [19,20], allowing for general static spherically symmetric spacetimes, and removing the condition that the point of collision coincides with the radial turning point.

## 3. Arbitrarily large center of mass energy

We immediately notice from the limit Eq. (8) that the center of mass energy diverges if one of the particles has zero energy. However, as evident from eq. (4), this means that geodesic motion is impossible. Instead, we may consider a particle with a very low energy. Assume that particle 1 has  $E_1 = \epsilon \ll 1$  and particle 2 is generic. Then from Eq. (8), one can observe,

$$\lim_{r \rightarrow r_H} \frac{E_{\text{cm}}^2}{2m^2} = \frac{E_2}{2\epsilon} \left(1 + \frac{L_1^2}{r_H^2}\right) + \mathcal{O}(\epsilon^0). \quad (9)$$

This result was basically derived in [19,20], although we did not explicitly assume that  $\dot{r}_1 = 0$  at the horizon. However, as mentioned above, geodesic motion with  $E_1 \ll 1$  is only possible if  $\dot{r}_1 = 0$  very close to the horizon. Note that if both particles have  $E \ll 1$ , the center of mass energy becomes again finite.

More generally, we can also obtain arbitrarily large center of mass energy for collisions close to the horizon, but outside of it. Let  $r_c$  be the point of collision with  $r_c - r_H \ll 1$ . We rewrite the metric function  $f$  as  $f(r) = (r - r_H)\tilde{f}(r)$ , in particular  $\tilde{f}(r) = 1/r$  for Schwarzschild spacetime. Let us choose for, say particle 1,  $E_1 = C_1\sqrt{r_c - r_H}$ . Here  $C_1$  is a positive constant, taken such that geodesic motion is possible in the region  $(r_H, r_c)$ . We then find the following expression from Eq. (7) at the point of collision  $r_c$ ,

$$\begin{aligned} \frac{E_{\text{cm}}^2}{2m^2} &= 1 - \frac{L_1 L_2}{r_c^2} + \frac{E_2 C_1 \sqrt{r_c - r_H}}{(r_c - r_H)\tilde{f}(r_c)} - \frac{1}{(r_c - r_H)\tilde{f}(r_c)} \\ &\times \left[ (r_c - r_H) \left( C_1^2 - \tilde{f}(r_c) \left(1 + \frac{L_1^2}{r_c^2}\right) \right) \right. \\ &\times \left. \left( E_2^2 - (r_c - r_H)\tilde{f}(r_c) \left(1 + \frac{L_2^2}{r_c^2}\right) \right) \right]^{\frac{1}{2}} \end{aligned} \quad (10)$$

$$\begin{aligned} &= 1 - \frac{L_1 L_2}{r_c^2} + \frac{E_2 C_1}{\sqrt{r_c - r_H}\tilde{f}(r_c)} \\ &- \frac{E_2 \sqrt{C_1^2 - \tilde{f}(r_c) \left(1 + \frac{L_1^2}{r_c^2}\right)}}{\sqrt{r_c - r_H}\tilde{f}(r_c)} + \mathcal{O}(\sqrt{r_c - r_H}). \end{aligned} \quad (11)$$

Therefore, for small but nonvanishing  $r_c - r_H$ , we can obtain arbitrarily large center of mass energy provided we can choose the constant  $C_1$  such that

$$C_1 - \sqrt{C_1^2 - \tilde{f}(r_c) \left(1 + \frac{L_1^2}{r_c^2}\right)} > 0. \quad (12)$$

For instance, we could choose

$$C_1 = \sqrt{1 + \frac{\tilde{f}(r_c)(r_c^2 + L_1^2)}{r_c^2}}. \quad (13)$$

Then, at the point of collision we find from Eq. (4) that  $\dot{r}^2|_{r=r_c} = (r_c - r_H) > 0$  as well.

A caveat of both calculations explained above is that the particle 1 is assumed to have a very small radial motion velocity although it is very close to the horizon. This is a very special situation, which will usually not be encountered, and probably will only arise if the particle originated from a foregoing collision.

#### 4. Summary and conclusion

In this short paper, we delved into the BSW effect and studied the collision of two geodesic particles near the horizon of a static spherically symmetric BH, generalising a scenario shortly discussed in [19,20]. It is assumed that one of the colliding particles (say particle 1) originates near the horizon of the BH and possesses a very small radial velocity and energy  $E_1 \ll 1$  (despite the fact that it is very close to  $r_H$ ). On the other hand, the second colliding particle with energy  $E_2$  is considered to be generic, for instance the usual particle coming from rest at infinity. It is shown that when the collision of the above mentioned geodesic particles takes place near the event horizon of a static spherically symmetric BH, it is still possible to have an arbitrarily high  $E_{cm}$ . This is in contrast to the case when the colliding particles need to have a spin as pointed out in [16,17], a charge as considered in [21], or to the case when one needs an external engine (i.e. source) to accelerate one of the colliding particles in order to achieve arbitrarily high  $E_{cm}$  for static spherically symmetric BHs as pointed out in [18].

It is noteworthy to mention that even a static spherically symmetric BH can also act as a particle accelerator for geodesic particles and one does not need special types of particles (i.e. spinning or accelerated) in order to obtain arbitrarily high  $E_{cm}$ . However, the scenario we discussed here is very special and is only possible if the particle 1 is originated near to horizon of BH by some process, maybe through a foregoing collision of particles.

In response to an earlier version of this paper, Zaslavskii [22] analysed the physical relevance of the scenario under discussion here. He found that for the Schwarzschild spacetime it is unphysical, requiring infinite forces to place the critical particle near the horizon. However, for the extremal Reissner-Nordström spacetime the critical (neutral) particle could be produced by a foregoing collision of charged particles. For other static and spherically symmetric spacetimes the question of physical relevance remains open.

#### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Acknowledgements

HN would like to thank Science and Engineering Research Board (SERB), New Delhi, India for financial support through grant

no. EMR/2017/000339. HN is also thankful to IUCAA, Pune, India (where a part of the work was completed) for support in form of academic visits under its Associateship programme. He would also like to acknowledge the facility at ICARD, Gurukula Kangri Vishwavidyalaya, Haridwar, India. PS would like to thank *Programa de Desarrollo Profesional Docente* (PRODEP) of the *Secretaría de Educación Pública* (SEP) of the Mexican government, for providing the financial support. EH is thankful for support from the research training group RTG 1620 “Models of Gravity” and the Cluster of Excellence EXC-2123 Quantum Frontiers - 390837967, both funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation).

#### References

- [1] M. Banados, J. Silk, S.M. West, Kerr black holes as particle accelerators to arbitrarily high energy, *Phys. Rev. Lett.* 103 (2009) 111102, <https://doi.org/10.1103/PhysRevLett.103.111102>, arXiv:0909.0169.
- [2] O.B. Zaslavskii, Acceleration of particles as universal property of rotating black holes, *Phys. Rev. D* 82 (2010) 083004, <https://doi.org/10.1103/PhysRevD.82.083004>, arXiv:1007.3678.
- [3] O.B. Zaslavskii, Acceleration of particles by black holes: general explanation, *Class. Quantum Gravity* 28 (2011) 105010, <https://doi.org/10.1088/0264-9381/28/10/105010>, arXiv:1011.0167.
- [4] S.-W. Wei, Y.-X. Liu, H. Guo, C.-E. Fu, Charged spinning black holes as particle accelerators, *Phys. Rev. D* 82 (2010) 103005, <https://doi.org/10.1103/PhysRevD.82.103005>, arXiv:1006.1056.
- [5] S.-W. Wei, Y.-X. Liu, H.-T. Li, F.-W. Chen, Particle collisions on stringy black hole background, *J. High Energy Phys.* 12 (2010) 066, [https://doi.org/10.1007/JHEP12\(2010\)066](https://doi.org/10.1007/JHEP12(2010)066), arXiv:1007.4333.
- [6] M. Kimura, K.-i. Nakao, H. Tagoshi, Acceleration of colliding shells around a black hole: validity of the test particle approximation in the Banados-Silk-West process, *Phys. Rev. D* 83 (2011) 044013, <https://doi.org/10.1103/PhysRevD.83.044013>, arXiv:1010.5438.
- [7] T. Harada, M. Kimura, Collision of two general geodesic particles around a Kerr black hole, *Phys. Rev. D* 83 (2011) 084041, <https://doi.org/10.1103/PhysRevD.83.084041>, arXiv:1102.3316.
- [8] A. Galajinsky, Near horizon geometry of extremal black holes and Banados-Silk-West effect, *Phys. Rev. D* 88 (2013) 027505, <https://doi.org/10.1103/PhysRevD.88.027505>, arXiv:1301.1159.
- [9] S.G. Ghosh, P. Sheoran, M. Amir, Rotating Ayón-Beato-García black hole as a particle accelerator, *Phys. Rev. D* 90 (2014) 103006, <https://doi.org/10.1103/PhysRevD.90.103006>, arXiv:1410.5588.
- [10] M. Amir, S.G. Ghosh, Rotating Hayward's regular black hole as particle accelerator, *J. High Energy Phys.* 07 (2015) 015, [https://doi.org/10.1007/JHEP07\(2015\)015](https://doi.org/10.1007/JHEP07(2015)015), arXiv:1503.08553.
- [11] S.G. Ghosh, M. Amir, Horizon structure of rotating Bardeen black hole and particle acceleration, *Eur. Phys. J. C* 75 (2015) 553, <https://doi.org/10.1140/epjc/s10052-015-3786-x>, arXiv:1506.04382.
- [12] A. Zakria, M. Jamil, Center of mass energy of the collision for two general geodesic particles around a Kerr-Newman-Taub-NUT black hole, *J. High Energy Phys.* 05 (2015) 147, [https://doi.org/10.1007/JHEP05\(2015\)147](https://doi.org/10.1007/JHEP05(2015)147), arXiv:1501.06306.
- [13] M. Amir, F. Ahmed, S.G. Ghosh, Collision of two general particles around a rotating regular Hayward's black holes, *Eur. Phys. J. C* 76 (2016) 532, <https://doi.org/10.1140/epjc/s10052-016-4365-5>, arXiv:1607.05063.
- [14] P.A. González, M. Olivares, E. Papantonopoulos, Y. Vásquez, Motion and collision of particles in a rotating linear dilaton black hole, *Phys. Rev. D* 97 (2018) 064034, <https://doi.org/10.1103/PhysRevD.97.064034>, arXiv:1802.01760.
- [15] K. Ogasawara, N. Tsukamoto, Effect of the self-gravity of shells on a high energy collision in a rotating Bañados-Teitelboim-Zanelli spacetime, *Phys. Rev. D* 99 (2019) 024016, <https://doi.org/10.1103/PhysRevD.99.024016>, arXiv:1810.03294.
- [16] C. Armas, M. Bañados, B. Koch, Collisions of spinning massive particles in a Schwarzschild background, *Class. Quantum Gravity* 33 (2016) 105014, <https://doi.org/10.1088/0264-9381/33/10/105014>, arXiv:1510.01223.
- [17] O.B. Zaslavskii, Schwarzschild black hole as particle accelerator of spinning particles, *Europhys. Lett.* 114 (2016) 30003, <https://doi.org/10.1209/0295-5075/114/30003>, arXiv:1603.09353.
- [18] O. Zaslavskii, Schwarzschild black hole as a particle accelerator, *JETP Lett.* 111 (2020) 260–263, <https://doi.org/10.1134/S0021364020050033>, arXiv:1910.04068.
- [19] A.A. Grib, Y.V. Pavlov, O.F. Piattella, On collisions with unlimited energies in the vicinity of Kerr and Schwarzschild black hole horizons, *Gravit. Cosmol.* 18 (2012) 70–75, <https://doi.org/10.1134/S0202289312010094>, arXiv:1203.4952.
- [20] A.A. Grib, Yu.V. Pavlov, O.F. Piattella, O.F. Piattella, High energy processes in the vicinity of the Kerr's black hole horizon, *Int. J. Mod. Phys. A* 26 (2011) 3856–3867, <https://doi.org/10.1142/S0217751X11054310>, arXiv:1105.

- 1540, Int. J. Mod. Phys. Conf. Ser. 03 (2011) 342, <https://doi.org/10.1142/S2010194511001449>.
- [21] O.B. Zaslavskii, Acceleration of particles by nonrotating charged black holes, JETP Lett. 92 (2010) 571–574, <https://doi.org/10.1134/S0021364010210010>, arXiv:1007.4598, Pis'ma Zh. Eksp. Teor. Fiz. 92 (2010) 635.
- [22] O. Zaslavskii, Special case of the Bañados-Silk-West effect, Phys. Rev. D 102 (2020) 044051, <https://doi.org/10.1103/PhysRevD.102.044051>, arXiv:2006.11552.