

An EFT hunter's guide to two-to-two scattering: HEFT and SMEFT on-shell amplitudes

Hongkai Liu,^a Teng Ma,^{a,b} Yael Shadmi^a and Michael Waterbury^a

^a*Physics Department, Technion — Israel Institute of Technology,
Technion city, Haifa 3200003, Israel*

^b*Department of Physics and IFAE-BIST, Universitat Autònoma de Barcelona,
E-08193 Bellaterra, Barcelona, Spain*

E-mail: liu.hongkai@campus.technion.ac.il, tma@ifae.es,
yshadmi@physics.technion.ac.il, mwaterbury@campus.technion.ac.il

ABSTRACT: We derive the contact terms contributing to the four-point amplitudes of the standard model particles, keeping terms with up to quartic energy growth. Imposing just the unbroken low-energy symmetry, and treating the electroweak gauge bosons and the Higgs as independent degrees of freedom, we obtain the most general four-point contact-term amplitudes, corresponding to the Higgs Effective Field Theory (HEFT) framework. The contact terms are spanned by a basis of Stripped Contact Terms, which carry the polarization information, multiplied by polynomials in the Mandelstam invariants. For terms with quadratic energy growth, we also derive the low-energy Standard Model Effective Field Theory (SMEFT) predictions, via on-shell Higgsing of the massless SMEFT contact terms. We discuss several aspects of bottom-up versus top-down on-shell derivations of the HEFT and SMEFT amplitudes, highlighting in particular the simple counting of HEFT dimensions in the on-shell approach and the transparent relation between perturbative unitarity and gauge-invariance in the little-group covariant massive spinor formalism. Our results provide a formulation of Effective Field Theory analyses directly in terms of observable quantities. For terms with quadratic energy growth, we also provide the mapping to the Warsaw basis.

KEYWORDS: Effective Field Theories, Scattering Amplitudes

ARXIV EPRINT: [2301.11349](https://arxiv.org/abs/2301.11349)

Contents

1	Introduction	1
2	Preliminaries	4
3	Four-point contact terms at $\mathcal{O}(E^2)$	6
3.1	HEFT contact terms	6
3.2	SMEFT contact terms	8
4	Four-point contact terms at E^3 and E^4	11
4.1	Bosonic amplitudes with all massive particles	12
4.1.1	$hhhh$	12
4.1.2	$Zhhh$	12
4.1.3	$ZZhh$	12
4.1.4	W^+W^-hh	13
4.1.5	W^+W^-Zh	13
4.1.6	$ZZZh$	13
4.1.7	W^+W^-ZZ	13
4.1.8	$W^+W^+W^-W^-$	14
4.1.9	$ZZZZ$	14
4.2	Fermionic amplitudes with all massive particles	14
4.2.1	$f^c f h h$	14
4.2.2	$f^c f Z h$ and $f^c f' W h$	14
4.2.3	$f^c f^c f f$	15
4.2.4	$W^+W^- f^c f$ and $W Z f^c f'$	15
4.2.5	$ZZ f^c f$	16
4.3	Bosonic amplitudes with massless vectors	16
4.3.1	$\gamma h h h$	16
4.3.2	$\gamma Z h h$	17
4.3.3	$g g h h$ and $\gamma \gamma h h$	17
4.3.4	$g g h$	17
4.3.5	$g g \gamma h$ and $\gamma \gamma \gamma h$	17
4.3.6	$g g Z h$ and $\gamma \gamma Z h$	17
4.3.7	$\gamma Z Z h$	17
4.3.8	$\gamma W W h$	18
4.3.9	$\gamma \gamma \gamma \gamma$	18
4.3.10	$g g g Z$, $g g g \gamma$ and $\gamma \gamma \gamma Z$	18
4.3.11	$g g Z Z$ and $\gamma \gamma Z Z$	18
4.3.12	$g g W W$ and $\gamma \gamma W W$	18
4.3.13	$\gamma Z W W$	19
4.3.14	$\gamma Z Z Z$	19

4.3.15	$gg\gamma\gamma$	19
4.3.16	$gg\gamma Z$	19
4.3.17	$gggg$	19
4.4	Fermionic amplitudes with massless vectors	20
4.4.1	$f^c f \gamma h$ and $f^c f gh$	20
4.4.2	$ggf^c f$ and $\gamma\gamma f^c f$	20
4.4.3	$\gamma g f^c f$	20
4.4.4	$\gamma Z f^c f, \gamma W f^c f', gZ f^c f$ and $gW f^c f'$	21
5	Conclusions	21
A	$WW hh$: on-shell construction of the HEFT and SMEFT amplitudes and on-shell Higgsing	22
A.1	The structure of the full amplitude	22
A.2	Four-point contact terms from on-shell Higgsing	24
A.3	The $WW h$ coupling from on-shell Higgsing	26
B	Energy-growth of factorizable amplitudes	28

1 Introduction

Precise measurements of the interactions of the standard-model (SM) particles, and in particular, the electroweak bosons and the top, will be a focus of the LHC program in the coming decade. These interactions can be systematically parameterized in terms of Effective Field Theory (EFT) Lagrangians, which in principle provide a model-independent framework for indirect searches for new physics. Much of the collider EFT program has been guided by the Standard Model EFT (SMEFT), whose starting point is the unbroken $SU(3)\times SU(2)\times U(1)$ theory with a single Higgs doublet, focusing in particular on dimension-six operators [1–3]. Even in the SMEFT framework, it is plausible however that a given set of heavy fields couple differently to different SM fields. Different SMEFT operator bases are therefore better suited to describe the effect of different UV models [3–6], and truncating the EFT at dimension-six may moreover leave out important effects (see for example [7–11]). Furthermore, some extensions of the SM are not captured by the SMEFT (at least at low-dimensions), and would lead at low energies to the framework known as the Higgs EFT (HEFT) [12–25]. These include for example models featuring fields which get their masses from electroweak symmetry breaking (EWSB), or fields which provide additional sources of this breaking [17, 26–28].

The on-shell bootstrap (see for example [29, 30]) provides a natural avenue for a bottom-up construction of EFTs. EFT extensions of the SM can be formulated in this approach directly in terms of the physical observables of interest, namely, the scattering amplitudes of the known SM particles, with a one-to-one mapping of EFT operators and contact-term

amplitudes [31–33]. Furthermore, since they are not obscured by field redefinitions and operator redundancies, questions such as the distinctions between different EFT extensions, or the assignments of EFT dimensions, are concretely phrased in terms of physical quantities.

In this paper, we derive the full set of four-point contact-term amplitudes, featuring the SM massive and massless particles, keeping terms with up to quartic energy growth. Together with the three-point amplitudes listed in [33], these determine the EFT predictions for four-point amplitudes with this energy growth. These amplitudes are the most interesting objects for phenomenological purposes, since two-to-two scattering processes, followed by two- and three-particle decays, are the ones where we can hope to get the most data. Our results are collected in tables 1 and 3 and section 4. The low-energy E^2 HEFT contact terms are given in table 1. The low-energy E^2 SMEFT contact terms appear in table 3 and are mapped to the massless SMEFT contact terms [34] collected in table 2, where we also give the relations to Warsaw basis operators. Section 4 contains the low-energy E^4 HEFT contact terms.

Indeed, on-shell scattering amplitudes have emerged in recent years as a powerful method for constructing EFT Lagrangians [31–42]. Bottom-up constructions of SM, or SM-like amplitudes were discussed in [33, 43–49]. The emergence of symmetry from the amplitude bootstrap, and its relation to the geometry of field space was studied for instance in [50–52].

We employ two types of on-shell constructions. The first is purely bottom-up and gives HEFT amplitudes. The second is top-down and starts from the massless amplitudes of the unbroken theory, yielding the SMEFT low-energy contact terms. We now sketch these in turn. Various methods for constructing generic contact-term bases for massless amplitudes were described in [26, 34, 38, 53, 54]. The construction of generic massive contact terms using little-group covariant spinors was discussed in [33, 43, 48, 55–57].

We first derive the most general four-point contact terms involving the SM particles consistent with $SU(3) \times U(1)_{EM}$ symmetry and baryon and lepton number conservation. Since they are built in terms of the broken-phase electroweak sector, with the physical Higgs h and the massive W and Z treated as independent degrees of freedom, the resulting amplitudes are valid beyond the SMEFT. In particular, any tree-amplitude featuring the SM particles with Wilson coefficients determined by the running to the energy scale of interest can be spanned by these contact terms. Thus, the contact terms derived in this way correspond to HEFT amplitudes. Our analysis extends [33], which derived the three-point SM amplitudes and one four-point example, to include the complete set of four-point contact terms.

To construct the independent contact terms, we use the strategy of [53, 55]: working with the little-group covariant massive spinor formalism [43], the basic building blocks of the basis are Stripped Contact Terms (SCTs), which are massive spinor structures with no additional factors of Mandelstam invariants. To get the full set of contact terms, each SCT is then multiplied by an expansion in these invariants. Note that the SCTs carry the little-group weights of the external particles, and encode the information on their polarizations. The expansion in the Mandelstam invariants on the other hand only depends on the scattering angles, and corresponds to the derivative expansion of EFT

Lagrangians. Generic four-point SCT bases for spins 0, 1/2, and 1 were given in [55]. Starting from these, we specify to the SM particle content, impose the low-energy symmetry and (anti)symmetrize over identical particles. Partial results on the electroweak sector contact terms were derived in [55], and our analysis extends these to the full set of massive and massless SM four-points. For each four-point contact term, we indicate the dimension of the corresponding HEFT operator, namely, the operator which generates this contact term at leading order, and the dimension at which this contact term can be generated in the SMEFT.

Turning to the construction of SMEFT contact terms, one way to proceed, which relies on low-energy input only, is to start from the HEFT contact terms and impose perturbative unitarity [33]. As we will see in section 2, the equivalence of perturbative unitarity and gauge invariance is clearly exposed when the amplitudes are written using the little-group covariant massive spinor formalism.

To recover all the SMEFT relations from this bottom-up approach, however, one needs to consider a sufficiently large set of amplitudes, including in particular higher-point amplitudes. Instead, one can start from the massless SMEFT contact terms and “Higgs” these to obtain the massive contact terms [47]. In the little group-covariant massive spinor formalism, massless SCTs featuring just fermions and vectors are simply bolded into massive SCTs. SCTs featuring an external scalar line give rise to two types of massive contact terms. Directly bolding the massless SCT gives a massive SCT with an external scalar line — a physical Higgs. Massless SCTs featuring a scalar momentum p bold into a massive vector line, with $p \rightarrow \mathbf{p}\rangle[\mathbf{p}$. This Higgsing relies on Lorentz symmetry, specifically the little group transformations of the SCTs in the massless and massive theory, and exploits the simple relations between the two in the massive formalism [43].

Two examples of four-point dimension ≤ 8 SMEFT amplitudes, namely $WWhh$ and $\bar{u}dWh$ were derived in [47]. Here we extend these results to all the SM particles, but only include dimension-six contributions. Our starting point is thus the dimension-six massless SMEFT contact terms derived in [34]. Higgsing these as described above, we obtain the massive SMEFT contact terms, recovering all the E^2 HEFT contact terms, with Wilson coefficients dictated by the SMEFT. In the bosonic sector, the number of dimension-six independent Wilson coefficients is reduced from eight in the HEFT to six in the SMEFT. Additional relations appear in the fermionic amplitudes.

Bottom-up derivations of HEFT *operators* appeared recently in [41, 58–60]. Ref. [60] presented the list of HEFT operators corresponding to four-point SCTs with Higgs external legs. (These operators are referred to as primary operators in [60].) Where they overlap, our results agree with the operator counting of [59, 60]. Hilbert series methods for counting independent EFT operators [2, 61, 62] were also extended recently to the case of massive theories and in particular to the HEFT [63].

This paper is organized as follows. In section 2, we review some elements of the little-group covariant massive spinors and SCT construction. We explain our normalization of SCTs with inverse powers of the mass and the cutoff, and the implications for identifying the operator dimensions and perturbative unitarity. We also discuss the equivalence of perturbative unitarity and gauge invariance, and comment on the differences between

SMEFT and HEFT amplitudes from the point of view of locality and analyticity. In section 3.1 we derive the generic — or HEFT — SM amplitudes with up to E^2 growth. These are shown in table 1. In section 3, we derive the analogous SMEFT amplitudes. The unbroken SMEFT contact terms are reviewed in table 2, where we also relate their Wilson coefficients to those of the Warsaw basis. We then list the massive SMEFT contact terms in table 3. Thus, each kinematic structure in the physical amplitudes can be associated with a specific operator in the Warsaw basis. In section 4, we derive the remaining HEFT amplitudes featuring cubic or quartic energy growth. For completeness, we flesh out the details of on-shell Higgsing in appendix A using the $WWhh$ amplitude as an example. We discuss the general structure of the low-energy amplitude, explain the derivation of four-point contact terms, and derive the dimension-six correction to the WWh amplitude from the relevant massless factorizable six-point amplitude. Finally, in appendix B, we list the leading high-energy behavior of the generic low-energy factorizable four-point amplitudes.

2 Preliminaries

Each four-point amplitude consists of a factorizable part, which depends on the three-point couplings; and a non-factorizable part, which is purely local and contains the four-point contact terms. The independent parameters entering the amplitude are thus the renormalizable and non-renormalizable three-point couplings, as well as the coefficients of independent four-point contact terms. Together with the three-point couplings given in ref. [33], the four-point couplings we will list here parameterize the most general SM EFT amplitudes, and allow for the construction of the full four-point amplitude.

We derive the contact terms of the massive and massless SM particles below the electroweak breaking scale. For the most part, we assume baryon- and lepton number conservation, but we will comment on the modifications to the contact terms in the presence of Majorana neutrinos. The low energy theory features several dimensionful parameters, namely the particle masses, and the cutoff, which we denote by $\bar{\Lambda}$. Neglecting fermion masses apart from the top, the masses are parametrically of the same order, and the contact terms can be written as a double expansion in m/E and $E/\bar{\Lambda}$, where m denotes the common mass scale and E is the energy. If we only impose $SU(3)\times U(1)$, the contact terms we derive a priori describe HEFT amplitudes. To obtain the low-energy SMEFT contact terms, we start from the SMEFT contact terms at high energies, with $SU(2)\times U(1)$ broken by the vacuum expectation value (VEV) v of a single Higgs doublet. The low-energy amplitudes then involve two dimensionful parameters, namely v and the SMEFT cutoff, which we denote by Λ . A large hierarchy between v and the cutoff is only possible in the SMEFT, where $SU(2)\times U(1)$ is linearly realized at Λ . Thus, for HEFT amplitudes it is appropriate to set $\bar{\Lambda} = v$. On the other hand, in the SMEFT, with lepton-number conservation, the massless amplitudes feature only even powers of Λ , and typically $1/\bar{\Lambda}^2$ in the low-energy SMEFT amplitudes maps to $1/\Lambda^2$, while $1/\bar{\Lambda}$ maps to v/Λ^2 .

The amplitudes are written in terms of spinor variables [64], using the little-group-covariant bolded spinor formalism of [43] for massive particles. We summarize the essentials of this formalism here. For more detail, and for explicit expressions for the spinors, we refer

the reader to [33, 43]. Our conventions for the spinors and their high energy limits follow ref. [33]. An external massive particle i of momentum p_i is described by a pair of massless spinors, $\mathbf{i}^{M=1,2}$. Here and in the following, \mathbf{i} stands for either \mathbf{i} or \mathbf{i} . For each external fermion i , the amplitude contains one factor of $\mathbf{i}^{M=1,2}$ with $M = 1, 2$ corresponding to positive and negative helicity respectively for square spinors, and conversely for angle spinors. For each external vector i , the amplitude contains $\mathbf{i}^{\{M\mathbf{i}\}^N}$. For square spinors, $(MN) = 11$, $(MN) = 22$, and $\{12\}$, correspond to positive, negative, and zero polarizations, respectively. Boldface indicates symmetrization over vector indices, but we use it for any massive spinor or momentum to distinguish them from massless ones.

The different spinor structures contributing to a given amplitude can be classified according to their *helicity category*, namely, the helicities of the external particles in the massless spinor structure obtained by naively unbolding the massive structure [55]. Thus for example, the structure $\mathbf{[12][12]}$, which can appear in amplitudes with two vectors, 1 and 2, and two scalars, is in the $++00$ helicity category, since the unbolded $[12][12]$ corresponds to a $++00$ helicity amplitude. In contrast, $\mathbf{[12]\langle 12 \rangle}$ is in the 0000 helicity category, since $[12]\langle 12 \rangle$ corresponds to a four-scalar massless amplitude.

To obtain the HEFT contact terms, we follow [55]. The contact terms are determined by Lorentz symmetry, which dictates their little-group transformations, locality, and the additional symmetries of the theory, in this case, $SU(3) \times U(1)_{EM}$ and baryon and lepton number. Manifestly-local contact terms can be constructed from the list of independent SCTs, namely spinor structures with no additional powers of the Mandelstams, and then appending a polynomial in the independent Mandelstam invariants, say s and t . Isolating the independent SCTs can be largely done by relying on the massless limit. However, once they are multiplied by the Mandelstams, some terms can become redundant. We refer the reader to [55] for more details. The relevant SCT bases for four-point of spins 0, 1/2, and 1 were presented in [55] and provide the basis for our analysis. Note that the SCTs carry all the polarization information of the external particles.

Since the low-energy amplitude features several mass scales, the energy growth of a certain contact term may not simply correlate with the dimension of the operator which generates it at leading order. In particular, spinor structures in longitudinal-vector helicity categories should be accompanied by an inverse factor of the vector mass in order to correctly infer the dimension at which these structures first appear at the Lagrangian level [55]. Concretely, a structure of the form $\cdots \mathbf{i} \langle \mathbf{i} \cdots$, where i is a vector line, should be normalized as $\cdots \mathbf{i} \langle \mathbf{i}/M_i \cdots$, where M_i is the mass of the vector i . This follows from the fact that $\mathbf{i} \langle \mathbf{i}/M_i$ is nothing but the polarization vector of the vector i . Another way to see this is to note that the longitudinal vector arises from a derivatively coupled Goldstone. Thus the corresponding operator contains ∂G , where G is the Goldstone field. To interpret this operator as a vector operator requires an inverse mass to get the correct dimension, that is, $\partial G \rightarrow \partial G/M_i$. Note that $\mathbf{i} \langle \mathbf{i}/M_i$ is finite in the high-energy limit for transverse vector polarizations, but scales as E/M_i for a longitudinal vector polarization.

In fact, the $1/M_V$ “poles” appearing in these contact terms (where M_V stands for a vector mass) reflect their non-local nature. These terms are required to cancel E/M_V growth in the factorizable massive amplitude in order to obtain a well-behaved theory above

v , and are therefore associated with the factorizable part of the amplitudes. In contrast, the non-factorizable parts of the amplitude consist of terms that are manifestly local, and which are therefore suppressed purely by powers of $\bar{\Lambda}$.

The distinction between $\bar{\Lambda}$ and M_V suppression is only sharp in the SMEFT, where these scales can be hierarchically separated. For the SMEFT amplitudes to be sensible at high-energy, $v \ll E \ll \Lambda$, positive powers of E/m must cancel. We list the leading power of E/m terms of the factorizable amplitudes in appendix B. The equivalence of perturbative unitarity and gauge invariance is very transparent in the massive spinor formalism. The sources of E/m behavior are factors such as $\mathbf{i}] \langle \mathbf{i}/M$. For zero vector polarization, this scales as E/M , and typically leads to amplitudes growing as a positive power of E/M . Such terms violate perturbative unitarity, which requires E^n growth to be suppressed by the same power of the cutoff. Choosing instead the vector polarization to be positive, the factor $\mathbf{i}] \mathbf{i}] / M_V$ is finite, and can be written as $\mathbf{i}] \xi_i \rangle$ where $\mathbf{i}]$ is the high-energy limit of $\mathbf{i}]^{I=1}$, and $\xi_i \rangle = \mathbf{i}]^{I=2} / M_V$ is finite. Requiring the high-energy amplitude to be independent of the arbitrary spinor $\xi_i \rangle$ is thus equivalent to requiring perturbative unitarity. On the other hand, in the massless, high-energy theory, $\xi_i \rangle$ is an arbitrary spinor, which is nothing but the reference spinor associated with the vector polarizations, and the condition that the amplitude is independent of $\xi_i \rangle$ translates to the condition that it is gauge invariant. We show one example of this type, namely the $WWhh$ amplitude in appendix A.

As mentioned above, inverse mass behavior signals the non-locality of amplitudes, associated with their factorizable parts. This is precisely the type of behavior we expect to see in the HEFT. The fact that states getting their mass from EWSB are integrated out, translates in the on-shell picture to $1/v$ non-analyticity of the amplitudes, which implies a cutoff of order v . In contrast, in the SMEFT, the full amplitudes, including factorizable and contact terms pieces, should be well behaved for $v \ll E \ll \Lambda$, with no E/M pieces.

3 Four-point contact terms at $\mathcal{O}(E^2)$

3.1 HEFT contact terms

In this section, we list the independent four-point contact terms of SM particles with E^2 energy growth, imposing $SU(3) \times U(1)_{EM}$ invariance and baryon- and lepton-number conservation. Bases of independent contact terms for four-point massive amplitudes of particles of spin ≤ 1 were derived in ref. [55]. Here we apply these results to the case of SM amplitudes. We list contact terms with E^2 growth here, and contact terms with E^4 growth in section 4. Together with the three-point electroweak amplitudes derived in ref. [33],¹ the four-point contact terms and their coefficients allow for a full parametrization of general EFT amplitudes up to E^4 .

The generic dimension-six contact terms are listed in table 1. The bolded products (\mathbf{ij}) stand for either square or angle brackets, as appropriate for the helicity category in question. The Wilson coefficients of these structures are denoted by capital C 's, with subscripts

¹Three-point gluons were not included in ref. [33], but can be obtained from the photon amplitudes by simply adding a color factor.

denoting the external particles, and superscripts denoting the helicity category. Here and in the following, f denotes any SM fermion, V denotes the W or the Z , and h denotes the physical Higgs.

Note that at this order, all the contact terms are given by spinor structures with no additional powers of the Mandelstam invariants. Thus they correspond to SCTs. In section 4, when we consider also E^4 terms, these expansions will appear. Recall that the SCTs carry the little group weights associated with the external particles and encode their polarization information. Amplitudes not shown in this table have their leading contributions from SCTs involving more than two spinor products.

Most of the contact terms in table 1 are suppressed by two powers of the cutoff, namely $1/\bar{\Lambda}^2$, and correspond to independent dimension-six operators. The exceptions are structures in longitudinal vector categories. As mentioned above, these are normalized as $\langle \mathbf{12} \rangle [\mathbf{12}] / M_V^2$ and $\langle \mathbf{13} \rangle [\mathbf{23}] / (M_V \bar{\Lambda})$ (and similarly for $1 \leftrightarrow 2$). With this normalization, we can read off the dimension of the low-energy operator which first generates these terms as 4 and 5 respectively. Indeed, $\langle \mathbf{12} \rangle [\mathbf{12}]$ is first generated at dimension-4, and corresponds to the operator $V^\mu V_\mu h^2$. It is required to cancel the high-energy growth of the massive SM factorizable amplitude.² We can split the coefficient of the contact term $\langle \mathbf{12} \rangle [\mathbf{12}]$ as $C_{WWhh}^{00} = C_{WWhh}^{00,\text{fac}} + C_{WWhh}^{00,\text{CT}}$ with the part $C_{WWhh}^{00,\text{fac}}$ canceling the E^2/M_V^2 part of the factorizable amplitude. Thus $C_{WWhh}^{00,\text{fac}}$ is determined by three-point couplings, while the remaining $C_{WWhh}^{00,\text{CT}}$ constitutes an independent Wilson coefficient. In the SMEFT, this split cleanly correlates with the counting of operator dimensions in the high-energy theory. C_{WWhh}^{00} is an expansion in v^2/Λ^2 , with the leading v^0 piece corresponding to $C_{WWhh}^{00,\text{fac}}$, and determined by the SM dimension-four gauge coupling. At dimension-six, both the three-point couplings and $C_{WWhh}^{00,\text{fac}}$ are shifted by v^2/Λ^2 corrections such that the cancellation still holds. On top of this, $C_{WWhh}^{00,\text{CT}}/\Lambda^2$ is an independent 4-point Wilson coefficient.

In the HEFT, on the other hand, the various couplings are just numbers, and there is no expansion in the VEV. Power counting can be done in various ways. Splitting C_{WWhh}^{00} as before, $C_{WWhh}^{00,\text{fac}}$ is naturally treated as dimension-four, such that upon adding it to the factorizable part, the full amplitude has no E growth. The coefficient $C_{WWhh}^{00,\text{CT}}$ can be viewed as dimension-six, since it generates E^2 terms. Alternatively, it can be viewed as dimension-four, since it corresponds to the operator $V^2 h^2$. In any case, the physical quantity is the numerical coefficient of each kinematic structure, and these differences are just a matter of theory interpretation. Moreover, there is no sharp distinction in the HEFT between the cutoff $\bar{\Lambda}$ and the electroweak mass scale v , with $\bar{\Lambda} \sim v$. In the following, when we refer to HEFT dimensions, we will refer to the dimension of the corresponding operator. The contact terms $\langle \mathbf{12} \rangle [\mathbf{12}]$ and $\langle \mathbf{13} \rangle [\mathbf{23}]$ are then dimension-4 and 5 respectively. Furthermore, it is easy to read off the minimal dimensions of these operators in the SMEFT. To leading order in the v expansion, $\bar{\Lambda}^{-2} = \Lambda^{-2}$, and $\bar{\Lambda}^{-1} = v\Lambda^{-2}$. Therefore, both of these contact terms can be first generated at dimension-six in the SMEFT. This is consistent with the fact that the factorizable fermion-fermion-vector-higgs amplitudes only feature E/M growth (see table 4), so $C_{ffVh}^{\pm\mp 0,\text{fac}} = 0$. Indeed, as was shown in [33], perturbative

²The leading high-energy behavior of each factorizable amplitude is shown in table 4.

Massive amplitudes	E^2 contact terms
$\mathcal{M}(WWhh)$	$C_{WWhh}^{00}(\mathbf{12})[\mathbf{12}], C_{WWhh}^{\pm\pm}(\mathbf{12})^2$
$\mathcal{M}(ZZhh)$	$C_{ZZhh}^{00}(\mathbf{12})[\mathbf{12}], C_{ZZhh}^{\pm\pm}(\mathbf{12})^2$
$\mathcal{M}(gghh)$	$C_{gghh}^{\pm\pm}(\mathbf{12})^2$
$\mathcal{M}(\gamma\gamma hh)$	$C_{\gamma\gamma hh}^{\pm\pm}(\mathbf{12})^2$
$\mathcal{M}(\gamma Zhh)$	$C_{\gamma Zhh}^{\pm}(\mathbf{12})^2$
$\mathcal{M}(hhhh)$	C_{hhhh}
$\mathcal{M}(f^c fhh)$	$C_{ffhh}^{\pm\pm}(\mathbf{12})$
$\mathcal{M}(f^c fWh)$	$C_{ffWh}^{+-0}[\mathbf{13}]\langle\mathbf{23}\rangle, C_{ffWh}^{-+0}(\mathbf{13})[\mathbf{23}], C_{ffWh}^{\pm\pm\pm}(\mathbf{13})(\mathbf{23})$
$\mathcal{M}(f^c fZh)$	$C_{ffZh}^{+-0}[\mathbf{13}]\langle\mathbf{23}\rangle, C_{ffZh}^{-+0}(\mathbf{13})[\mathbf{23}], C_{ffZh}^{\pm\pm\pm}(\mathbf{13})(\mathbf{23})$
$\mathcal{M}(f^c f\gamma h)$	$C_{ff\gamma h}^{\pm\pm\pm}(\mathbf{13})(\mathbf{23})$
$\mathcal{M}(q^c qgh)$	$C_{qqh}^{\pm\pm}(\mathbf{13})(\mathbf{23})$
$\mathcal{M}(f^c f f^c f)$	$C_{ffff}^{\pm\pm\pm\pm,1}(\mathbf{12})(\mathbf{34}), C_{ffff}^{--++}\langle\mathbf{12}\rangle[\mathbf{34}], C_{ffff}^{-++-}\langle\mathbf{13}\rangle[\mathbf{24}], C_{ffff}^{-++-}\langle\mathbf{14}\rangle[\mathbf{23}]$ $C_{ffff}^{\pm\pm\pm\pm,2}(\mathbf{13})(\mathbf{24}), C_{ffff}^{++--}[\mathbf{12}]\langle\mathbf{34}\rangle, C_{ffff}^{+-+-}[\mathbf{13}]\langle\mathbf{24}\rangle, C_{ffff}^{+-+-}[\mathbf{14}]\langle\mathbf{23}\rangle$

Table 1. Contact terms with E^2 growth. The C 's stand for independent HEFT coefficients, and are mostly generated at $\bar{\Lambda}^{-2}$, corresponding to $d = 6$ operators. The only exceptions are C_{WWhh}^{00} and $C_{ffVh}^{\pm\mp 0}$ which appear with M_V^{-2} and $(M_V \bar{\Lambda})^{-1}$ respectively, corresponding to $d = 4$ and $d = 5$ operators (for details see text). Color structures and indices are not shown but can be added unambiguously. For identical Majorana neutrinos, the structures $C_{ffZh}^{\pm\pm\pm}(\mathbf{13})(\mathbf{23})$ and $C_{ff\gamma h}^{\pm\pm\pm}(\mathbf{13})(\mathbf{23})$ do not appear.

unitarity of this amplitude only implies relations between SM couplings, specifically, the relation between the fermion mass, the Yukawa coupling, and the Higgs VEV.

3.2 SMEFT contact terms

To obtain the SMEFT contact terms, we start with the massless dimension-six SMEFT contact terms. These were derived in [34] and we list them for completeness in table 2.

For each amplitude in table 2, we show the kinematic and group theory structure. We also list the Warsaw basis operator, or combination of operators, \mathcal{O} , that generates this structure, and the corresponding Wilson coefficient c . We use H to denote the Higgs doublet, g , W and B for an SU(3), SU(2) or U(1) gauge boson respectively, Q (L) for SU(2)-doublet quarks (leptons), and u , d (e) for SU(2)-singlet quarks (leptons). The different group theory factors are denoted as follows: σ^I are the Pauli matrices, λ^A are the Gell-Mann matrices, $T_{ij}^{\pm kl} \equiv 1/2(\delta_i^k \delta_j^l \pm \delta_j^k \delta_i^l)$, $T_{ijk}^{lmn} \equiv \delta_i^l \delta_j^m \delta_k^n + \delta_i^l \delta_k^m \delta_j^n + \delta_j^l \delta_i^m \delta_k^n + \delta_j^l \delta_k^m \delta_i^n + \delta_k^l \delta_j^m \delta_i^n + \delta_k^l \delta_i^m \delta_j^n$, and ϵ^{IJK} and f^{ABC} are the SU(2)_L and SU(3)_c structure constants respectively.

Parameterizing the Higgs doublet as

$$H = \left(G^+, \frac{1}{\sqrt{2}}(v + h + iG^0) \right)^T, \quad (3.1)$$

Amplitude	Contact term	Warsaw basis operator	Coefficient
$\mathcal{A}(H_i^c H_j^c H_k^c H^l H^m H^n)$	T_{ijk}^{+lmn}	$\mathcal{O}_H/6$	$c_{(H^\dagger H)^3}$
$\mathcal{A}(H_i^c H_j^c H^k H^l)$	$s_{12} T_{ij}^{+kl}$	$\mathcal{O}_{HD}/2 + \mathcal{O}_{H\Box}/4$	$c_{(H^\dagger H)^2}^{(+)}$
$\mathcal{A}(H_i^c H_j^c H^k H^l)$	$(s_{13} - s_{23}) T_{ij}^{-kl}$	$\mathcal{O}_{HD}/2 - \mathcal{O}_{H\Box}/4$	$c_{(H^\dagger H)^2}^{(-)}$
$\mathcal{A}(B^\pm B^\pm H_i^c H^j)$	$(12)^2 \delta_i^j$	$(\mathcal{O}_{HB} \pm i\mathcal{O}_{H\tilde{B}})/2$	$c_{BBHH}^{\pm\pm}$
$\mathcal{A}(B^\pm W^{I\pm} H_i^c H^j)$	$(12)^2 (\sigma^I)_i^j$	$\mathcal{O}_{HWB} \pm i\mathcal{O}_{H\tilde{W}B}$	$c_{BWHH}^{\pm\pm}$
$\mathcal{A}(W^{I+} W^{J+} H_i^c H^j)$	$(12)^2 \delta^{IJ} \delta_i^j$	$(\mathcal{O}_{HW} \pm i\mathcal{O}_{H\tilde{W}})/2$	$c_{WWHH}^{\pm\pm}$
$\mathcal{A}(g^{A\pm} g^{B\pm} H_i^c H^j)$	$(12)^2 \delta^{AB} \delta_i^j$	$(\mathcal{O}_{HG} \pm i\mathcal{O}_{H\tilde{G}})/2$	$c_{GGHH}^{\pm\pm}$
$\mathcal{A}(L_i^c e H_j^c H^k H^l)$	$[12] T_{ij}^{+kl}$	$\mathcal{O}_{eH}/2$	c_{LeHHH}^{++}
$\mathcal{A}(Q_{a,i}^c d^b H_j^c H^k H^l)$	$[12] T_{ij}^{+kl} \delta_a^b$	$\mathcal{O}_{dH}/2$	c_{QdHHH}^{++}
$\mathcal{A}(Q_{a,i}^c u^b H_j^c H^k H^l)$	$[12] \varepsilon_{im} T_{jk}^{+ml} \delta_a^b$	$\mathcal{O}_{uH}/2$	c_{QuHHH}^{++}
$\mathcal{A}(e^c e H_i^c H^j)$	$\langle 142 \rangle \delta_i^j$	$\mathcal{O}_{He}/2$	c_{eeHH}^{-+}
$\mathcal{A}(u_a^c u^b H_i^c H^j)$	$\langle 142 \rangle \delta_i^j \delta_a^b$	$\mathcal{O}_{Hu}/2$	c_{uuHH}^{-+}
$\mathcal{A}(d_a^c d^b H_i^c H^j)$	$\langle 142 \rangle \delta_i^j \delta_a^b$	$\mathcal{O}_{Hd}/2$	c_{ddHH}^{-+}
$\mathcal{A}(u_a^c d^b H^i H^j)$	$\langle 142 \rangle \epsilon^{ij} \delta_a^b$	$\mathcal{O}_{Hud}/2$	c_{udHH}^{-+}
$\mathcal{A}(L_i^c L^j H_k^c H^l)$	$[142] T_{ik}^{+jl}$	$(\mathcal{O}_{HL}^{(1)} + \mathcal{O}_{HL}^{(3)})/8$	$c_{LLHH}^{+,-,+}$
$\mathcal{A}(L_i^c L^j H_k^c H^l)$	$[142] T_{ik}^{-jl}$	$(\mathcal{O}_{HL}^{(1)} - \mathcal{O}_{HL}^{(3)})/8$	$c_{LLHH}^{+,-,-}$
$\mathcal{A}(Q_{a,i}^c Q^{bj} H_k^c H^l)$	$[142] T_{ik}^{+jl} \delta_a^b$	$(3\mathcal{O}_{HQ}^{(1)} + \mathcal{O}_{HQ}^{(3)})/8$	$c_{QQHH}^{+,-,+}$
$\mathcal{A}(Q_{a,i}^c Q^{bj} H_k^c H^l)$	$[142] T_{ik}^{-jl} \delta_a^b$	$(\mathcal{O}_{HQ}^{(1)} - \mathcal{O}_{HQ}^{(3)})/8$	$c_{QQHH}^{+,-,-}$
$\mathcal{A}(L_i^c e B^+ H^j)$	$[13][23] \delta_i^j$	$-i\mathcal{O}_{eB}/(2\sqrt{2})$	c_{LeBH}^{+++}
$\mathcal{A}(Q_{a,i}^c d^b B^+ H^j)$	$[13][23] \delta_i^j \delta_a^b$	$-i\mathcal{O}_{dB}/(2\sqrt{2})$	c_{QdBH}^{+++}
$\mathcal{A}(Q_{a,i}^c u^b B^+ H^j)$	$[13][23] \epsilon_{ij} \delta_a^b$	$-i\mathcal{O}_{uB}/(2\sqrt{2})$	c_{QuBH}^{+++}
$\mathcal{A}(L_i^c e W^{I+} H^j)$	$[13][23] (\sigma^I)_i^j$	$-i\mathcal{O}_{eW}/(2\sqrt{2})$	c_{LeWH}^{+++}
$\mathcal{A}(Q_{a,i}^c d^b W^{I+} H^j)$	$[13][23] (\sigma^I)_i^j \delta_a^b$	$-i\mathcal{O}_{dW}/(2\sqrt{2})$	c_{QdWH}^{+++}
$\mathcal{A}(Q_{a,i}^c u^b W^{I+} H^j)$	$[13][23] (\sigma^I)_{ik} \epsilon_j^k \delta_a^b$	$-i\mathcal{O}_{uW}/(2\sqrt{2})$	c_{QuWH}^{+++}
$\mathcal{A}(Q_{a,i}^c d^b g^{A+} H^j)$	$[13][23] \delta_i^j (\lambda^A)_a^b$	$-i\mathcal{O}_{dG}/(2\sqrt{2})$	c_{QdGH}^{+++}
$\mathcal{A}(Q_{a,i}^c u^b g^{A+} H^j)$	$[13][23] \epsilon_{ij} (\lambda^A)_a^b$	$-i\mathcal{O}_{uG}/(2\sqrt{2})$	c_{QuGH}^{+++}
$\mathcal{A}(W^{I\pm} W^{J\pm} W^{K\pm})$	$(12)(23)(31) \epsilon^{IJK}$	$(\mathcal{O}_W \pm i\mathcal{O}_{\tilde{W}})/6$	$c_{WWW}^{\pm\pm\pm}$
$\mathcal{A}(g^{A\pm} g^{B\pm} g^{C\pm})$	$(12)(23)(31) f^{ABC}$	$(\mathcal{O}_G \pm i\mathcal{O}_{\tilde{G}})/6$	$c_{GGG}^{\pm\pm\pm}$

Table 2. Massless $d = 6$ SMEFT contact terms [34] and their relations to Warsaw basis operators [3]. For each operator (or operator combination) \mathcal{O} in the third column, $c\mathcal{O}$ generates the structure in the second column with the coefficient c given in the fourth column. c -superscripts denote charge conjugation.

we can obtain the high-energy amplitudes featuring the Goldstones G^\pm and the radial mode h on the external legs. Each one of the massless contact terms is then ‘‘Higgsed’’ to obtain the corresponding massive contact term(s), as described in ref. [47]. Massless contact terms featuring only fermions and vectors are simply bolded to give massive contact terms with fermions and vectors, in transverse vector helicity categories. Massless contact terms featuring a Higgs leg give rise to contact terms with a massive scalar leg, in which case they are simply bolded; or to contact terms with a massive vector leg. Thus for example, based on kinematics alone, it is easy to see that at order E^2 , the $Q^\dagger Q H^\dagger H$ contact term gives rise to a $Q^\dagger Q Z h$ contact term, but does not generate a contact term with two physical Higgses. The massless amplitude features [132]. We can then read off the massive structure using,

$$[132] = [13]\langle 32 \rangle \rightarrow [\mathbf{13}]\langle \mathbf{32} \rangle, \tag{3.2}$$

which contributes to the $Q^\dagger Q Z h$ amplitude. Note that only a structure with a momentum insertion p_3 can give rise to a vector amplitude. Indeed [132] is consistent with being a Goldstone amplitude since it is derivatively coupled. On the other hand, [132] cannot contribute to a low-energy amplitude with two physical Higgses: Bose symmetry would require $[132] \rightarrow [1(3+4)2]$ which is vanishing.

This procedure reproduces the full set of structures of table 1, and relates their coefficients to the massless SMEFT coefficients. We collect the massive SMEFT contact terms and their coefficients in table 3. Four-fermion contact terms are not shown here because their matching to the high-energy amplitudes is straightforward. Each of the Wilson coefficients C in table 3 is $d = 6$, and is suppressed by Λ^2 . As explained in section 3.1, the low-energy amplitudes may also contain mass-suppressed contact terms in longitudinal vector helicity categories, which are associated with the factorizable part of the amplitude. Thus for example, the structure $\langle \mathbf{12} \rangle [\mathbf{12}]$ in the $WWhh$ amplitude has two pieces: one comes with a coefficient $C_{WWhh}^{00,\text{fac}}$, which is determined by three-point couplings, and one which is an independent SMEFT $d = 6$ four-point coupling, $C_{WWhh}^{00,\text{CT}}$. Only the latter is given in table 3, but we omit the superscript CT for simplicity.

Note furthermore that high-energy four-point contact terms with Higgs legs may also correct the three-point couplings. The $d = 6$ SMEFT corrections to the three-points were derived in ref. [33] by matching to the Feynman diagram result obtained using ref. [65]. These corrections can also be obtained by on-shell Higgsing. For an explicit example, see appendix A, where we calculate the v^2/Λ^2 correction to the WWh coupling from the massless $H^2(H^\dagger)^2 WW$ amplitude.

For the $d = 6$ bosonic contact terms of table 3, the only change compared to the HEFT contact terms of table 1 is in the $\pm\pm$ helicity categories of $VVhh$, where six $d = 6$ SMEFT parameters control eight HEFT parameters. Additional relations appear among the fermion SMEFT amplitudes, where the coefficients of up- and down-quark (or antiquark) amplitudes featuring i , (or \bar{i}) are equal, since they originate from the same doublet (anti)-quark amplitude. The coefficients of lepton-doublet amplitudes are similarly related.

Massive $d = 6$ amplitudes	SMEFT Wilson coefficients
$\mathcal{M}(W_L^+ W_L^- hh) = C_{WWhh}^{00} \langle \mathbf{12} \rangle [\mathbf{12}]$	$C_{WWhh}^{00} = (c_{(H^\dagger H)^2}^{(+)} - 3c_{(H^\dagger H)^2}^{(-)})/2$
$\mathcal{M}(W_\pm^+ W_\pm^- hh) = C_{WWhh}^{\pm\pm} (\mathbf{12})^2$	$C_{WWhh}^{\pm\pm} = 2c_{WWHH}^{\pm\pm}$
$\mathcal{M}(Z_L Z_L hh) = C_{ZZhh}^{00} \langle \mathbf{12} \rangle [\mathbf{12}]$	$C_{ZZhh}^{00} = -2c_{(H^\dagger H)^2}^{(+)}$
$\mathcal{M}(Z_\pm Z_\pm hh) = C_{ZZhh}^{\pm\pm} (\mathbf{12})^2$	$C_{ZZhh}^{\pm\pm} = c_W^2 c_{WWHH}^{\pm\pm} + s_W^2 c_{BBHH}^{\pm\pm} + c_W s_W c_{BWHH}^{\pm\pm}$
$\mathcal{M}(g_\pm g_\pm hh) = C_{gghh}^{\pm\pm} (\mathbf{12})^2$	$C_{gghh}^{\pm\pm} = c_{GGHH}^{\pm\pm}$
$\mathcal{M}(\gamma_\pm \gamma_\pm hh) = C_{\gamma\gamma hh}^{\pm\pm} (\mathbf{12})^2$	$C_{\gamma\gamma hh}^{\pm\pm} = s_W^2 c_{WWHH}^{\pm\pm} + c_W^2 c_{BBHH}^{\pm\pm} - c_W s_W c_{BWHH}^{\pm\pm}$
$\mathcal{M}(\gamma_\pm Z hh) = C_{\gamma Zh h}^\pm (\mathbf{12})^2$	$C_{\gamma Zh h}^\pm = s_W c_W c_{WWHH}^{\pm\pm} - s_W c_W c_{BBHH}^{\pm\pm} + \frac{1}{2}(s_W^2 - c_W^2) c_{BWHH}^{\pm\pm}$
$\mathcal{M}(hhhh) = C_{hhhh}$	$C_{hhhh} = -3c_{(H^\dagger H)^2} + 45 v^2 c_{(H^\dagger H)^3}$
$\mathcal{M}(f_\pm^c f_\pm hh) = C_{ffhh}^{\pm\pm} (\mathbf{12})$	$C_{ffhh}^{\pm\pm} = 3c_{\Psi\Psi HHH}^{\pm\pm} v / (2\sqrt{2})$
$\mathcal{M}(f_+^c f_- W_L h) = C_{ffWh}^{+-0} [\mathbf{13}] [\mathbf{23}]$	$C_{ffWh}^{+-0} = (c_{\Psi\Psi HH}^{+-, (+)} - c_{\Psi\Psi HH}^{+-, (-)})/2$
$\mathcal{M}(f_-^c f_+ W_L h) = C_{ffWh}^{-+0} [\mathbf{13}] [\mathbf{23}]$	$C_{ffWh}^{-+0} = c_{\psi_R \psi_R' HH}^-$
$\mathcal{M}(f_\pm^c f_\pm^\dagger W_\pm h) = C_{ffWh}^{\pm\pm\pm} (\mathbf{13}) (\mathbf{23})$	$C_{ffWh}^{\pm\pm\pm} = c_{\Psi\Psi WH}^{\pm\pm\pm} / 2$
$\mathcal{M}(f_+^c f_- Z_L h) = C_{ffZh}^{+-0} [\mathbf{13}] \langle \mathbf{23} \rangle$	$C_{e_L e_L Zh}^{+-0} = -i\sqrt{2} c_{\Psi\Psi HH}^{+-, (+)}, C_{\nu_L \nu_L Zh}^{+-0} = -i(c_{\Psi\Psi HH}^{+-, (+)} + c_{\Psi\Psi HH}^{+-, (-)})/\sqrt{2}$
$\mathcal{M}(f_-^c f_+ Z_L h) = C_{ffZh}^{-+0} \langle \mathbf{13} \rangle [\mathbf{23}]$	$C_{ffZh}^{-+0, CT} = -i\sqrt{2} c_{\psi\psi HH}^-$
$\mathcal{M}(f_\pm^c f_\pm Z_\pm h) = C_{ffZh}^{\pm\pm\pm} (\mathbf{13}) (\mathbf{23})$	$C_{ffZh}^{\pm\pm\pm} = -(s_W c_{\Psi\psi BH}^{\pm\pm\pm} + c_W c_{\Psi\psi WH}^{\pm\pm\pm})/\sqrt{2}$
$\mathcal{M}(f_\pm^c f_\pm \gamma_\pm h) = C_{ff\gamma h}^{\pm\pm\pm} (\mathbf{13}) (\mathbf{23})$	$C_{ff\gamma h}^{\pm\pm\pm} = (-s_W c_{\Psi\psi WH}^{\pm\pm\pm} + c_W c_{\Psi\psi BH}^{\pm\pm\pm})/\sqrt{2}$
$\mathcal{M}(q_\pm^c q_\pm g_\pm^A h) = C_{qqgh}^{\pm\pm\pm} \lambda^A (\mathbf{13}) (\mathbf{23})$	$C_{qqgh}^{\pm\pm\pm} = c_{\Psi\psi GH}^{\pm\pm\pm} / \sqrt{2}$

Table 3. The low-energy E^2 contact terms (left column) and their $d = 6$ coefficients in the SMEFT (right column). $c_{(H^\dagger H)^2}$ without a superscript is the renormalizable four-Higgs coupling. The mapping for four fermion contact terms is trivial, so we do not include them here.

4 Four-point contact terms at E^3 and E^4

In this section, we derive the remaining contact terms contributing to the SM amplitudes up to and including quartic energy growth. These include additional SCTs beyond those listed in table 2, as well as variations of the SCTs in table 2 multiplied by powers of the Mandelstam invariants. For generic four-point amplitudes with spins ≤ 1 , the list of independent SCTs is exhausted at quartic energy growth. However, for the SM particle content, some of these only contribute at higher orders, when multiplied by additional powers of the invariants, due to (anti)symmetrization over identical particles. We comment on these additional contributions where relevant.

For each amplitude, we show the independent contact terms, and the dimension of the corresponding HEFT operator, following the discussion in section 3.1. Recall that apart from longitudinal vector categories, all structures are suppressed by the appropriate power of $\bar{\Lambda}$, namely $\bar{\Lambda}^3$ or $\bar{\Lambda}^4$ here. On the other hand, each longitudinal vector i comes with a factor $i) [i/M_i$. In the HEFT, $\bar{\Lambda} = v \sim M_V$, but the M_V factors allow us to infer the dimension of

the corresponding operator. We also show the lowest dimension at which each structure may be generated in the SMEFT, using the fact that any single power of $\bar{\Lambda}$ can be written as $1/\bar{\Lambda} = v/\Lambda^2$. Thus for example, in the $WWZh$ amplitude, the structure $[\mathbf{13}][\mathbf{12}]\langle\mathbf{23}\rangle$ is accompanied by $1/(M_W M_Z \bar{\Lambda})$, and its HEFT and possible SMEFT dimensions are given as (5, 8).

Where appropriate, we only show “half” the allowed structures, with the rest obtained by a parity flip (PF), switching all angle and square brackets. The number of independent structures is also given, following the HEFT and SMEFT operator dimensions. In the HEFT, the coefficients of the terms listed here are all independent. In the SMEFT, many of them are related. These relations can be derived by “Higgsing” the massless amplitudes. This was done for $\bar{u}dWh$ and $WWhh$ in ref. [47]. We also comment on how the contact terms are modified when Majorana neutrinos are involved.

4.1 Bosonic amplitudes with all massive particles

4.1.1 $hhhh$

There is no E^2 contact term due to the Bose symmetry of the Higgs legs. The first contact term appears at E^4 and is,

$$\tilde{s}_{12}^2 + \tilde{s}_{13}^2 + \tilde{s}_{14}^2 \quad (8, 8) \quad \# = 1 \quad (4.1)$$

Here and in the following, the numbers in the parenthesis indicate the dimensions of the corresponding HEFT and SMEFT operators respectively, and $\#$ is the number of independent contact terms.

4.1.2 $Zhhh$

Once we symmetrize over h legs, there is no E^2 contact term. At E^4 there is a single structure,

$$0 : \quad \tilde{s}_{12}[\mathbf{121}] + \tilde{s}_{13}[\mathbf{131}] + \tilde{s}_{14}[\mathbf{141}] \quad (7; 8) \quad \# = 1 \quad (4.2)$$

The Mandelstams are necessary due to symmetrization over h . The symmetric sum of $\tilde{s}_{13}[\mathbf{121}]$ is $(\tilde{s}_{13} + \tilde{s}_{14})[\mathbf{121}] + (\tilde{s}_{12} + \tilde{s}_{14})[\mathbf{131}] + (\tilde{s}_{13} + \tilde{s}_{14})[\mathbf{141}]$ which simplifies to the above structure. Note that there is no LE factorizable amplitude.

There is an additional SCT in this case, which first contributes at dimension 13, $(\tilde{s}_{12} - \tilde{s}_{13})(\tilde{s}_{12} - \tilde{s}_{14})(\tilde{s}_{13} - \tilde{s}_{14})([\mathbf{1231}] - \langle\mathbf{1231}\rangle)$.

4.1.3 $ZZhh$

$$\begin{aligned} 00 : & \quad [\mathbf{131}][\mathbf{232}] + [\mathbf{141}][\mathbf{242}], \tilde{s}_{12}[\mathbf{12}]\langle\mathbf{12}\rangle \quad (6; 8) \quad \# = 2 \\ ++ : & \quad \tilde{s}_{12}[\mathbf{12}]^2; \text{ PF} \quad (8; 8) \quad \# = 2 \\ +- : & \quad [\mathbf{1(3-4)2}]^2 + \langle\mathbf{1(3-4)2}\rangle^2 \quad (8; 8) \quad \# = 1 \end{aligned} \quad (4.3)$$

Since there is no $E^4/(M^2\bar{\Lambda}^2)$ growth in the factorizable amplitude, there are no $M^2\Lambda^2$ -suppressed contact terms in the SMEFT. All independent $vvss$ SCTs appear at E^4 order.

4.1.4 W^+W^-hh

$$\begin{aligned}
 00 : & \quad [\mathbf{131}][\mathbf{242}] + [\mathbf{141}][\mathbf{232}], \tilde{s}_{12}[\mathbf{12}]\langle\mathbf{12}\rangle & (6;8) \# = 2 \\
 ++ : & \quad \tilde{s}_{12}[\mathbf{12}]^2; \text{PF} & (8;8) \# = 2 \\
 +- : & \quad [\mathbf{1(3-4)2}]^2; \text{PF} & (8;8) \# = 2
 \end{aligned} \tag{4.4}$$

All independent $vvss$ SCTs appear at order E^4 .

4.1.5 W^+W^-Zh

$$\begin{aligned}
 000 : & \quad [\mathbf{12}][\mathbf{343}]\langle\mathbf{12}\rangle, (1 \leftrightarrow 3), (2 \leftrightarrow 3) & (5;8) \# = 3 \\
 +00 : & \quad [\mathbf{12}]\langle\mathbf{23}\rangle[\mathbf{31}]; \text{Perm}(+00); \text{PF} & (5;8) \# = 6 \\
 ++0 : & \quad \{[\mathbf{12}]^2[\mathbf{313}], [\mathbf{12}]^2[\mathbf{323}]\}; \text{Perm}(++0); \text{PF} & (7;8) \# = 12 \\
 +-0 : & \quad [\mathbf{13}][\mathbf{142}]\langle\mathbf{23}\rangle, \text{Perm}(+-0) & (7;8) \# = 6 \\
 +++ : & \quad [\mathbf{12}][\mathbf{13}][\mathbf{23}]; \text{PF} & (7,8) \# = 2
 \end{aligned} \tag{4.5}$$

Above, ‘‘Perm’’ stands for the different possible helicity assignments, eg, $(+00)$, $(0+0)$, $(00+)$. For the $(+-0)$ helicity category, two of the six structures can be exchanged for other $\mathcal{O}(E^4)$ SCTs times Mandelstams. Since the latter are beyond quartic order and therefore not included in our counting, all six structures $(+-0)$ are independent.

4.1.6 $ZZZh$

$$\begin{aligned}
 000 : & \quad [\mathbf{12}][\mathbf{343}]\langle\mathbf{12}\rangle + \text{Perm}(123) & (5;8) \# = 1 \\
 ++0 : & \quad [\mathbf{12}]^2[\mathbf{343}] + \text{Perm}(++0); \text{PF} & (7;8) \# = 2 \\
 +-0 : & \quad [\mathbf{13}][\mathbf{142}]\langle\mathbf{23}\rangle + \text{Perm}(+-0) & (7;8) \# = 1
 \end{aligned} \tag{4.6}$$

Here, $\text{Perm}(123)$ means all permutations of the momenta. The remaining SCTs which appear in $WWZh$ require additional Mandelstams to satisfy the Bose symmetry of the Z bosons. The $(+00)$ helicity category first appears at E^5 as $(s_{12} - s_{13})[\mathbf{12}]\langle\mathbf{23}\rangle[\mathbf{31}]$. With the parity flipped structure, this introduces two independent coefficients. The $(+++)$ helicity category first appears at E^9 from $(s_{12} - s_{13})(s_{13} - s_{23})(s_{21} - s_{23})[\mathbf{12}][\mathbf{13}][\mathbf{23}]$, with an additional independent structure from parity.

4.1.7 W^+W^-ZZ

$$\begin{aligned}
 0000 : & \quad [\mathbf{12}][\mathbf{34}]\langle\mathbf{12}\rangle\langle\mathbf{34}\rangle, [\mathbf{13}][\mathbf{24}]\langle\mathbf{13}\rangle\langle\mathbf{24}\rangle + (3 \leftrightarrow 4) & (4;8) \# = 2 \\
 ++00 : & \quad [\mathbf{12}]^2[\mathbf{34}]\langle\mathbf{34}\rangle; \text{PF} & (6;8) \# = 2 \\
 +0+0 : & \quad \{[\mathbf{12}][\mathbf{34}][\mathbf{13}]\langle\mathbf{24}\rangle, [\mathbf{14}][\mathbf{23}][\mathbf{13}]\langle\mathbf{24}\rangle\} + (3 \leftrightarrow 4); (1 \leftrightarrow 2); \text{PF} & (6;8) \# = 8 \\
 00++ : & \quad [\mathbf{34}]^2[\mathbf{12}]\langle\mathbf{12}\rangle; \text{PF} & (6;8) \# = 2 \\
 +-00 : & \quad [\mathbf{13}][\mathbf{14}]\langle\mathbf{23}\rangle\langle\mathbf{24}\rangle; \text{PF} & (6;8) \# = 2 \\
 +0-0 : & \quad \{[\mathbf{12}][\mathbf{14}]\langle\mathbf{23}\rangle\langle\mathbf{34}\rangle + (3 \leftrightarrow 4), (1 \leftrightarrow 2)\}; \text{PF} & (6;8) \# = 4 \\
 00+- : & \quad [\mathbf{13}][\mathbf{23}]\langle\mathbf{14}\rangle\langle\mathbf{24}\rangle + (3 \leftrightarrow 4) & (6;8) \# = 1 \\
 ++++ : & \quad \{[\mathbf{12}]^2[\mathbf{34}]^2, [\mathbf{13}]^2[\mathbf{24}]^2 + (3 \leftrightarrow 4)\}; \text{PF} & (8;8) \# = 4 \\
 ++-- : & \quad [\mathbf{12}]^2\langle\mathbf{34}\rangle^2; \text{PF} & (8;8) \# = 2 \\
 -+-+ : & \quad [\mathbf{14}]^2\langle\mathbf{23}\rangle^2 + (3 \leftrightarrow 4); \text{PF} & (8;8) \# = 2
 \end{aligned} \tag{4.7}$$

At order E^5 several new $vvvv$ SCTs become independent in the $(+000)$, $(+++0)$, and $(++-0)$ helicity categories.

4.1.8 $W^+W^+W^-W^-$

$$\begin{aligned}
 0000 : & \quad [\mathbf{12}][\mathbf{34}]\langle\mathbf{12}\rangle\langle\mathbf{34}\rangle, [\mathbf{13}][\mathbf{24}]\langle\mathbf{13}\rangle\langle\mathbf{24}\rangle + (3 \leftrightarrow 4) & (4;8) \# = 2 \\
 ++00 : & \quad [\mathbf{12}]^2[\mathbf{34}]\langle\mathbf{34}\rangle; \text{PF} & (6;8) \# = 2 \\
 +0+0 : & \quad \{[\mathbf{12}][\mathbf{34}][\mathbf{13}]\langle\mathbf{24}\rangle, [\mathbf{14}][\mathbf{23}][\mathbf{13}]\langle\mathbf{24}\rangle\} + (1 \leftrightarrow 2) + (3 \leftrightarrow 4); \text{PF} & (6;8) \# = 4 \\
 00++ : & \quad [\mathbf{34}]^2[\mathbf{12}]\langle\mathbf{12}\rangle; \text{PF} & (6;8) \# = 2 \\
 +-00 : & \quad [\mathbf{13}][\mathbf{14}]\langle\mathbf{23}\rangle\langle\mathbf{24}\rangle + (1 \leftrightarrow 2) & (6;8) \# = 1 \\
 +0-0 : & \quad [\mathbf{12}][\mathbf{14}]\langle\mathbf{23}\rangle\langle\mathbf{34}\rangle + (1 \leftrightarrow 2) + (3 \leftrightarrow 4); \text{PF} & (6;8) \# = 2 \\
 00+- : & \quad [\mathbf{13}][\mathbf{23}]\langle\mathbf{14}\rangle\langle\mathbf{24}\rangle + (3 \leftrightarrow 4) & (6;8) \# = 1 \\
 ++++ : & \quad \{[\mathbf{13}]^2[\mathbf{24}]^2 + (1 \leftrightarrow 2), [\mathbf{13}][\mathbf{14}][\mathbf{23}][\mathbf{24}]\}; \text{PF} & (8;8) \# = 4 \\
 ++-- : & \quad [\mathbf{12}]^2\langle\mathbf{34}\rangle^2; \text{PF} & (8;8) \# = 2 \\
 -+ -+ : & \quad [\mathbf{24}]^2\langle\mathbf{13}\rangle^2 + [\mathbf{14}]^2\langle\mathbf{23}\rangle^2 + (3 \leftrightarrow 4) & (8;8) \# = 1
 \end{aligned} \tag{4.8}$$

At E^5 several new $vvvv$ SCTs become independent in the $(+000)$, $(++ +0)$, and $(++ -0)$ helicity categories.

4.1.9 $ZZZZ$

$$\begin{aligned}
 0000 : & \quad [\mathbf{13}][\mathbf{24}]\langle\mathbf{13}\rangle\langle\mathbf{24}\rangle + \text{Perm}(1234) & (4;8) \# = 1 \\
 ++00 : & \quad [\mathbf{12}]^2[\mathbf{34}]\langle\mathbf{34}\rangle + \text{Perm}(1234); \text{PF} & (6;8) \# = 2 \\
 +-00 : & \quad [\mathbf{13}][\mathbf{14}]\langle\mathbf{23}\rangle\langle\mathbf{24}\rangle + \text{Perm}(1234) & (6;8) \# = 1 \\
 ++++ : & \quad [\mathbf{12}]^2[\mathbf{34}]^2 + [\mathbf{13}]^2[\mathbf{24}]^2 + [\mathbf{14}]^2[\mathbf{23}]^2; \text{PF} & (8;8) \# = 2 \\
 ++-- : & \quad [\mathbf{12}]^2\langle\mathbf{34}\rangle^2 + \text{Perm}(1234) & (8;8) \# = 1
 \end{aligned} \tag{4.9}$$

At E^5 several new $vvvv$ SCTs become independent in the $(+000)$, $(++ +0)$, and $(++ -0)$ helicity categories.

4.2 Fermionic amplitudes with all massive particles

4.2.1 $f^c fhh$

$$\begin{aligned}
 ++ : & \quad \tilde{s}_{12}[\mathbf{12}]; \text{PF} & (7;8) \# = 2 \\
 +- : & \quad \{\tilde{s}_{14}[\mathbf{132}] + \tilde{s}_{13}[\mathbf{142}]\}; \text{PF} & (8;8) \# = 2
 \end{aligned} \tag{4.10}$$

All SCT bases are covered at E^4 . For Majorana neutrinos, there is only a single independent coefficient in the $(+-)$ category.

4.2.2 $f^c fZh$ and $f^c f'Wh$

$$\begin{aligned}
 ++0 : & \quad \{[\mathbf{12}][\mathbf{313}], [\mathbf{12}][\mathbf{323}]\}; \text{PF} & (6;8) \# = 4 \\
 +-+ : & \quad \{[\mathbf{13}][\mathbf{312}], [\mathbf{23}][\mathbf{321}]\}; \text{PF} & (7;8) \# = 4 \\
 +-0 : & \quad [\mathbf{13}]\langle\mathbf{23}\rangle \times \{\tilde{s}_{12}, \tilde{s}_{13}\}; \text{PF} & (7;8) \# = 4 \\
 +++ : & \quad [\mathbf{13}][\mathbf{23}] \times \{\tilde{s}_{12}, \tilde{s}_{13}\}; \text{PF} & (8;8) \# = 4 \\
 +-+ : & \quad [\mathbf{12}]\langle\mathbf{3123}\rangle; \text{PF} & (8;8) \# = 2
 \end{aligned} \tag{4.11}$$

All SCT bases are covered at E^4 .

For identical Majorana neutrinos, the $\nu\nu Zh$ structures are modified to,

$$\begin{aligned}
 ++0 : & \quad [\mathbf{12}][\mathbf{313}]; \text{ PF} & (6;8) \# = 2 \\
 +-+ : & \quad [\mathbf{13}][\mathbf{312}]; \text{ PF} & (7;8) \# = 2 \\
 +-0 : & \quad ([\mathbf{13}]\langle\mathbf{23}\rangle - (1 \leftrightarrow 2)) \times \tilde{s}_{12}, ([\mathbf{13}]\langle\mathbf{23}\rangle + (1 \leftrightarrow 2)) \times (\tilde{s}_{13} - \tilde{s}_{23}) & (7;8) \# = 2 \\
 +++ : & \quad [\mathbf{13}][\mathbf{23}] \times (\tilde{s}_{13} - \tilde{s}_{23}); \text{ PF} & (8;8) \# = 2
 \end{aligned} \tag{4.12}$$

where we only show the HEFT operator dimensions. The $(++-)$ helicity category only appears at E^6 , with the two independent E^4 structures multiplied by $s_{13} - s_{14}$.

4.2.3 $f^c f^c f f$

When the four fermions are distinguishable, the contact terms are,

$$\begin{aligned}
 +++- : & \quad \{[\mathbf{12}][\mathbf{324}], \text{Perm}(+++-)\}; \text{ PF} & (7;8) \# = 8 \\
 ++++ : & \quad \{\tilde{s}_{13}[\mathbf{13}][\mathbf{24}], \tilde{s}_{13}[\mathbf{14}][\mathbf{23}], \tilde{s}_{14}[\mathbf{14}][\mathbf{23}]\}; \text{ PF} & (8;8) \# = 6 \\
 +++- : & \quad \{[\mathbf{12}]\langle\mathbf{34}\rangle, \text{Perm}(+++ -)\} \times \{\tilde{s}_{12}, \tilde{s}_{13}\} & (8;8) \# = 12
 \end{aligned} \tag{4.13}$$

All SCTs are covered at E^4 . For four Dirac fermions of the same flavor, $f_1^c f_1^c f_1 f_1$, the basis is to modified to,

$$\begin{aligned}
 ++++ : & \quad \{[\mathbf{12}][\mathbf{34}] \times \tilde{s}_{12}, ([\mathbf{13}][\mathbf{24}] + (1 \leftrightarrow 2)) \times (\tilde{s}_{13} - \tilde{s}_{14})\}; \text{ PF} & (8;8) \# = 4 \\
 +++- : & \quad [\mathbf{12}]\langle\mathbf{34}\rangle \times \tilde{s}_{12}; \text{ PF} & (8;8) \# = 2 \\
 +-+- : & \quad [([\mathbf{13}]\langle\mathbf{24}\rangle - (3 \leftrightarrow 4)) - (1 \leftrightarrow 2)] \times \tilde{s}_{12}, \\
 & \quad [([\mathbf{13}]\langle\mathbf{24}\rangle + (3 \leftrightarrow 4)) + (1 \leftrightarrow 2)] \times (\tilde{s}_{13} - \tilde{s}_{14}) & (8;8) \# = 2
 \end{aligned} \tag{4.14}$$

For four identical Majorana neutrinos, one has

$$\begin{aligned}
 ++++ : & \quad [\mathbf{12}][\mathbf{34}] \times \tilde{s}_{12} + \text{Perm}(1234); \text{ PF} & (8;8) \# = 2 \\
 +++- : & \quad [\mathbf{12}]\langle\mathbf{34}\rangle \times \tilde{s}_{12} + \text{Perm}(1234) & (8;8) \# = 1
 \end{aligned} \tag{4.15}$$

For the same flavor and Majorana neutrinos, the missing SCTs in the $(+++-)$ helicity category appear at E^5 .

4.2.4 $W^+W^- f^c f$ and $WZ f^c f'$

$$\begin{aligned}
 00++ : & \quad \{\langle\mathbf{12}\rangle[\mathbf{12}][\mathbf{34}], \langle\mathbf{12}\rangle[\mathbf{13}][\mathbf{24}]\}; \text{ PF} & (5;8) \# = 4 \\
 0+-+ : & \quad \langle\mathbf{14}\rangle[\mathbf{12}][\mathbf{23}]; (1 \leftrightarrow 2); (3 \leftrightarrow 4); \text{ PF} & (6;8) \# = 8 \\
 00+- : & \quad \{\langle\mathbf{14}\rangle\langle\mathbf{231}\rangle[\mathbf{23}], (1 \leftrightarrow 2)\}; \text{ PF} & (6;8) \# = 4 \\
 ++++ : & \quad \{[\mathbf{12}]^2[\mathbf{34}], [\mathbf{12}][\mathbf{13}][\mathbf{24}]\}; \text{ PF} & (7;8) \# = 4 \\
 +++- : & \quad [\mathbf{12}]^2\langle\mathbf{34}\rangle; \text{ PF} & (7;8) \# = 2 \\
 0-++ : & \quad \{\langle\mathbf{12}\rangle[\mathbf{34}]\langle\mathbf{241}\rangle, (1 \leftrightarrow 2)\}; \text{ PF} & (7;8) \# = 4 \\
 0+++ : & \quad \{[\mathbf{132}][\mathbf{12}][\mathbf{34}], \langle\mathbf{132}\rangle[\mathbf{13}][\mathbf{24}]\}; (1 \leftrightarrow 2); \text{ PF} & (7;8) \# = 8 \\
 +++- : & \quad \{[\mathbf{12}]^2[\mathbf{314}], (3 \leftrightarrow 4)\}; \text{ PF} & (8;8) \# = 4 \\
 +-+- : & \quad \{[\mathbf{14}][\mathbf{132}]\langle\mathbf{23}\rangle, (1 \leftrightarrow 2)\}; (3 \leftrightarrow 4) & (8;8) \# = 4
 \end{aligned} \tag{4.16}$$

There is a non-trivial reduction of the spinor basis for the $(0-++)$ helicity category, but the reduction appears as a linear combination of terms with higher energy growth which we

have neglected. Thus all of the structures appear with independent coefficients in our basis. All independent SCTs appear at E^4 order for distinguishable fermions.

For identical Majorana neutrinos, $W^+W^-\nu\nu$,

$$\begin{aligned}
 00++ &: \langle \mathbf{12} \rangle [\mathbf{12}] [\mathbf{34}]; \text{PF} & (5;8) - 2 \\
 0+ +- &: \{ \langle \mathbf{14} \rangle [\mathbf{12}] [\mathbf{23}] - (3 \leftrightarrow 4), (1 \leftrightarrow 2) \}; \text{PF} & (6;8) - 4 \\
 00+ - &: \langle \mathbf{14} \rangle \langle \mathbf{231} \rangle [\mathbf{23}] - (3 \leftrightarrow 4), (1 \leftrightarrow 2) & (6;8) \# = 2 \\
 +++ + &: [\mathbf{12}]^2 [\mathbf{34}]; \text{PF} & (7;8) \# = 2 \\
 ++ - - &: [\mathbf{12}]^2 \langle \mathbf{34} \rangle; \text{PF} & (7;8) \# = 2 \\
 0+ ++ &: \{ \langle \mathbf{132} \rangle [\mathbf{13}] [\mathbf{24}], (1 \leftrightarrow 2) \}; \text{PF} & (7;8) \# = 4 \\
 ++ + - &: [\mathbf{12}]^2 \langle \mathbf{314} \rangle - (3 \leftrightarrow 4); \text{PF} & (8;8) \# = 2 \\
 + - - + &: [\mathbf{14}] [\mathbf{132}] \langle \mathbf{23} \rangle - (3 \leftrightarrow 4), (1 \leftrightarrow 2) & (8;8) \# = 2
 \end{aligned} \tag{4.17}$$

The missing $(0 - ++)$ helicity category SCT first appears at E^6 as $(s_{13} - s_{14}) \langle \mathbf{12} \rangle [\mathbf{34}] \langle \mathbf{2(3-4)1} \rangle$ with four independent coefficients.

4.2.5 ZZf^cf

$$\begin{aligned}
 00++ &: \langle \mathbf{12} \rangle [\mathbf{12}] [\mathbf{34}]; \text{PF} & (5;8) \# = 2 \\
 0+ +- &: \{ \langle \mathbf{14} \rangle [\mathbf{12}] [\mathbf{23}] + (1 \leftrightarrow 2), (3 \leftrightarrow 4) \}; \text{PF} & (6;8) \# = 4 \\
 00+ - &: \langle \mathbf{14} \rangle \langle \mathbf{231} \rangle [\mathbf{23}] + (1 \leftrightarrow 2); \text{PF} & (6;8) \# = 2 \\
 +++ + &: [\mathbf{12}]^2 [\mathbf{34}]; \text{PF} & (7;8) \# = 2 \\
 ++ - - &: [\mathbf{12}]^2 \langle \mathbf{34} \rangle; \text{PF} & (7;8) \# = 2 \\
 0- ++ &: \langle \mathbf{12} \rangle [\mathbf{34}] \langle \mathbf{241} \rangle + (1 \leftrightarrow 2); \text{PF} & (7;8) \# = 2 \\
 0+ ++ &: \{ \langle \mathbf{132} \rangle [\mathbf{12}] [\mathbf{34}], \langle \mathbf{132} \rangle [\mathbf{13}] [\mathbf{24}] \} + (1 \leftrightarrow 2); \text{PF} & (7;8) \# = 4 \\
 + - - + &: [\mathbf{14}] [\mathbf{132}] \langle \mathbf{23} \rangle + (1 \leftrightarrow 2), (3 \leftrightarrow 4) & (8;8) \# = 2
 \end{aligned} \tag{4.18}$$

All independent SCTs appear at E^4 .

For identical Majorana neutrinos,

$$\begin{aligned}
 00++ &: \langle \mathbf{12} \rangle [\mathbf{12}] [\mathbf{34}]; \text{PF} & (5;8) \# = 2 \\
 0+ +- &: [\langle \mathbf{14} \rangle [\mathbf{12}] [\mathbf{23}] + (1 \leftrightarrow 2)] - (3 \leftrightarrow 4); \text{PF} & (6;8) \# = 2 \\
 00+ - &: [\langle \mathbf{14} \rangle \langle \mathbf{231} \rangle [\mathbf{23}] + (1 \leftrightarrow 2)] - (3 \leftrightarrow 4) & (6;8) \# = 1 \\
 +++ + &: [\mathbf{12}]^2 [\mathbf{34}]; \text{PF} & (7;8) \# = 2 \\
 ++ - - &: [\mathbf{12}]^2 \langle \mathbf{34} \rangle; \text{PF} & (7;8) \# = 2 \\
 0+ ++ &: [\langle \mathbf{132} \rangle [\mathbf{13}] [\mathbf{24}] + (1 \leftrightarrow 2)] - (3 \leftrightarrow 4); \text{PF} & (7;8) \# = 2 \\
 + - - + &: [[\mathbf{14}] [\mathbf{132}] \langle \mathbf{23} \rangle + (1 \leftrightarrow 2)] - (3 \leftrightarrow 4) & (8;8) \# = 1
 \end{aligned} \tag{4.19}$$

The missing $(0 - ++)$ helicity category SCT first appears at E^6 from symmetrizing $(s_{13} - s_{14}) \langle \mathbf{12} \rangle [\mathbf{34}] \langle \mathbf{2(3-4)1} \rangle$ with two independent coefficients.

4.3 Bosonic amplitudes with massless vectors

4.3.1 γhhh

There is a single structure appearing at dimension 13, using $(\tilde{s}_{12} - \tilde{s}_{13})(\tilde{s}_{12} - \tilde{s}_{14})(\tilde{s}_{13} - \tilde{s}_{14})([\mathbf{1231}] - \langle \mathbf{1231} \rangle)$.

4.3.2 $\gamma Zh h$

$$\begin{aligned} ++ : & \quad \tilde{s}_{12}[12]^2; \text{PF} & (8;8) \# = 2 \\ +- : & \quad [132]^2 + (3 \leftrightarrow 4); \text{PF} & (8;8) \# = 2 \end{aligned} \tag{4.20}$$

All independent $vvss$ SCTs appear already at E^4 .

4.3.3 $gghh$ and $\gamma\gamma hh$

$$\begin{aligned} ++ : & \quad \tilde{s}_{12}[12]^2; \text{PF} & (8;8) \# = 2 \\ +- : & \quad [[132]^2 + (3 \leftrightarrow 4)] + (1 \leftrightarrow 2) & (8;8) \# = 1 \end{aligned} \tag{4.21}$$

The δ^{AB} color factors are suppressed in $gghh$. All independent $vvss$ SCTs appear already at E^4 . These amplitudes were derived in [31] up to dimension-10.

4.3.4 $gggh$

$$+++ : \quad f^{ABC}[12][13][23]; \text{PF} \quad (7;8) \# = 2 \tag{4.22}$$

The $(++-)$ helicity amplitude is first generated at E^5 as $f^{ABC}[12]^3\langle 13 \rangle \langle 23 \rangle$ with 2 independent coefficients. The $gggh$ amplitudes are given to dimension-13 in [31].

4.3.5 $gg\gamma h$ and $\gamma\gamma\gamma h$

The only structures with E^n , $n \leq 4$ are $[12][13][23]$ which are manifestly antisymmetric under 1 – 2 exchange and therefore do not appear. This SCT structure first appears at E^9 by multiplying by $(s_{12} - s_{13})(s_{21} - s_{23})(s_{31} - s_{32})$ with two independent coefficients. The $++-$ helicity amplitude is first generated at E^7 with $[12]^3\langle 13 \rangle \langle 23 \rangle (s_{13} - s_{23})$. The $\gamma\gamma\gamma h$ amplitudes can be easily obtained from the $gggh$ amplitudes given in [31].

4.3.6 $ggZh$ and $\gamma\gamma Zh$

$$\begin{aligned} ++0 : & \quad [12]^2[313] + (1 \leftrightarrow 2); \text{PF} & (7;8) \# = 2 \\ +-0 : & \quad [13][142]\langle 23 \rangle + (1 \leftrightarrow 2) & (7;8) \# = 1 \end{aligned} \tag{4.23}$$

We suppressed the δ^{AB} color factor in $ggZh$. For same-helicity gluons or photons, the helicity category $(\pm\pm\pm)$ first appears at E^5 with two independent coefficients. For opposite gluon or photon helicities, $(\pm\mp\pm)$ first appears at E^5 with two independent coefficients. (Note that $(\pm\pm\mp)$ which could appear at E^7 , is reducible to a linear combination of other structures multiplied by Mandelstams and the Z mass [55].)

4.3.7 $\gamma ZZ h$

$$\begin{aligned} ++0 : & \quad \{[12]^2[313] + (2 \leftrightarrow 3), [12]^2[323] + (2 \leftrightarrow 3)\}; \text{PF} & (7;8) \# = 4 \\ +-0 : & \quad [13][142]\langle 23 \rangle + (2 \leftrightarrow 3); \text{PF} & (7;8) \# = 2 \end{aligned} \tag{4.24}$$

The helicity category $(+++)$ first appears at E^5 with two independent coefficients, while $(-++)$ first appears at E^7 with two independent coefficients. The $(+-+)$ structure first appears at E^7 but is reducible to a linear combination of other structures multiplied by Mandelstams and the Z mass [55].

4.3.8 $\gamma WW h$

$$\begin{aligned}
 +00 : & \quad [12]\langle 23 \rangle [31]; \text{ PF} & (5; 8) \quad \# = 2 \\
 ++0 : & \quad \{[12]^2[313], [12]^2[323]\}; (2 \leftrightarrow 3); \text{ PF} & (7; 8) \quad \# = 8 \\
 +-0 : & \quad [13][142]\langle 23 \rangle, (2 \leftrightarrow 3) & (7; 8) \quad \# = 4 \\
 +++ : & \quad [12][13][23]; \text{ PF} & (7, 8) \quad \# = 2
 \end{aligned} \tag{4.25}$$

The helicity category $(-++)$ first appears at E^7 with four independent coefficients. The $(++-)$ structure first appears at E^5 but is reducible to a linear combination of other structures multiplied by Mandelstams and the W mass [55].

4.3.9 $\gamma\gamma\gamma$

$$\begin{aligned}
 +++ : & \quad [12]^2[34]^2 + [13]^2[24]^2 + [14]^2[23]^2; \text{ PF} & (8; 8) \quad \# = 2 \\
 +- - : & \quad [12]^2\langle 34 \rangle^2 + \text{Perm}(1234) & (8; 8) \quad \# = 1
 \end{aligned} \tag{4.26}$$

There is an additional structure in the helicity category $(+++)$ which first appears at E^6 .

4.3.10 $gggZ$, $ggg\gamma$ and $\gamma\gamma Z$

$$\begin{aligned}
 +- - : & \quad [12]^2\langle 34 \rangle^2 + \text{Perm}(123); \text{ PF} & (8; 8) \quad \# = 2 \\
 +++ : & \quad [12]^2[34]^2 + [13]^2[24]^2 + [14]^2[23]^2; \text{ PF} & (8; 8) \quad \# = 2
 \end{aligned} \tag{4.27}$$

We suppressed the d^{ABC} color factor. There are no contact terms with an f^{ABC} structure at this order. The $gggZ$ amplitudes were worked out in [31] up to dimension-12. There are additional structures in the $(+++)$, $(++-)$, and $(+-+)$ helicity categories which first appear at E^5 , E^7 , and E^{10} respectively. Note that the $(+++)$ and $(++-)$ structures are not present for the $ggg\gamma$ contact term, and the $(+-+)$ helicity category structure is reducible. $ggg\gamma$ can be obtained by unbolding. The $\gamma\gamma Z$ contact terms can be obtained from $gggZ$ with $f^{ABC} = 0$ and $d^{ABC} = 1$.

4.3.11 $ggZZ$ and $\gamma\gamma ZZ$

$$\begin{aligned}
 ++00 : & \quad \{[12]^2[34]\langle 34 \rangle\}; \text{ PF} & (6; 8) \quad \# = 2 \\
 +-00 : & \quad [13][14]\langle 23 \rangle\langle 24 \rangle + (1 \leftrightarrow 2) & (6; 8) \quad \# = 1 \\
 ++++ : & \quad \{[13]^2[24]^2 + (1 \leftrightarrow 2), [13][14][23][24]\}; \text{ PF} & (8; 8) \quad \# = 4 \\
 +- - - : & \quad [12]^2\langle 34 \rangle^2; \text{ PF} & (8; 8) \quad \# = 2 \\
 -+ - + : & \quad ([24]^2\langle 13 \rangle^2 + [14]^2\langle 23 \rangle^2) + (3 \leftrightarrow 4) & (8; 8) \quad \# = 1
 \end{aligned} \tag{4.28}$$

For $ggZZ$, there is a δ^{AB} color factor. There are additional structures in the $(+++)$, $(++-)$, and $(+-+)$ helicity categories which first appear at E^5 , E^7 , and E^6 respectively. Note that the $(+-+)$ helicity category structures are reducible.

4.3.12 $ggWW$ and $\gamma\gamma WW$

$$\begin{aligned}
 ++00 : & \quad \{[12]^2[34]\langle 34 \rangle\}; \text{ PF} & (6; 8) \quad \# = 2 \\
 +-00 : & \quad [13][14]\langle 23 \rangle\langle 24 \rangle + (1 \leftrightarrow 2) & (6; 8) \quad \# = 1 \\
 ++++ : & \quad \{[12]^2[34]^2, [13]^2[24]^2 + (1 \leftrightarrow 2)\}; \text{ PF} & (8; 8) \quad \# = 4 \\
 +- - - : & \quad \{[12]^2\langle 34 \rangle^2\}; \text{ PF} & (8; 8) \quad \# = 2 \\
 -+ - + : & \quad \{[14]^2\langle 23 \rangle^2 + (1 \leftrightarrow 2)\}; \text{ PF} & (8; 8) \quad \# = 2
 \end{aligned} \tag{4.29}$$

For $ggWW$ there is a δ^{AB} color factor. There are additional structures in the $(+++0)$, $(++-0)$, and $(+-++)$ helicity categories which first appear at E^5 , E^7 , and E^6 respectively. Note that the $(+++-)$ helicity category structures are reducible.

4.3.13 γZWW

$$\begin{aligned}
 ++00 &: \{[12]^2[34]\langle 34 \rangle, [12][13][24]\langle 34 \rangle\}; \text{Perm}(+00); \text{PF} & (6;8) \# = 12 \\
 +-00 &: \{[13][14]\langle 23 \rangle\langle 24 \rangle, \text{Perm}(-00)\}; \text{PF} & (6;8) \# = 6 \\
 ++++ &: \{[12]^2[34]^2, [13]^2[24]^2, [14]^2[23]^2\}; \text{PF} & (8;8) \# = 6 \\
 ++-- &: \{[12]^2\langle 34 \rangle^2, \text{Perm}(+-)\}; \text{PF} & (8;8) \# = 6
 \end{aligned} \tag{4.30}$$

There are additional structures in the $(+++0)$, $(++-0)$, and $(+++-)$ helicity categories which first appear at E^5 , E^5 , and E^6 respectively. Note that the $(-+++)$ helicity category structures are reducible.

4.3.14 γZZZ

$$\begin{aligned}
 ++00 &: [12]^2[34]\langle 34 \rangle + [13]^2[24]\langle 24 \rangle + [14]^2[23]\langle 23 \rangle; \text{PF} & (6;8) \# = 2 \\
 +-00 &: [13][14]\langle 23 \rangle\langle 24 \rangle + \text{Perm}(234); \text{PF} & (6;8) \# = 2 \\
 ++++ &: [12]^2[34]^2 + [13]^2[24]^2 + [14]^2[23]^2; \text{PF} & (8;8) \# = 2 \\
 ++-- &: [12]^2\langle 34 \rangle^2 + \text{Perm}(234); \text{PF} & (8;8) \# = 2
 \end{aligned} \tag{4.31}$$

There are additional structures in the $(+++0)$, $(++-0)$, and $(+++-)$ helicity categories which first appear at E^5 , E^5 , and E^6 respectively. Note that the $(-+++)$ helicity category structures are reducible.

4.3.15 $gg\gamma\gamma$

$$\begin{aligned}
 ++++ &: \{[12]^2[34]^2, [13]^2[24]^2 + (1 \leftrightarrow 2)\}; \text{PF} & (8;8) \# = 4 \\
 ++-- &: \{[12]^2\langle 34 \rangle^2\}; \text{PF} & (8;8) \# = 2 \\
 +-+- &: [[13]^2\langle 24 \rangle^2 + (1 \leftrightarrow 2)] + (3 \leftrightarrow 4) & (8;8) \# = 1
 \end{aligned} \tag{4.32}$$

We suppressed the δ^{AB} group factor for the gluons. The $(+++-)$ helicity category structure first appears at E^6 .

4.3.16 $gg\gamma Z$

$$\begin{aligned}
 ++++ &: \{[12]^2[34]^2, [13]^2[24]^2 + (1 \leftrightarrow 2)\}; \text{PF} & (8;8) \# = 4 \\
 ++-- &: \{[12]^2\langle 34 \rangle^2\}; \text{PF} & (8;8) \# = 2 \\
 +-+- &: \{[14]^2\langle 23 \rangle^2 + (1 \leftrightarrow 2)\}; \text{PF} & (8;8) \# = 2
 \end{aligned} \tag{4.33}$$

We suppressed the δ^{AB} group factor for the gluons. There are additional structures in the $(+++0)$, $(++-0)$, and $(+++-)$ helicity categories which first appear at E^5 , E^7 , and E^6 respectively. Note that the $(+++-)$ helicity category structures are reducible.

4.3.17 $gggg$

$$\begin{aligned}
 ++++ &: \{\mathcal{G} \times [12]^2[34]^2 + \text{Perm}(1234), f^{ABE}f^{CDE}[13]^2[24]^2 + \text{Perm}(1234)\} & (8;8) \# = 3 \\
 ---- &: \{\mathcal{G} \times \langle 12 \rangle^2\langle 34 \rangle^2 + \text{Perm}(1234), f^{ABE}f^{CDE}\langle 13 \rangle^2\langle 24 \rangle^2 + \text{Perm}(1234)\} & (8;8) \# = 3 \\
 ++-- &: \{\mathcal{G}, f^{ACE}f^{BDE} + f^{BCE}f^{ADE}\} \times ([12]^2\langle 34 \rangle^2 + \text{Perm}(1234)) & (8;8) \# = 3
 \end{aligned} \tag{4.34}$$

Here $\mathcal{G} = \{\delta^{AB}\delta^{CD}, d^{ABE}d^{CDE}\}$ is a set of SU(3) structures. There are additional structures in the helicity category $(+++-)$ which first appears at E^6 .

4.4 Fermionic amplitudes with massless vectors

4.4.1 $f^c f \gamma h$ and $f^c f g h$

$$\begin{aligned}
 +++ &: [\mathbf{13}][\mathbf{23}] \times \{s_{12}, s_{13}\}; \text{PF } (8;8) \# = 4 \\
 +-+ &: \{[\mathbf{13}][\mathbf{312}], (1 \leftrightarrow 2)\}; \text{PF } (7;8) \# = 4 \\
 ++- &: [\mathbf{12}]\langle\mathbf{3123}\rangle; \text{PF } (8;8) \# = 2
 \end{aligned} \tag{4.35}$$

The gluon amplitude is only nonzero when the fermions are quarks, in which case it appears with the color factor $(\lambda^A)_a^b$. All SCT bases are covered at E^4 . For identical Majorana neutrinos, the independent structures are,

$$\begin{aligned}
 +++ &: [\mathbf{13}][\mathbf{23}] \times (s_{13} - s_{23}); \text{PF } (8;8) \# = 2 \\
 +-+ &: [\mathbf{13}][\mathbf{312}] - (1 \leftrightarrow 2); \text{PF } (7;8) \# = 2
 \end{aligned} \tag{4.36}$$

In this case the $(++-)$ helicity category only appears at E^6 with the two independent E^4 SCTs multiplied by $s_{13} - s_{14}$.

4.4.2 $ggf^c f$ and $\gamma\gamma f^c f$

$$\begin{aligned}
 ++++ &: [12]^2[\mathbf{34}]; \text{PF } (7;8) \# = 2 \\
 +++- &: [12]^2\langle\mathbf{34}\rangle; \text{PF } (7;8) \# = 2 \\
 +-+- &: [14][\mathbf{132}]\langle\mathbf{23}\rangle + (1 \leftrightarrow 2); (3 \leftrightarrow 4) (8;8) \# = 2
 \end{aligned} \tag{4.37}$$

The gluon contact terms are proportional to δ^{AB} . The fermions form an SU(3) singlet in both cases. For identical Majorana neutrinos one has

$$\begin{aligned}
 ++++ &: [12]^2[\mathbf{34}]; \text{PF } (7;8) \# = 2 \\
 +++- &: [12]^2\langle\mathbf{34}\rangle; \text{PF } (7;8) \# = 2 \\
 +-+- &: ([14][\mathbf{132}]\langle\mathbf{23}\rangle + (1 \leftrightarrow 2)) - (3 \leftrightarrow 4) (8;8) \# = 1
 \end{aligned} \tag{4.38}$$

There are structures in the $(+++-)$ and $(-+++)$ helicity categories which first appear at E^6 and E^5 respectively.

4.4.3 $\gamma g f^c f$

$$\begin{aligned}
 ++++ &: \{[12]^2[\mathbf{34}], [12][\mathbf{13}][\mathbf{24}]\}; \text{PF } (7;8) \# = 4 \\
 +++- &: [12]^2\langle\mathbf{34}\rangle; \text{PF } (7;8) \# = 2 \\
 +++- &: \{[12]^2[\mathbf{314}], (3 \leftrightarrow 4)\}; \text{PF } (8;8) \# = 4 \\
 +-+- &: \{[14][\mathbf{132}]\langle\mathbf{23}\rangle, (1 \leftrightarrow 2)\}; (3 \leftrightarrow 4) (8;8) \# = 4
 \end{aligned} \tag{4.39}$$

This amplitude is only nonzero for quarks, and involves the color factor $(\lambda^A)_a^b$. There is an SCT in the $(-+++)$ helicity categories which first contributes at E^5 .

4.4.4 $\gamma Z f^c f$, $\gamma W f^c f'$, $g Z f^c f$ and $g W f^c f'$

$$\begin{aligned}
 +0+-: & \quad \{[12][13]\langle 24 \rangle, (3 \leftrightarrow 4)\}; \text{ PF} & (6;8) \quad \# = 4 \\
 + + + +: & \quad \{[12]^2[34], [12][13][24]\}; \text{ PF} & (7;8) \quad \# = 4 \\
 + + --: & \quad [12]^2\langle 34 \rangle; \text{ PF} & (7;8) \quad \# = 2 \\
 -0++: & \quad \langle 12 \rangle[34]\langle 142 \rangle; \text{ PF} & (7;8) \quad \# = 2 \\
 +0++: & \quad \{\langle 231 \rangle[12][34], \langle 231 \rangle[23][14]\}; \text{ PF} & (7;8) \quad \# = 4 \\
 + + +-: & \quad \{[12]^2[314], (3 \leftrightarrow 4)\}; \text{ PF} & (8;8) \quad \# = 4 \\
 + - - +: & \quad \{[14][132]\langle 23 \rangle, (3 \leftrightarrow 4)\}; \text{ PF} & (8;8) \quad \# = 4
 \end{aligned} \tag{4.40}$$

The gluon amplitude is only non-zero for quarks, and appears with $(\lambda^A)_a^b$. There is an additional structure in the $(-+++)$ helicity category which first appears at E^5 .

For identical Majorana neutrinos, $\gamma Z \nu \nu$, one has instead,

$$\begin{aligned}
 +0+-: & \quad [12][13]\langle 24 \rangle - (3 \leftrightarrow 4); \text{ PF} & (6;8) \quad \# = 2 \\
 + + + +: & \quad [12]^2[34]; \text{ PF} & (7;8) \quad \# = 2 \\
 + + --: & \quad [12]^2\langle 34 \rangle; \text{ PF} & (7;8) \quad \# = 2 \\
 +0++: & \quad \langle 231 \rangle[23][14] - (3 \leftrightarrow 4); \text{ PF} & (7;8) \quad \# = 2 \\
 + + +-: & \quad [12]^2[314] - (3 \leftrightarrow 4); \text{ PF} & (8;8) \quad \# = 2 \\
 + - - +: & \quad [14][132]\langle 23 \rangle - (3 \leftrightarrow 4); \text{ PF} & (8;8) \quad \# = 2
 \end{aligned} \tag{4.41}$$

The $(-0++)$ helicity-category SCT first appears at E^6 .

5 Conclusions

The on-shell bootstrap is ideally suited to the derivation of low-energy effective amplitudes, since the latter are fully determined by the particle content and assumed symmetry. In this paper, we have applied these methods to parametrize the four-point local amplitudes of the known particles, keeping structures scaling with the energy as $E^{n \leq 4}$. These are given by a set of independent contact terms, which can be organized in terms of linear combinations of independent, manifestly-local spinor structures, or SCTs, multiplied by expansions in the Mandelstam invariants.

Our results provide the basic building blocks for collider EFT searches. Two-to-two EFT scattering amplitudes can be derived from the set of SM three-point amplitudes obtained in [33], and the four-point contact terms derived here. Together, these contact terms also parametrize two- and three-particle decay amplitudes. The resulting EFT formulation involves just physical quantities. We also discuss the mapping of the contact terms to the HEFT and the SMEFT. In particular, in table 3, we derive the low-energy SMEFT contact terms, and relate them to the Warsaw basis. These results can be used to distinguish between the HEFT and SMEFT frameworks, and to identify observables that are particularly sensitive to different types of UV models.

Acknowledgments

We thank Csaba Csaki, Gauthier Durieux, Christophe Grojean and Markus Luty for discussions. YS and MW thank the Mainz Institute for Theoretical Physics (MITP) of the

Cluster of Excellence PRISMA+ (Project ID 39083149), for its hospitality and support during the workshop Amplitudes meet BSM. YS thanks the Aspen Center for physics, which is supported by the National Science Foundation (grant PHY-1607611), where parts of this work were completed. Research supported in part by the Israel Science Foundation (Grant No. 751/19), and by the NSF-BSF (Grant No. 2020-785). The research of TM is also supported by “Study in Israel” Fellowship for Outstanding Post-Doctoral Researchers from China and India by PBC of CHE. The research of MW is also supported by a Zuckerman Fellowship. HL is supported by the ISF, BSF and the Azrieli Foundation

A *WWhh*: on-shell construction of the HEFT and SMEFT amplitudes and on-shell Higgsing

In this appendix we provide a detailed derivation of the *WWhh* HEFT and SMEFT amplitudes. This illustrates several points:

- Mass-suppressed contact terms, which appear for longitudinal-vector helicity categories are associated with the factorizable part of the SMEFT amplitude.
- The coefficients of these terms are determined by the three-point couplings based on the high-energy limit of the amplitude. This can be done in two equivalent ways: requiring that the zero-polarization amplitude has no E/m growth, or requiring that the transverse-polarization amplitude does not depend on spurious spinors. This latter requirement is nothing but gauge invariance, so the equivalence of gauge invariance and perturbative unitarity are manifest.
- The remaining contact terms can then be determined by “Higgsing” the massless $4 + n_H$ contact terms.

A.1 The structure of the full amplitude

The on-shell construction of the *WWhh* amplitude requires as inputs the *WWh* and *hhh* 3-point couplings, as well as the 4-point *WWhh* contact terms. In the HEFT, the 3- and 4-point couplings are the most general ones consistent with the symmetry of the low-energy theory. In the SMEFT, both the 3-points and the 4-points can be derived by Higgsing the massless amplitudes in the unbroken phase.

The most general three points consistent with the symmetries of the low-energy theory are [33],

$$\mathcal{M}(W^+, W^-, h) = C_{WWh}^{00} \frac{\langle \mathbf{12} \rangle [\mathbf{12}]}{M_W} + C_{WWh}^{++} \frac{[\mathbf{12}] [\mathbf{12}]}{\bar{\Lambda}} + C_{WWh}^{--} \frac{\langle \mathbf{12} \rangle \langle \mathbf{12} \rangle}{\bar{\Lambda}}, \quad (\text{A.1})$$

$$\mathcal{M}(h, h, h) = m_h C_{hhh}. \quad (\text{A.2})$$

The three-point couplings C_{WWh} and C_{hhh} are numbers, since there is no kinematic dependence in 3-point amplitudes. In the SMEFT, they are given as an expansion in

v/Λ . In this subsection, we neglect the terms with $C_{WW_h}^{\pm\pm}$ because they will not affect our discussion.³ The $WWhh$ amplitude is then,

$$\begin{aligned}
 \mathcal{M}(W^+, W^-, h, h) = & -\frac{m_h M_W C_{WW_h}^{00} c_{hhh} [\mathbf{12}]\langle\mathbf{12}\rangle}{s_{12} - m_h^2} \frac{1}{M_W^2} \\
 & + \frac{C_{WW_h}^{00}}{s_{13} - M_W^2} \left(\frac{\langle\mathbf{131}\rangle\langle\mathbf{242}\rangle}{2M_W^2} - [\mathbf{12}]\langle\mathbf{12}\rangle \right) \\
 & + \frac{C_{WW_h}^{00}}{s_{14} - M_W^2} \left(\frac{\langle\mathbf{141}\rangle\langle\mathbf{232}\rangle}{2M_W^2} - [\mathbf{12}]\langle\mathbf{12}\rangle \right) \\
 & + C_{WWhh}^{00, fac} \frac{[\mathbf{12}]\langle\mathbf{12}\rangle}{M_W^2} + C_{WWhh}^{00, CT} \frac{[\mathbf{12}]\langle\mathbf{12}\rangle}{\bar{\Lambda}^2} \\
 & + C_{WWhh}^{\prime 00, fac} \frac{[\mathbf{12}]\langle\mathbf{12}\rangle s_{12}}{M_W^2 \bar{\Lambda}^2} + C_{WWhh}^{\prime\prime 00, fac} \frac{\langle\mathbf{131}\rangle\langle\mathbf{242}\rangle + \langle\mathbf{141}\rangle\langle\mathbf{232}\rangle}{M_W^2 \bar{\Lambda}^2} \quad (\text{A.3})
 \end{aligned}$$

where we have kept terms up to E^4 . The first two lines of this expression are obtained by gluing the 3-point amplitudes.⁴ The last two lines of eq. (A.3) contain the most general manifestly-local structures, or contact terms.⁵ As explained above, to read off the operator dimension at which each longitudinal vector structure first appears, we normalize it with a factor of $1/M_W$. In the SMEFT, each of the couplings C_{WWhh} entering the amplitude is given as an expansion in v/Λ , and in particular, each power of $\bar{\Lambda}$ is given in terms of Λ and potentially v .

Note that we have isolated the $1/M_W^2$ pieces of the longitudinal-vector contact terms, and labeled them with the superscript *fac*: these pieces are associated with the *factorizable* parts of the amplitudes, and in the SMEFT arise from the factorizable parts of the massless amplitudes. Thus, these pieces are determined by the three-point couplings. The remaining pieces are labeled by CT for contact terms: at each dimension, these are the novel inputs in the theory, which arise in the SMEFT from massless contact terms. In the HEFT on the other hand, $\bar{\Lambda} = v \sim M_W$, there is no real separation between the two types of terms.

To determine the $1/M_W^2$ pieces, consider the expansion of the amplitude in terms of $s_{ij}/\bar{\Lambda}^2$ and s_{ij}/M_W^2 . For the *longitudinal-W* amplitude, this expansion starts as,

$$\begin{aligned}
 \mathcal{M}(W^{+(0)}, W^{-(0)}, h, h) \sim & -C_{WWhh}^{00, fac} \frac{s_{12}}{M_W^2} - C_{WWhh}^{00, CT} \frac{s_{12}}{\bar{\Lambda}^2} - \frac{C_{WW_h}^{00}}{2} \frac{s_{12}}{M_W^2} \\
 & - C_{WWhh}^{\prime 00, fac} \frac{s_{12}^2}{M_W^2 \bar{\Lambda}^2} + C_{WWhh}^{\prime\prime 00, fac} \frac{s_{13}^2 + s_{14}^2}{M_W^2 \bar{\Lambda}^2} \quad (\text{A.4})
 \end{aligned}$$

where we only kept terms which grow with the energy. The superscripts (0) denote the W polarization. At dimension-six, the amplitude should be unitary up to $E^2/\bar{\Lambda}^2$ terms. Thus,

³Our focus here is on the C_{WWhh}^{00} terms, and the relevant helicity amplitudes to consider are $\pm\mp$ and 00. The $C_{WW_h}^{\pm\pm}$ contributions are subleading in the high-energy limit of these amplitudes. One can show, either by Higgsing the high-energy amplitudes or by requiring good high-energy behavior of the amplitudes with the same W helicities that $C_{WW_h}^{\pm\pm} \propto v$.

⁴For details of this gluing, see ref. [33]. Note that this simple gluing can only be done for massive legs (see eg [49] for a recent discussion).

⁵In the notation of ref. [33], the coefficients C_{WWhh} of the spinor structures are expansions in the Mandelstams, but here we expanded these out.

pieces with E^2/M_W^2 should vanish, fixing,

$$C_{WWhh}^{00,fac} = -\frac{C_{WWh}^{00}{}^2}{2} \quad C_{WWhh}^{\prime 00,fac} = C_{WWhh}^{\prime\prime 00,fac} = 0. \quad (\text{A.5})$$

Therefore, at dimension-six, these coefficients are determined in terms of the three-point couplings. At dim-8, this implies cancellations between terms in $C_{WWhh}^{00,fac}$ and $C_{WWhh}^{\prime 00,fac}$. In the HEFT, one cannot take the high energy limit, since the cutoff cannot be much higher than the electroweak scale. Thus, the reasoning above only serves to differentiate between dimension-4, 6, etc contributions to the Wilson coefficients. In particular, since we can set $\bar{\Lambda} = v$, there are still independent dimension-six contact terms suppressed by v , such as $C_{WWhh}^{00,CT}[\mathbf{12}]\langle\mathbf{12}\rangle/v^2$.

In contrast, in the SMEFT, we can really take the high-energy limit as in eq. (A.4), with $M_W \sim v \ll E \ll \Lambda$. Furthermore, we can alternatively obtain the relations in eq. (A.5) by considering the high-energy limit of the amplitude with *transverse* W 's (which are finite in this limit),

$$A(W^{+(+)}, W^{(-)}, h, h) \sim c_{WWhh}^{00,fac} \frac{[1_k 2_q]\langle 1_q 2_k \rangle}{M_W^2} + \frac{c_{WWh}^{00}{}^2}{2M_W^2} \frac{[1_k 3 1_q]\langle 2_q 4 2_k \rangle}{s_{13}} + \frac{c_{WWh}^{00}{}^2}{2M_W^2} \frac{[1_k 4 1_q]\langle 2_q 3 2_k \rangle}{s_{14}}. \quad (\text{A.6})$$

The spinors 1_q) and 2_q) scale as the mass, and can be written as (see eg [47])

$$i_q) = \frac{M_W}{(i_k \xi_i)} \xi_i) \quad (\text{A.7})$$

where $\xi_i)$ is an arbitrary constant spinor. For the amplitude to be well-defined at high energies, it must be independent of the arbitrary spinors. Requiring that the amplitude in eq. (A.6) is independent of these arbitrary spinors, one recovers the relation

$$C_{WWhh}^{00,fac} = -\frac{C_{WWh}^{00}{}^2}{2}. \quad (\text{A.8})$$

Indeed, in the SMEFT, the high energy EFT has the full unbroken gauge symmetry, and the role of the arbitrary spinors ξ_i is clear — these are the reference spinors for the two massless vector polarizations [47] (see also [66]). Thus the equivalence of perturbative unitarity and the restoration of gauge symmetry is manifest in this example.

A.2 Four-point contact terms from on-shell Higgsing

As we saw above, the independent couplings appearing in the amplitude are the three-point couplings and the Λ -suppressed contact terms (with no M_W suppression). In the SMEFT, these can be determined by Higgsing the massless $SU(3) \times SU(2) \times U(1)$ -symmetric amplitudes [47]. Two examples, namely $WWhh$ and $\bar{u}dWh$ were worked out in detail in ref. [47] up to dimension-8. Here we briefly repeat the $WWhh$ derivation for completeness, keeping only dimension-six terms. The results for all the four-points at dimension-six appear in table 3.

The dimension-six contributions to the three points were derived in ref. [33] by matching the amplitudes to the broken-phase SMEFT using [65]. We do not repeat the derivation using on-shell Higgsing. Instead, we just show a few examples deriving the three-points from on-shell Higgsing in the next subsection.

The low-energy $WWhh$ contact terms originate from a number of high-energy contact terms. The coefficient of each structure in a given helicity category is determined, to leading order in v/Λ , by the corresponding helicity amplitude in the high-energy theory. For instance, the massless $WWH^\dagger H$ amplitudes give the leading-order contribution to the $++$ and $--$ helicity categories, while $(H^\dagger H)^2$ is the leading-order contribution to the 00 helicity category. Sub-leading v^2/Λ^2 contributions originate from higher-point *contact terms* such as $(H^\dagger H)^3$. Here, we just focus on their leading order pieces. It is worth noting that the high-energy contact terms fully determine the coefficients of the low-energy massive contact terms. Each massive contact term is of course a little-group tensor, and the derivation of its coefficient C_{low} essentially relies on the matching of the leading-energy component to the contact term of the corresponding high energy amplitude, with Wilson coefficient c_{high} . The sub-leading components of the massive structure are generated by factorizable high-energy amplitude featuring the coupling c_{high} .

Let us begin with the contribution from $(H^\dagger H)^2$. The high-energy amplitude is

$$\mathcal{A}(H^i H_k^\dagger H^j H_l^\dagger) \supset c_{HHHH}^+ \frac{s_{13}}{\Lambda^2} T^{+ij}_{kl} + c_{HHHH}^- \frac{s_{12} - s_{14}}{\Lambda^2} T^{-ij}_{kl} \quad (\text{A.9})$$

where $T^{\pm ij}_{kl} = (\delta_k^i \delta_l^j \pm \delta_l^i \delta_k^j)/2$ are the symmetric and anti-symmetric $SU(2)$ structures. Parameterizing the Higgs doublet as in (3.1), we find that

$$\mathcal{A}(G^+ G^- hh) = \frac{1}{2} \left(\mathcal{A}(H^1 H_1^\dagger H^2 H_2^\dagger) + \mathcal{A}(H^1 H_1^\dagger H_2^\dagger H^2) \right) = - \frac{c_{(H^\dagger H)^2}^+ - 3c_{(H^\dagger H)^2}^-}{2} \frac{s_{12}}{2\Lambda^2} \quad (\text{A.10})$$

which *bolds* into

$$- \frac{c_{(H^\dagger H)^2}^+ - 3c_{(H^\dagger H)^2}^-}{2} \frac{s_{12}}{2\Lambda^2} \rightarrow \frac{c_{(H^\dagger H)^2}^+ - 3c_{(H^\dagger H)^2}^-}{2} \frac{[\mathbf{12}]\langle \mathbf{12} \rangle}{\Lambda^2}. \quad (\text{A.11})$$

Thus there is a contact term in the massive EFT of the form,

$$C_{WWhh}^{00,CT} \frac{[\mathbf{12}]\langle \mathbf{12} \rangle}{\Lambda^2} = \frac{c_{(H^\dagger H)^2}^+ - 3c_{(H^\dagger H)^2}^-}{2} \frac{[\mathbf{12}]\langle \mathbf{12} \rangle}{\Lambda^2}. \quad (\text{A.12})$$

We now turn to the contribution of the massless $WWH^\dagger H$ contact terms,

$$\mathcal{A}(W^{a,\pm} W^{b,\pm} H_i^\dagger H^j) = c_{WWHH}^{\pm\pm} \frac{(12)^2}{\Lambda^2} \left(T^{ab} \right)_i^j. \quad (\text{A.13})$$

Here \pm are the helicities of the W 's, $(12) = [12]$ for $++$, $(12) = \langle 12 \rangle$ for $--$, and $\left(T^{ab} \right)_i^j = \delta_i^{ab} \delta_i^j$ is required by the symmetry of the spinor structure. The kinematics bold trivially $(12) \rightarrow \mathbf{(12)}$, and one finds the low-energy contact terms

$$C_{WWhh}^{\pm\pm} \frac{\mathbf{(12)}}{\Lambda^2} = 2c_{WWHH}^{\pm\pm} \frac{\mathbf{(12)}}{\Lambda^2}. \quad (\text{A.14})$$

A.3 The WWh coupling from on-shell Higgsing

The WWh coupling C_{WWh}^{00} in eq. (A.1) is given by the gauge coupling to leading order, with a v^2/Λ^2 correction at dimension-six. Here we show how both of these contributions can be obtained from on-shell Higgsing. The relevant massless amplitudes to consider are $WH^\dagger H$ or $WWH^\dagger H$ to determine the dimension-4 part, and $H^2(H^\dagger)^2 WW$ to determine the dimension-6 shift.

Our discussion closely parallels the derivation of three-points from the massless amplitudes of a toy model with higgsed U(1) symmetry in ref. [47], and we refer the reader to that paper for more details.

To determine C_{WWh}^{00} one can start from either of its components, which map to different high-energy amplitudes. Apriori, the obvious amplitude to start from is the massless three point $WH^\dagger H$, which for positive W helicity is,

$$\mathcal{A}((W^+)^+ G^- h) = \frac{g}{\sqrt{2}} \frac{[12][13]}{[23]} \tag{A.15}$$

where we parametrized the Higgs doublet as in eq. (3.1). Bolding this amplitude is not entirely straightforward because it is non-local, and indeed, its non-locality translates in the massive amplitude to the $1/M_W$ ‘‘pole’’. However, we can proceed by multiplying and dividing eq. (A.15) by $\langle 3\xi \rangle$ for some arbitrary ξ . After some manipulations using momentum conservation, we get

$$\mathcal{A}((W^+)^+ G^- h) = \frac{g}{\sqrt{2}} \frac{[12]\langle \xi 2 \rangle}{\langle 1\xi \rangle}. \tag{A.16}$$

Identifying $1_q] = \xi]$, this maps to the longitudinal W component of

$$C_{WWh}^{00} \frac{\langle \mathbf{12} \rangle [\mathbf{12}]}{M_W} \tag{A.17}$$

with $C_{WWh}^{00} = g$. Alternatively, a simpler way to get this coupling is to start from the 4-point amplitude $H^2(H^\dagger)^2$, which matches the all-longitudinal component of the spinor structure. At the renormalizable level, this amplitude,

$$\mathcal{A}_{\text{SM}}((W^-)^- (W^+)^+ hh) = -\frac{g^2}{2} \frac{\langle 132 \rangle}{\langle 231 \rangle}. \tag{A.18}$$

Identifying $i_q) = (M_W/(i_k 3)) 3)$ for $i = 1, 2$ and taking $p_4 \rightarrow 0$ gives,

$$\mathcal{A}_{\text{SM}}^m(hW^+W^-) = \frac{g^2 v}{2} \frac{[\mathbf{23}]\langle \mathbf{23} \rangle}{M_W^2} \tag{A.19}$$

where we also used $\langle i_k i_q \rangle = [i_q i_k] = M_W$. Therefore, at the leading order, $C_{WWh}^{00} = g^2 v/M_W$. Note that the arbitray ξ spinor that we identified with the q spinors above naturally arises in this case from the soft higgs leg, with $\hat{1}_q$ and $\hat{2}_q$ along $\hat{4}$ (see also ref. [66]).

Turning to the dimension-six correction to C_{WWh}^{00} , this originates, to leading order, from the 6-point $H^2(H^\dagger)^2 WW$ factorizable amplitude with a single insertion of a $H^2(H^\dagger)^2$ contact term. The massless amplitude can be written as,

$$\mathcal{A}_{d=6}^{\text{tot}}((H^0)^4 (W^+)^+ (W^-)^-) = \mathcal{A}^{(3)}((H^0)^4 (W^+)^+ (W^-)^-) + \mathcal{A}^{(4)}((H^0)^4 (W^+)^+ (W^-)^-),$$




Figure 1. Feynman-diagram of $H^2(H^\dagger)^2 WW$ factorizable amplitude.

with

$$\begin{aligned} \mathcal{A}^{(3)}((H^0)^4(W^+)^+(W^-)^-) &= \frac{8g^2C}{4} \left[\frac{[142]}{\langle 12 \rangle s_{41}} s_{63} + \frac{[162]}{\langle 12 \rangle s_{61}} s_{43} + \frac{[132]}{\langle 12 \rangle s_{31}} s_{64} \right] \frac{\langle 251 \rangle}{[21] s_{52}} \\ &\quad + (5 \leftrightarrow 4, 6, 3), \\ \mathcal{A}^{(4)}((H^0)^4(W^+)^+(W^-)^-) &= \frac{8g^2C}{4} \frac{\langle 25 \rangle [15]}{\langle 15 \rangle [25]} + (5 \leftrightarrow 3, 4, 6), \end{aligned} \tag{A.20}$$

and

$$C = (c_{(HH^\dagger)^2}^+ - 3c_{(HH^\dagger)^2}^-)/2. \tag{A.21}$$

The piece $\mathcal{A}^{(3)}((H^0)^4(W^+)^+(W^-)^-)$ is the sum over Feynman diagrams in which the vector legs attach to different scalar legs (left panel of figure 1), and $\mathcal{A}^{(4)}((H^0)^4(W^+)^+(W^-)^-)$ is the piece with the vector legs attached to the same scalar legs (see the right figure in figure 1). (Of course, only the sum is gauge invariant.) Taking the momenta of three H^0 legs-4,5,6 to be soft, only $\mathcal{A}^{(4)}((H^0)^4(W^+)^+(W^-)^-)$ survives,

$$\lim_{q_{4,5,6} \rightarrow 0} \mathcal{A}^{(4)}(H^0(q_4)H^0(q_5)H^0(q_6)H^0(3)(W^+)^+(1)(W^-)^-(2)) = 6g^2v^3C \frac{[1_k \xi 2_k]}{[2_k \xi 1_k]}, \tag{A.22}$$

where at the last stage we set $q_i \propto \xi$ for some arbitrary ξ as before. This holds to the same massive structure as above. Altogether, after adding in the renormalizable contribution we have,

$$\mathcal{M}_{d=6}^m(h(W^+)^+(W^-)^-) = g(1 + v^2C) \frac{[12] \langle 12 \rangle}{M_W}. \tag{A.23}$$

Note that this correction is nothing but the correction to the Higgs wave-function renormalization induced by the four-Higgs contact term C .

One can derive the SMEFT corrections to the remaining three-point low-energy amplitudes along the same lines. However, since the most general 3-point couplings were listed in ref. [33] based on Feynman diagrams matching, we do not do so here.

B Energy-growth of factorizable amplitudes

Massive amplitudes	E growth
$\mathcal{M}(Zh\bar{h}h)$	no factorizable contribution
$\mathcal{M}(VVhh)$	E^2/M^2
$\mathcal{M}(WWZh)$	$E^2/M^2, E^3/(M^2\bar{\Lambda})$
$\mathcal{M}(ZZZh)$	no factorizable contribution
$\mathcal{M}(WWWW/ZZWW)$	$E^3/(M^2\bar{\Lambda}), E^4/M^4, E^4/(M^2\bar{\Lambda}^2)$
$\mathcal{M}(ZZZZ)$	$E^2/M^2, E^2/(M\bar{\Lambda})$
$\mathcal{M}(ffVh)$	E/M
$\mathcal{M}(ffWW)$	$E^2/M^2, E^3/(M^2\bar{\Lambda})$
$\mathcal{M}(ffZZ)$	$E/M, E^2/(M\bar{\Lambda})$

Table 4. Leading energy growth in the massive factorizable amplitudes.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl. Phys. B* **268** (1986) 621 [[INSPIRE](#)].
- [2] E.E. Jenkins and A.V. Manohar, *Algebraic Structure of Lepton and Quark Flavor Invariants and CP Violation*, *JHEP* **10** (2009) 094 [[arXiv:0907.4763](#)] [[INSPIRE](#)].
- [3] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085 [[arXiv:1008.4884](#)] [[INSPIRE](#)].
- [4] G.F. Giudice, C. Grojean, A. Pomarol and R. Rattazzi, *The Strongly-Interacting Light Higgs*, *JHEP* **06** (2007) 045 [[hep-ph/0703164](#)] [[INSPIRE](#)].
- [5] R.S. Gupta, A. Pomarol and F. Riva, *BSM Primary Effects*, *Phys. Rev. D* **91** (2015) 035001 [[arXiv:1405.0181](#)] [[INSPIRE](#)].
- [6] A. Falkowski et al., *Rosetta: an operator basis translator for Standard Model effective field theory*, *Eur. Phys. J. C* **75** (2015) 583 [[arXiv:1508.05895](#)] [[INSPIRE](#)].
- [7] L. Berthier and M. Trott, *Towards consistent Electroweak Precision Data constraints in the SMEFT*, *JHEP* **05** (2015) 024 [[arXiv:1502.02570](#)] [[INSPIRE](#)].
- [8] L. Berthier and M. Trott, *Consistent constraints on the Standard Model Effective Field Theory*, *JHEP* **02** (2016) 069 [[arXiv:1508.05060](#)] [[INSPIRE](#)].
- [9] C. Hays, A. Helset, A. Martin and M. Trott, *Exact SMEFT formulation and expansion to $\mathcal{O}(v^4/\Lambda^4)$* , *JHEP* **11** (2020) 087 [[arXiv:2007.00565](#)] [[INSPIRE](#)].

- [10] T. Corbett, A. Helset, A. Martin and M. Trott, *EWPD in the SMEFT to dimension eight*, *JHEP* **06** (2021) 076 [[arXiv:2102.02819](#)] [[INSPIRE](#)].
- [11] S. Dawson, S. Homiller and M. Sullivan, *Impact of dimension-eight SMEFT contributions: A case study*, *Phys. Rev. D* **104** (2021) 115013 [[arXiv:2110.06929](#)] [[INSPIRE](#)].
- [12] F. Feruglio, *The chiral approach to the electroweak interactions*, *Int. J. Mod. Phys. A* **8** (1993) 4937 [[hep-ph/9301281](#)] [[INSPIRE](#)].
- [13] J. Bagger et al., *The strongly interacting $W W$ system: Gold plated modes*, *Phys. Rev. D* **49** (1994) 1246 [[hep-ph/9306256](#)] [[INSPIRE](#)].
- [14] V. Koulovassilopoulos and R.S. Chivukula, *The phenomenology of a nonstandard Higgs boson in $W_L W_L$ scattering*, *Phys. Rev. D* **50** (1994) 3218 [[hep-ph/9312317](#)] [[INSPIRE](#)].
- [15] C.P. Burgess, J. Matias and M. Pospelov, *A Higgs or not a Higgs? What to do if you discover a new scalar particle*, *Int. J. Mod. Phys. A* **17** (2002) 1841 [[hep-ph/9912459](#)] [[INSPIRE](#)].
- [16] B. Grinstein and M. Trott, *A Higgs-Higgs bound state due to new physics at a TeV*, *Phys. Rev. D* **76** (2007) 073002 [[arXiv:0704.1505](#)] [[INSPIRE](#)].
- [17] R. Alonso et al., *The Effective Chiral Lagrangian for a Light Dynamical “Higgs Particle”*, *Phys. Lett. B* **722** (2013) 330 [Erratum *ibid.* **726** (2013) 926] [[arXiv:1212.3305](#)] [[INSPIRE](#)].
- [18] D. Espriu, F. Mescia and B. Yengo, *Radiative corrections to $W_L W_L$ scattering in composite Higgs models*, *Phys. Rev. D* **88** (2013) 055002 [[arXiv:1307.2400](#)] [[INSPIRE](#)].
- [19] G. Buchalla, O. Catà and C. Krause, *Complete Electroweak Chiral Lagrangian with a Light Higgs at NLO*, *Nucl. Phys. B* **880** (2014) 552 [[arXiv:1307.5017](#)] [[INSPIRE](#)].
- [20] I. Brivio et al., *Disentangling a dynamical Higgs*, *JHEP* **03** (2014) 024 [[arXiv:1311.1823](#)] [[INSPIRE](#)].
- [21] R. Alonso, E.E. Jenkins and A.V. Manohar, *A Geometric Formulation of Higgs Effective Field Theory: Measuring the Curvature of Scalar Field Space*, *Phys. Lett. B* **754** (2016) 335 [[arXiv:1511.00724](#)] [[INSPIRE](#)].
- [22] R. Alonso, E.E. Jenkins and A.V. Manohar, *Geometry of the Scalar Sector*, *JHEP* **08** (2016) 101 [[arXiv:1605.03602](#)] [[INSPIRE](#)].
- [23] G. Buchalla et al., *Complete One-Loop Renormalization of the Higgs-Electroweak Chiral Lagrangian*, *Nucl. Phys. B* **928** (2018) 93 [[arXiv:1710.06412](#)] [[INSPIRE](#)].
- [24] R. Alonso, K. Kanshin and S. Saa, *Renormalization group evolution of Higgs effective field theory*, *Phys. Rev. D* **97** (2018) 035010 [[arXiv:1710.06848](#)] [[INSPIRE](#)].
- [25] J. de Blas, O. Eberhardt and C. Krause, *Current and Future Constraints on Higgs Couplings in the Nonlinear Effective Theory*, *JHEP* **07** (2018) 048 [[arXiv:1803.00939](#)] [[INSPIRE](#)].
- [26] A. Falkowski and R. Rattazzi, *Which EFT*, *JHEP* **10** (2019) 255 [[arXiv:1902.05936](#)] [[INSPIRE](#)].
- [27] T. Cohen, N. Craig, X. Lu and D. Sutherland, *Is SMEFT Enough?*, *JHEP* **03** (2021) 237 [[arXiv:2008.08597](#)] [[INSPIRE](#)].
- [28] I. Banta et al., *Non-decoupling new particles*, *JHEP* **02** (2022) 029 [[arXiv:2110.02967](#)] [[INSPIRE](#)].
- [29] H. Elvang, D.Z. Freedman and M. Kiermaier, *A simple approach to counterterms in $N = 8$ supergravity*, *JHEP* **11** (2010) 016 [[arXiv:1003.5018](#)] [[INSPIRE](#)].

- [30] C. Cheung et al., *A Periodic Table of Effective Field Theories*, *JHEP* **02** (2017) 020 [[arXiv:1611.03137](#)] [[INSPIRE](#)].
- [31] Y. Shadmi and Y. Weiss, *Effective Field Theory Amplitudes the On-Shell Way: Scalar and Vector Couplings to Gluons*, *JHEP* **02** (2019) 165 [[arXiv:1809.09644](#)] [[INSPIRE](#)].
- [32] R. Aoude and C.S. Machado, *The Rise of SMEFT On-shell Amplitudes*, *JHEP* **12** (2019) 058 [[arXiv:1905.11433](#)] [[INSPIRE](#)].
- [33] G. Durieux, T. Kitahara, Y. Shadmi and Y. Weiss, *The electroweak effective field theory from on-shell amplitudes*, *JHEP* **01** (2020) 119 [[arXiv:1909.10551](#)] [[INSPIRE](#)].
- [34] T. Ma, J. Shu and M.-L. Xiao, *Standard model effective field theory from on-shell amplitudes**, *Chin. Phys. C* **47** (2023) 023105 [[arXiv:1902.06752](#)] [[INSPIRE](#)].
- [35] I. Low and Z. Yin, *Soft Bootstrap and Effective Field Theories*, *JHEP* **11** (2019) 078 [[arXiv:1904.12859](#)] [[INSPIRE](#)].
- [36] I. Low and Z. Yin, *New Flavor-Kinematics Dualities and Extensions of Nonlinear Sigma Models*, *Phys. Lett. B* **807** (2020) 135544 [[arXiv:1911.08490](#)] [[INSPIRE](#)].
- [37] A. Falkowski, G. Isabella and C.S. Machado, *On-shell effective theory for higher-spin dark matter*, *SciPost Phys.* **10** (2021) 101 [[arXiv:2011.05339](#)] [[INSPIRE](#)].
- [38] M. Accettulli Huber and S. De Angelis, *Standard Model EFTs via on-shell methods*, *JHEP* **11** (2021) 221 [[arXiv:2108.03669](#)] [[INSPIRE](#)].
- [39] H.-L. Li et al., *Complete set of dimension-eight operators in the standard model effective field theory*, *Phys. Rev. D* **104** (2021) 015026 [[arXiv:2005.00008](#)] [[INSPIRE](#)].
- [40] H.-L. Li et al., *Complete set of dimension-nine operators in the standard model effective field theory*, *Phys. Rev. D* **104** (2021) 015025 [[arXiv:2007.07899](#)] [[INSPIRE](#)].
- [41] H. Sun, M.-L. Xiao and J.-H. Yu, *Complete NLO operators in the Higgs effective field theory*, *JHEP* **05** (2023) 043 [[arXiv:2206.07722](#)] [[INSPIRE](#)].
- [42] I. Low, J. Shu, M.-L. Xiao and Y.-H. Zheng, *Amplitude/operator basis in chiral perturbation theory*, *JHEP* **01** (2023) 031 [[arXiv:2209.00198](#)] [[INSPIRE](#)].
- [43] N. Arkani-Hamed, T.-C. Huang and Y.-T. Huang, *Scattering amplitudes for all masses and spins*, *JHEP* **11** (2021) 070 [[arXiv:1709.04891](#)] [[INSPIRE](#)].
- [44] N. Christensen and B. Field, *Constructive standard model*, *Phys. Rev. D* **98** (2018) 016014 [[arXiv:1802.00448](#)] [[INSPIRE](#)].
- [45] B. Bachu and A. Yellespur, *On-Shell Electroweak Sector and the Higgs Mechanism*, *JHEP* **08** (2020) 039 [[arXiv:1912.04334](#)] [[INSPIRE](#)].
- [46] N. Christensen, B. Field, A. Moore and S. Pinto, *Two-, three-, and four-body decays in the constructive standard model*, *Phys. Rev. D* **101** (2020) 065019 [[arXiv:1909.09164](#)] [[INSPIRE](#)].
- [47] R. Balkin et al., *On-shell Higgsing for EFTs*, *JHEP* **03** (2022) 129 [[arXiv:2112.09688](#)] [[INSPIRE](#)].
- [48] S. De Angelis, *Amplitude bases in generic EFTs*, *JHEP* **08** (2022) 299 [[arXiv:2202.02681](#)] [[INSPIRE](#)].
- [49] N. Christensen et al., *Challenges with Internal Photons in Constructive QED*, [arXiv:2209.15018](#) [[INSPIRE](#)].

- [50] C. Cheung and Z. Moss, *Symmetry and Unification from Soft Theorems and Unitarity*, *JHEP* **05** (2021) 161 [[arXiv:2012.13076](#)] [[INSPIRE](#)].
- [51] D. Liu and Z. Yin, *Gauge invariance from on-shell massive amplitudes and tree-level unitarity*, *Phys. Rev. D* **106** (2022) 076003 [[arXiv:2204.13119](#)] [[INSPIRE](#)].
- [52] C. Cheung, A. Helset and J. Parra-Martinez, *Geometric soft theorems*, *JHEP* **04** (2022) 011 [[arXiv:2111.03045](#)] [[INSPIRE](#)].
- [53] G. Durieux and C.S. Machado, *Enumerating higher-dimensional operators with on-shell amplitudes*, *Phys. Rev. D* **101** (2020) 095021 [[arXiv:1912.08827](#)] [[INSPIRE](#)].
- [54] B. Henning and T. Melia, *Constructing effective field theories via their harmonics*, *Phys. Rev. D* **100** (2019) 016015 [[arXiv:1902.06754](#)] [[INSPIRE](#)].
- [55] G. Durieux et al., *Constructing massive on-shell contact terms*, *JHEP* **12** (2020) 175 [[arXiv:2008.09652](#)] [[INSPIRE](#)].
- [56] Z.-Y. Dong, T. Ma and J. Shu, *Constructing on-shell operator basis for all masses and spins*, [arXiv:2103.15837](#) [[INSPIRE](#)].
- [57] Z.-Y. Dong, T. Ma, J. Shu and Y.-H. Zheng, *Constructing generic effective field theory for all masses and spins*, *Phys. Rev. D* **106** (2022) 116010 [[arXiv:2202.08350](#)] [[INSPIRE](#)].
- [58] H. Sun, M.-L. Xiao and J.-H. Yu, *Complete NNLO operator bases in Higgs effective field theory*, *JHEP* **04** (2023) 086 [[arXiv:2210.14939](#)] [[INSPIRE](#)].
- [59] Z.-Y. Dong, T. Ma, J. Shu and Z.-Z. Zhou, *The New Formulation of Higgs Effective Field Theory*, [arXiv:2211.16515](#) [[INSPIRE](#)].
- [60] S. Chang, M. Chen, D. Liu and M.A. Luty, *Primary Observables for Indirect Searches at Colliders*, [arXiv:2212.06215](#) [[INSPIRE](#)].
- [61] L. Lehman and A. Martin, *Low-derivative operators of the Standard Model effective field theory via Hilbert series methods*, *JHEP* **02** (2016) 081 [[arXiv:1510.00372](#)] [[INSPIRE](#)].
- [62] B. Henning, X. Lu, T. Melia and H. Murayama, *2, 84, 30, 993, 560, 15456, 11962, 261485, ...: Higher dimension operators in the SM EFT*, *JHEP* **08** (2017) 016 [Erratum *ibid.* **09** (2019) 019] [[arXiv:1512.03433](#)] [[INSPIRE](#)].
- [63] L. Gráf et al., *Hilbert series, the Higgs mechanism, and HEFT*, *JHEP* **02** (2023) 064 [[arXiv:2211.06275](#)] [[INSPIRE](#)].
- [64] F.A. Berends et al., *Single Bremsstrahlung Processes in Gauge Theories*, *Phys. Lett. B* **103** (1981) 124 [[INSPIRE](#)].
- [65] A. Dedes et al., *Feynman rules for the Standard Model Effective Field Theory in R_ξ -gauges*, *JHEP* **06** (2017) 143 [[arXiv:1704.03888](#)] [[INSPIRE](#)].
- [66] N. Craig, H. Elvang, M. Kiermaier and T. Slatyer, *Massive amplitudes on the Coulomb branch of $N = 4$ SYM*, *JHEP* **12** (2011) 097 [[arXiv:1104.2050](#)] [[INSPIRE](#)].