



Quasi-integrability of deformations of the KdV equation

F. ter Braak^a, L.A. Ferreira^{b,*}, W.J. Zakrzewski^a

^a Department of Mathematical Sciences, University of Durham, Durham DH1 3LE, UK

^b Instituto de Física de São Carlos, IFSC/USP, Universidade de São Paulo, Caixa Postal 369, CEP 13560-970, São Carlos-SP, Brazil

Received 30 October 2017; received in revised form 26 November 2018; accepted 6 December 2018

Available online 10 December 2018

Editor: Hubert Saleur

Abstract

We investigate the quasi-integrability properties of various deformations of the Korteweg–de Vries (KdV) equation, depending on two parameters ε_1 and ε_2 , which include among them the regularized long-wave (RLW) and modified regularized long-wave (mRLW) equations. We show, using analytical and numerical methods, that the charges, constructed from a deformation of the zero curvature equation for the KdV equation, are asymptotically conserved for various values of the deformation parameters. By this we mean that, despite the fact that the charges do vary in time during the scattering of solitons, they return after the scattering to the same values they had before it. This property was tested numerically for the scattering of two and three solitons, and analytically for the scattering of two solitons in the mRLW theory ($\varepsilon_2 = \varepsilon_1 = 1$). In addition we show that for any values of ε_1 and ε_2 the Hirota method leads to analytical one-soliton solutions of our deformed equation but for $\varepsilon_1 \neq 1$ such solutions have the dispersion relation which depends on the parameter ε_1 . We also discuss some properties of soliton-radiation interactions seen in some of our simulations.

© 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The objective of the present paper is to study the quasi-integrability properties of deformations of the Korteweg–de Vries (KdV) equation [9,23] that include as particular cases the so-called reg-

* Corresponding author.

E-mail address: laf@ifsc.usp.br (L.A. Ferreira).

ularized long-wave equation (RLW), proposed by Peregrine [36] and T.B. Benjamin, J.L. Bona and J.J. Mahoney [5], and also the modified regularized long-wave equation (mRLW) introduced and studied by J.D. Gibbon, J.C. Eilbeck and R.K. Dodd [19]. Concretely, the model we consider involves a scalar field u satisfying the equation

$$u_t + u_x + \left[\frac{\alpha}{2} u^2 + \varepsilon_2 \frac{\alpha}{4} w_x v_t + u_{xx} - \varepsilon_1 (u_{xt} + u_{xx}) \right]_x = 0. \quad (1.1)$$

In which ε_1 , ε_2 , and α are real parameters, and where

$$u = w_t = v_x. \quad (1.2)$$

Note that the integrable KdV equation [9,23], corresponds to the case where the deformation parameters vanish, *i.e.* $\varepsilon_1 = \varepsilon_2 = 0$, and it is given by¹

$$u_t + u_x + \left[\frac{\alpha}{2} u^2 + u_{xx} \right]_x = 0. \quad (1.3)$$

As is well known, this equation describes nonlinear waves in shallow water traveling in the positive direction of the x -axis only. If one considers the linearization of the KdV equation one finds that its traveling wave solutions satisfy a dispersion relation of the form $\omega = k - k^3$, and so the phase velocity $\omega/k = 1 - k^2$, and group velocity $\frac{d\omega}{dk} = 1 - 3k^2$, which become negative, and in fact unbounded, for large enough k .

Motivated by this fact, Peregrine [36] and T.B. Benjamin, et al. [5] proposed the so-called regularized long wave equation (RLW)

$$u_t + u_x + \left(\frac{\alpha}{2} u^2 - u_{xt} \right)_x = 0, \quad (1.4)$$

which corresponds to (1.1) for the case $\varepsilon_1 = 1$ and $\varepsilon_2 = 0$. The advantage of the RLW equation over the KdV equation is that the RLW equation yields a dispersion relation of the form $\omega = k/(1 + k^2)$, and so a phase velocity that is bounded and tends to zero for short wavelengths. Its disadvantage is that the RLW equation is not integrable and that it possesses only one analytical solution, namely the one-soliton solution. The two and three-soliton solutions for RLW are only known numerically and were constructed by Eilbeck and McGuire [12,13].

The mRLW equation, introduced by Gibbon, Eilbeck and Dodd [19], can be written in terms of the u field as

$$u_t + u_x + \left[\frac{\alpha}{2} u^2 + \frac{\alpha}{4} w_x v_t - u_{xt} \right]_x = 0 \quad (1.5)$$

and so it corresponds to (1.1) for the case $\varepsilon_1 = \varepsilon_2 = 1$. The linearized version of such an equation has the same dispersion relation as the RLW equation, and the same exact analytical one-soliton solution as RLW. The remarkable property of the mRLW equation, however, is that it also possesses analytic two-soliton solutions, even though it is not integrable in the sense of possessing an infinite number of conserved quantities.

The analytical one-soliton solution for the RLW equation, and the analytical one- and two-soliton solutions for the mRLW can be constructed using the Hirota direct method where the relation between the τ -function and the u -field is of the form $u \sim -(\ln \tau)_{,xt}$. Therefore, by

¹ The KdV equation as presented here is not in its standard form, but one can perform a change of variables to get the standard form of the KdV equation (see, for instance, the notes by M. Dunajski for more information [11]). The same applies for the RLW and mRLW equations. In fact, to get the original notation used by the authors one should choose $\alpha = 12$, in (1.4), and $\alpha = 8$ in (1.5).

changing the τ -function as $\tau \rightarrow f(x) g(t) \tau$, does not change the solution for the u -field. So, integrating (1.2) one gets that $w \sim -(\ln \tau)_x + h(x)$, and $v \sim -(\ln \tau)_t + j(t)$. Consequently, as long as Hirota’s solutions are concerned the integration “constants” $h(x)$ and $j(t)$, can be reabsorbed in the redefinition of the τ -function. Therefore, when constructing the soliton solutions, either analytically or numerically, it is convenient to work with the field q defined as

$$u \equiv -\frac{8}{\alpha} q_{xt}. \tag{1.6}$$

Dropping the integration “constants” as explained above one can then write $w_x = -\frac{8}{\alpha} q_{xx}$ and $v_t = -\frac{8}{\alpha} q_{tt}$. Replacing the u -field by the q -field into (1.1) one gets an equation for q which can in fact be written as the x -derivative of the equation

$$q_{tt} + q_{xt} - 4q_{xt}^2 - 2\varepsilon_2 q_{xx} q_{tt} + q_{xxx} - \varepsilon_1 (q_{xxt} + q_{xxt}) = 0. \tag{1.7}$$

Therefore, any solution of (1.7) leads to a solution of (1.1), in which the integration “constants” in w and v have been absorbed as explained above. So in this paper we base our discussion on the study of (1.7).

First, we show in this paper that (1.7) admits at least two types of analytical Hirota one-solitons. The first one is obtained in section 4 by the Hirota’s method and it is given by

$$q_{\text{type 1}} = \frac{3}{(2 + \varepsilon_2)(1 + (1 - \varepsilon_1)k^2)} \left[\ln 2 + \frac{\Gamma}{2} + \ln \cosh \left(\frac{\Gamma}{2} \right) \right], \tag{1.8}$$

with

$$\Gamma = kx - \omega_1 t + \delta; \quad \omega_1 = \frac{k + (1 - \varepsilon_1)k^3}{1 - \varepsilon_1 k^2}. \tag{1.9}$$

The second type is constructed in Appendix A by a direct method and it is given by

$$q_{\text{type 2}} = \frac{3[1 + (\varepsilon_1 - 1)k^2]}{(2 + \varepsilon_2)} \left[\ln \left(\cosh \left(\frac{\xi}{2\sqrt{1 + (\varepsilon_1 - 1)k^2}} \right) \right) + b\xi + c \right] \tag{1.10}$$

with

$$\xi = kx - \omega_2 t + \delta; \quad \omega_2 = \frac{k}{1 - k^2} \tag{1.11}$$

where b and c are arbitrary parameters, and where we have to assume that $1 + (\varepsilon_1 - 1)k^2 > 0$, for the solution to be real. In terms of the u -field these solutions take the form

$$u_{\text{type 1}} = \frac{6}{\alpha} \frac{k^2}{(2 + \varepsilon_2)(1 - \varepsilon_1 k^2)} \operatorname{sech}^2 \left[\frac{1}{2} (kx - \omega_1 t + \delta) \right] \tag{1.12}$$

and

$$u_{\text{type 2}} = \frac{6}{\alpha} \frac{k^2}{(2 + \varepsilon_2)(1 - k^2)} \operatorname{sech}^2 \left[\frac{kx - \omega_2 t + \delta}{2\sqrt{1 + (\varepsilon_1 - 1)k^2}} \right]. \tag{1.13}$$

Clearly they become the same for the case where $\varepsilon_1 = 1$.

Note that (1.12) and (1.13) constitute two-parameter families of one-soliton solutions, labeled by the deformation parameters ε_1 and ε_2 . They interpolate between the one-soliton solution of the RLW model (for $\varepsilon_1 = 1, \varepsilon_2 = 0$ and $\alpha = 12$) constructed in [5,36], and the one-soliton solution of the mRLW model (for $\varepsilon_1 = \varepsilon_2 = 1$ and $\alpha = 8$) in [19]. Note also that, as pointed out

in [19], the expressions for the one-soliton solutions of the RLW and mRLW, indeed look the same for the u -field, when the rescaling parameter α is chosen as above, *i.e.* $\alpha = 12$ for RLW and $\alpha = 8$ for mRLW.

The analytical Hirota two-soliton solution, however, exists only for the case $\varepsilon_1 = \varepsilon_2 = 1$, and it is the solution constructed in [19] and given by

$$q = \ln(1 + e^{\Gamma_1} + e^{\Gamma_2} + A_{12} e^{\Gamma_1 + \Gamma_2}) ; \quad \varepsilon_1 = \varepsilon_2 = 1, \quad (1.14)$$

where

$$\Gamma_i = k_i x - \omega_i t + \delta_i, \quad \omega_i = \frac{k_i}{1 - k_i^2}, \quad i = 1, 2, \quad (1.15)$$

and

$$A_{12} = - \frac{(\omega_1 - \omega_2)^2 (k_1 - k_2)^2 + (\omega_1 - \omega_2)(k_1 - k_2) - (\omega_1 - \omega_2)^2}{(\omega_1 + \omega_2)^2 (k_1 + k_2)^2 + (\omega_1 + \omega_2)(k_1 + k_2) - (\omega_1 + \omega_2)^2}. \quad (1.16)$$

The previous paper by two of us (FtB and WJZ) [6], looked at the scattering properties of various soliton-like configurations of the modified regularized long-wave (mRLW) equation, originally introduced and studied by J.D. Gibbon, J.C. Eilbeck and R.K. Dodd [19], and given by equation (1.7) for $\varepsilon_1 = \varepsilon_2 = 1$. As stated in the aforementioned paper, the three soliton-solutions are not known and, in fact, they cannot be found using Hirota's method.

The main results of the previous paper [6] involved the observation that the scattering of two initially well-separated solitons, which, in fact, is described by an exact solution of the equations of motion (see equation (1.14)), could be approximated very well by the numerical time-evolution of a linear superposition of two one-soliton solutions. In fact, the approximation was so good that it was virtually impossible to see any difference between them. Then [6] also extended this idea to the simulation of various three-soliton interactions by numerically evolving linear superpositions of three (initially well-separated) single-soliton solutions. Such scattering were also found to be very elastic (in the sense that the solitons preserved their shapes and velocities and there was no visible loss of radiation). Moreover, the phase shifts in the scattering of three solitons were, again, very well approximated by the sums of the successive two two-soliton scattering. This property holds for integrable models and the fact that it holds also for the mRLW equation, which is not an Hirota integrable system, suggests that the mRLW equation may be 'close' to being integrable.

To test this further we have decided to extend the investigations presented in the previous paper by considering this model as a perturbation of the integrable KdV equation since then we can look at the conserved quantities of the integrable model and see how they change when we perturb this model to become the mRLW model, or for that matter any other model 'nearby'.

We discuss all of this in the context of quasi-integrability, a concept introduced in [14], and developed further, for other models, in [3,15–18]. As it is well known, in a theory in $(1 + 1)$ -dimensions, a soliton [39] is a classical solution of its equations of motion that travels with constant velocity and without dissipation of energy. But more importantly, when two or more solitons are scattered they do not destroy each other, and the only effect of their scattering is a shift in their positions relative to those they would have had, had the scattering not had occurred. A well accepted explanation for such a behaviour is that soliton theories possess an infinite number of exactly conserved charges that constrain the dynamics in such a way that the solitons have no option but to retain their shapes and velocities after the scattering. Such theories are, in fact, exactly integrable admitting a zero curvature representation of their equations of motion, *i.e.* a Lax–Zakharov–Shabat equation [25].

A few years ago, having performed many numerical simulations, we noticed that some deformations of exactly integrable soliton theories, like the sine-Gordon model, possess solutions that approximately behave like solitons and this observation has led us to propose the concept of quasi-integrability, which can be formulated in the following form: We call a theory quasi-integrable if it admits a representation of their equations of motion in terms of an anomalous zero curvature condition, which in turn, leads to an infinite number of anomalous conservation laws. In addition, the theory has to possess one-soliton like solutions that travel with constant velocity without the dissipation of energy. Therefore, the conservation of energy is not anomalous but exact. When two or more of these one-soliton like solutions scatter the anomalous charges (and there is an infinite number of them) are conserved asymptotically, *i.e.* they do vary in time during the scattering process, and sometimes quite a lot, but they all return, after the scattering, to the values they had before the scattering took place. The asymptotic conservation of such an infinite number of charges has the same effect, as in true soliton theories, of constraining the dynamics in such a way that the soliton like configurations leave the scattering process basically as they have entered it. We have tested these ideas, analytically and numerically, in several deformations of exact integrable soliton theories and, within the limitations of our methods, they seem to have worked [3,14–18].

The dynamical mechanism responsible for such a behaviour of these quasi-solitons is far from having been understood yet. We have observed, however, some patterns that might be relevant for the study of such a nonlinear phenomenon. First of all, the anomalies of the conservation laws vanish for the one-soliton like solutions, and so the one-soliton sectors of these theories do have an infinite number of exact conservation laws. Secondly, the anomalies also vanish for configurations in which one-soliton like solutions are well separated from each other. Moreover, the anomalies are non-zero only when the solitons are close together and interact with each other. Thirdly, the observed phenomenon seems to occur when the multi-soliton solutions of the equation of motion possess very specific symmetry properties under a space–time parity transformation which can be described as follows. In the situations when the phenomenon occurs the two or three-soliton configurations are eigenstates, either odd or even, of an inversion transformation of the space and time coordinates around a point in space–time that depends on the parameters of the solutions, like speed, width, localization, *etc.* The quasi-integrability seems to occur when the densities of the anomalies of the quasi-conserved charges are odd under this parity transformation. Thus, integrating these densities in a rectangle in space–time centered at the particular point around which the inversion is performed, one gets zero. In consequence, the charges possess a mirror-like symmetry in the sense that they have the same values at times equidistant from this particular point in the time axis, around which the parity transformation is performed. So, we have not only an asymptotic conservation of the charges but also a mirror symmetry. We do not know whether such a parity property is the cause of quasi-integrability, but it seems to be present whenever the theory is quasi-integrable.

The densities of the quasi-conserved charges are in general polynomials in the derivatives of the fields, and so such densities vanish away from the core of the solitons. Therefore, when the solitons are well separated, like in the “in states” of a scattering process, the value of any one of the infinite number of charges should be the sum of the charges associated with each individual soliton. After the scattering, when the solitons are well separated again, the values of these charges should become again the sum of the charges of each “out soliton”. Consequently, one is led to believe that the asymptotic conservation of our infinite number of quasi-charges is quite reasonable. However, for that to be the case the solitons have to leave the scattering, more or less, in the same configurations that they have entered it, *i.e.* possessing the same speeds,

widths, shape, *etc.* The more reasonable thing to expect is that they could radiate some energy away, change their shapes and velocities, *etc.* This is the mystery still to be understood. What is the dynamical mechanism that drives the solitons to remain the same after the scattering?

The answer to this question is not known but there are many results in the literature that are related to it. First of all, there is the seminal paper of Zabusky and Kruskal [39], that has, in fact, introduced the word soliton, in which the authors described their discovery that if one takes a sinusoidal wave as an initial configuration and then evolves in a discretized version of the KdV equation, one finds that instead of the wave crest breaking, like sea waves, the effect of the non-linearity modifies it to evolve into a train of KdV solitons that continue to propagate, overtaking one another and interacting, and not destroying each other. So, the solitons in the integrable KdV model seem to be the “fixed points” of the dynamics and describe the states ‘where the system prefers to be’, like natural normal modes of linear systems. Since Zabusky and Kruskal’s paper [39] a lot of other dispersive equations have been studied, including non-integrable ones, and it has been found that the above described behaviour seems to be quite common in nonlinear dispersive partial differential equations. This large number of studies is so robust that it has given rise to a conjecture, the so-called *Soliton Resolution Conjecture*. This conjecture basically says that with a generic data as initial condition the nonlinear systems, integrable or not, should evolve into a finite number of solitons plus some radiation. The conjecture, itself, has not been proved yet and it is not even easy to formulate it more rigorously. For a more concise discussion on this topic see [26,38].²

The other concept that might be important to what we observe in quasi-integrable theories is the notion of orbital asymptotic stability, or simply *asymptotic stability* [34,35], that has, in fact, evolved from *orbital stability* put forward by Benjamin [4]. Here one does not look at several solitons, but just at one solitary wave. One takes a given solitary wave solution u_{sol} of the equation under consideration, and considers the orbit of solutions $g u_{\text{sol}}$ of the equation, under the group G of symmetries of this equation, with $g \in G$. The solitary wave is called *orbitally asymptotic stable* if a small deviation of $g u_{\text{sol}}$ from the initial data implies that there exists a $g(t) \in G$, such that $g(t) \rightarrow g_{\infty}$, and the solution converges to $g_{\infty} u_{\text{sol}}$, in some norm, as $t \rightarrow \infty$ [34]. There is a vast literature about asymptotic stability, in particular, on the RLW equation (1.4). It is also worth mentioning the concept of asymptotic stability in the sense of the asymptotic expansion of the PDE up to the first nonlinear correction [8,22]. Such a concept might also be relevant to quasi-integrability.

The stability of the RLW solitary waves has been considered, for example, in [29,30]. In addition the concept of asymptotic stability has been extended to two-soliton-like solutions and in the case of RLW some particular two-soliton configurations have been discussed and shown to describe inelastic or quasi-elastic scattering in [27,28]. Such results certainly fit together with what we describe in the present paper for the RLW equation (1.4), and in some sense also for the more general deformations of the KdV equations given by equation (1.1). If solitary waves can be shown to be asymptotically stable then one could intuitively (and naively) think that if the deformations induced by the scattering of two of them are not too large then, when these waves separate after the scattering and propagate without influencing each other, they would asymptotically converge to solitary waves again. Of course, this is only a picture of the phenomenon but

² We are very grateful to the referee for drawing our attention to this Soliton Resolution Conjecture and to the notion of Asymptotic Stability that we discuss next.

certainly such possible connections between asymptotic stability and quasi-integrability deserve further studies.

The paper is organized as follows: In the next section we introduce our Lax–Zakharov–Shabat equation which we then use to study the quasi-integrability of the model described by equation (1.1) for various values of ε_1 and ε_2 . We perform the usual gauge transformations and finally define the charges which are truly conserved when $\varepsilon_1 = \varepsilon_2 = 0$ (i.e., for the KdV model). In section 3 we introduce the parity argument for the travelling wave solutions of (1.1) and show that if the multi-soliton field configurations of the model possess this symmetry, then the charges are quasi-conserved. The next section discusses the Hirota method of finding solutions of some non-linear equations and applying it to our equation (1.1). In it we show that this method gives a one soliton solution for any ε_2 with $\varepsilon_1 = 1$ being constant. This is the case for the usual dispersion relation ($\omega = \frac{k}{1-k^2}$), and for $\varepsilon_1 \neq 1$ the dispersion relation has to be modified. However, if we insist on the usual dispersion relation the Hirota method does not work but a one-soliton solution can still be found.³ We discuss a construction of such a solution in the appendix A.

The next two sections discuss the analytical quasi-integrability of the mRLW model and present arguments for the integrability based on the evenness parity properties of the multi-soliton functions. Section 7 discusses the soliton solutions of the KdV equation, obtained via the Hirota method, and the parity properties of two- and three-soliton solutions of this model. The lengthy section 8 presents some results of our numerical simulations. These simulations were performed using a specially constructed numerical program based on implicit and explicit methods of solving (1.7). The calculations involved studying the time evolution of various field configurations initially corresponding to two- or three-soliton systems and then checking whether the observed results supported quasi-integrability of the model for various values of ε_1 and ε_2 . We finish the paper with our conclusions and a few short appendices presenting more information about our numerical techniques and providing some details on the construction of conserved or quasi-conserved quantities discussed in section 2 of the paper.

2. The anomalous Lax–Zakharov–Shabat equation

Our motivation is to study equation (1.1) in the context of quasi-integrability as a deformation of the integrable KdV equation expressed by equation (1.3). Of course, there is a vast literature on deformations of integrable theories (see for instance [2,21]), using different techniques. Here we work with the concept of anomalous zero curvature equations. To this end, we start by constructing the quasi-zero curvature equation, and subsequently produce the quasi-conserved charges from it. To do this we introduce the Lax potentials A_x and A_t , as

$$A_x = - \left[b_1 - \frac{\alpha}{12} u (b_{-1} - F_{-1}) \right], \tag{2.1}$$

$$A_t = - \left[-4b_3 - b_1 + \frac{\alpha}{6} u_x F_0 - \frac{\alpha}{3} u F_1 + \frac{G}{2} (b_{-1} - F_{-1}) \right], \tag{2.2}$$

where the generators b_{2n+1} and F_n , are defined in the appendix C, and where

$$G \equiv \frac{\alpha^2}{18} u^2 + \frac{\alpha^2}{24} \varepsilon_2 w_x v_t - \frac{\alpha}{6} \varepsilon_1 u_{xt} + \frac{\alpha}{6} (1 - \varepsilon_1) u_{xx} + \frac{\alpha}{6} u. \tag{2.3}$$

The curvature of the Lax connection is given by

³ We thank the referee for drawing our attention to this possibility.

$$F_{tx} = \partial_t A_x - \partial_x A_t + [A_t, A_x] = -X F_0 + \frac{\alpha}{12} Y (b_{-1} - F_{-1}), \quad (2.4)$$

where $Y = 0$ corresponds to the equation of motion of our deformed model (1.1)

$$Y \equiv u_t + u_x + \left[\frac{\alpha}{2} u^2 + \varepsilon_2 \frac{\alpha}{4} w_x v_t + u_{xx} - \varepsilon_1 (u_{xt} + u_{xx}) \right]_x, \quad (2.5)$$

and where

$$X \equiv \frac{\alpha}{6} \left[\frac{\alpha}{4} \varepsilon_2 w_x v_t - \varepsilon_1 (u_{xt} + u_{xx}) \right]. \quad (2.6)$$

Thus, if we now assume that u is a solution of the equations of motion (*i.e.*, $Y = 0$), then X represents the anomaly of the zero curvature equation. In the case of the KdV equation (*i.e.*, $\varepsilon_2 = \varepsilon_1 = 0$), the anomaly vanishes resulting in the well-known infinite number of truly conserved charges.

2.1. Quasi-conserved charges

In order to construct the quasi-conserved charges we follow the procedure discussed in [14], which is an adaptation to quasi-integrable theories, of the so-called abelianization procedure for exactly integrable theories [1,10,32]. The key ingredient is that the generator b_1 appearing in A_x in (2.1), is a semi-simple element of the loop algebra and so splits this algebra into its kernel and image under its adjoint action as explained in the appendix C. Then, we perform a gauge transformation to rotate the Lax potentials into an infinite abelian subalgebra of the $\mathfrak{sl}(2)$ algebra, generated by b_{2n+1} . The gauge transformation transforms the potentials as

$$A_x \rightarrow a_x = g A_x g^{-1} - (\partial_x g) g^{-1}, \quad (2.7)$$

$$A_t \rightarrow a_t = g A_t g^{-1} - (\partial_t g) g^{-1}, \quad (2.8)$$

where the group element g is chosen to be

$$g = \exp \left(\sum_{n=1}^{\infty} \zeta_n F_{-n} \right) \equiv \exp \left(\sum_{n=1}^{\infty} \mathcal{F}_{-n} \right), \quad (2.9)$$

where ζ_n are parameters to be determined (see below and appendix D), and F_{-n} are generators of the $\mathfrak{sl}(2)$ loop algebra (see appendix C).

Performing the expansion we find that a_x can be expressed as

$$\begin{aligned} a_x = & A_x + \left[\sum_{n=1}^{\infty} \mathcal{F}_{-n}, A_x \right] + \frac{1}{2!} \left[\sum_{n=1}^{\infty} \mathcal{F}_{-n}, \left[\sum_{m=1}^{\infty} \mathcal{F}_{-m}, A_x \right] \right] + \dots \\ & - \sum_{m=1}^{\infty} \partial_x \mathcal{F}_{-m} - \frac{1}{2!} \left[\sum_{n=1}^{\infty} \mathcal{F}_{-n}, \sum_{m=1}^{\infty} \partial_x \mathcal{F}_{-m} \right] \\ & - \frac{1}{3!} \left[\sum_{k=1}^{\infty} \mathcal{F}_{-k}, \left[\sum_{n=1}^{\infty} \mathcal{F}_{-n}, \sum_{m=1}^{\infty} \partial_x \mathcal{F}_{-m} \right] \right] + \dots \end{aligned} \quad (2.10)$$

Using the algebra defined by eqs. (C.5) to (C.10), we can write a_x as

$$\begin{aligned}
 a_x &= -b_1 \\
 &+ \zeta_1 [b_1, F_{-1}] \\
 &+ \zeta_2 [b_1, F_{-2}] + \frac{\alpha}{12} u (b_{-1} - F_{-1}) - \frac{1}{2} \zeta_1^2 [F_{-1}, [F_{-1}, b_1]] - \partial_x \zeta_1 F_{-1} \\
 &+ \dots,
 \end{aligned} \tag{2.11}$$

where we have written down all the terms of the same grade on the same line and the grading is defined by the grading operator d given in (C.11). Note that the parameter ζ_k , in the expansion (2.11), multiplying the commutator $[b_1, F_{-k}]$, and so first appears among the terms of grade $-k + 1$. Thus, one can choose the parameters ζ_k , recursively, to cancel the image part of a_x at the grade $-k + 1$, *i.e.* its component in the direction of F_{-k+1} . The expressions for the parameters ζ_k , obtained this way, are given in the appendix D, and they are polynomials in u and its x -derivatives. Having chosen the parameters ζ_k this way, the expression for a_x becomes

$$a_x = -b_1 + \sum_{n=0}^{\infty} a_x^{(-2n-1)} b_{-2n-1}, \tag{2.12}$$

where the first values of $a_x^{(-2n-1)}$ are given by

$$a_x^{(-1)} = \frac{\alpha}{2^2 3} u, \tag{2.13}$$

$$a_x^{(-3)} = \frac{\alpha^2}{2^5 3^2} u^2, \tag{2.14}$$

$$a_x^{(-5)} = \frac{\alpha^3}{2^7 3^3} u^3 + \frac{\alpha^2}{2^7 3^2} uu_{xx}, \tag{2.15}$$

$$a_x^{(-7)} = \frac{5\alpha^4}{2^{11} 3^4} u^4 + \frac{\alpha^3}{2^7 3^3} u^2 u_{xx} + \frac{1}{2^9 3^2} (\alpha^3 uu_x^2 + \alpha^2 uu_{xxx}). \tag{2.16}$$

Note that in our gauge transformation we have not used the equations of motion $Y = 0$, with Y given in (2.5). In the transformation (2.8) of the A_t component of the Lax connection, the group element g is already fixed, but we still can use the equations of motion $Y = 0$ to simplify it, *i.e.* we are performing an on-shell gauge transformation. The on-shell result is then given by

$$a_t = 4b_3 + b_1 + \sum_{n=0}^{\infty} a_t^{(-2n-1)} b_{-2n-1} + \sum_{n=-2}^{\infty} c_t^{(-n)} F_{-n}. \tag{2.17}$$

In the appendix D we give explicit expressions for the first few quantities $a_t^{(-2n-1)}$ and $c_t^{(-n)}$. Note that, due to the anomaly X , the quantities $c_t^{(-n)}$ do not vanish, and so the potential a_t is not really rotated to the abelian subalgebra generated by b_{2n+1} . That is a difference with respect to the case of exactly integrable theories, but it will not be a concern for us as we explain below.

The on-shell gauge-transformed curvature then becomes

$$F_{tx} \rightarrow f_{tx} = \partial_t a_x - \partial_x a_t + [a_t, a_x] = g F_{tx} g^{-1} = -X g F_0 g^{-1} \tag{2.18}$$

and so it takes the form

$$f_{tx} = -X \left(\sum_{n=0}^{\infty} \gamma^{(-2n-1)} b_{-2n-1} + \sum_{n=0}^{\infty} \beta^{(-n)} F_{-n} \right), \tag{2.19}$$

where we have assumed that the equations of motion are satisfied (*i.e.*, $Y = 0$), and where

$$\begin{aligned}
\gamma^{(-1)} &= 0, \\
\gamma^{(-3)} &= -\partial_x \left[\frac{\alpha}{2^3 3} u \right], \\
\gamma^{(-5)} &= -\partial_x \left[\frac{\alpha^2}{2^6 3} u^2 + \frac{\alpha}{2^5 3} u_{xx} \right], \\
\gamma^{(-7)} &= -\partial_x \left[\frac{5\alpha^3}{2^8 3^3} u^3 + \frac{5\alpha^2}{2^8 3^2} u_x^2 + \frac{5\alpha^2}{2^7 3^2} u_{xx} + \frac{\alpha}{2^7 3} u_{xxxx} \right].
\end{aligned} \tag{2.20}$$

In addition we have that $\beta^{(0)} = 1$, and $\beta^{(-m)} = 0$ for $m = 1, 2, 3, 4$. The first nonvanishing $\beta^{(i)}$ is $\beta^{(-5)}$, and it is given by

$$\beta^{(-5)} = \partial_x \left[\frac{\alpha^2}{2^7 3^2} u^2 \right]. \tag{2.21}$$

From the commutation relations eqs. (C.5) to (C.10) one can deduce that the commutators of a given element b_{2n+1} with any element of the algebra never produce an element of the abelian subalgebra generated by b_{2n+1} . Therefore, the commutator $[a_x, a_t]$ does not contain any terms in the direction of the b_{-2n-1} generators, since a_x lies in this abelian subalgebra. Thus, if we now consider only the terms in the direction of the b_{-2n-1} generators of the gauge transformed curvature, we find that

$$\partial_t a_x^{(-2n-1)} - \partial_x a_t^{(-2n-1)} = -X \gamma^{(-2n-1)}, \quad \forall n \in \mathbb{Z}_0^+, \tag{2.22}$$

which can be rewritten as

$$\frac{dQ^{(-2n-1)}}{dt} = a_t^{(-2n-1)} \Big|_{x=-\infty}^{\infty} + \alpha^{(-2n-1)}, \tag{2.23}$$

where

$$Q^{(-2n-1)} \equiv \int_{-\infty}^{\infty} dx a_x^{(-2n-1)} \quad \text{and} \quad \alpha^{(-2n-1)} \equiv - \int_{-\infty}^{\infty} dx X \gamma^{(-2n-1)}. \tag{2.24}$$

Since all the terms of the parameters ζ_n depend on u and its x -derivatives, and $u \rightarrow 0$ when $x \rightarrow \pm\infty$, we see that $g \rightarrow \mathbb{1}$ as $x \rightarrow \pm\infty$. This implies that

$$\lim_{x \rightarrow \pm\infty} a_t = \lim_{x \rightarrow \pm\infty} A_t \tag{2.25}$$

and so from (2.2), we get that

$$a_t^{(-2n-1)} \Big|_{x=-\infty}^{\infty} = 0. \tag{2.26}$$

Hence, equation (2.23) becomes

$$\frac{dQ^{(-2n-1)}}{dt} = \alpha^{(-2n-1)}, \tag{2.27}$$

and so the quantities $Q^{(-2n-1)}$ are candidates for our quasi-conserved charges.

It is important to point out that the Lax potential A_x , given in (2.1), does not depend upon the deformation parameters ε_1 and ε_2 , and so it is the same as the Lax potential for the integrable KdV equation. Therefore, the charge densities $a_x^{(-2n-1)}$ and, consequently, the charges $Q^{(-2n-1)}$, are the same as those for the KdV theory. The dependence upon the deformation parameters in

(2.27), comes only from the anomaly X . Furthermore, we note that the lowest of these charges is exactly conserved, *i.e.*

$$Q^{(-1)} = \frac{\alpha}{12} \int_{-\infty}^{\infty} dx u, \quad \frac{d Q^{(-1)}}{dt} = 0. \tag{2.28}$$

Indeed, from (2.20) one sees that $\gamma^{(-1)} = 0$, and this implies that $Q^{(-1)}$ is conserved and so u is a density of a conserved quantity for any value of $\varepsilon_1, \varepsilon_2$, and α .

The remaining charges are not properly conserved. However, as we show below, for some special soliton solutions such charges are quasi-conserved. By that we mean that in the process of a soliton scattering these charges do vary in time but, after the scattering, they all return to the same values they had prior to this scattering (*i.e.* the values before and after the scattering are the same). This property is not well understood yet, but we have found that it is accompanied by a space–time parity symmetry of the soliton solutions, and this is useful in trying to gain an understanding why the anomalies of the charges vanish when integrated over time during the whole scattering process. In section 3 we explain how this works for the case under consideration here.

3. The parity argument

Here we explain in detail how the quasi-integrability concept is related to the existence of some parity symmetries of the soliton solutions. Before we show how it works for some specific soliton solutions, let us present here a general argument for the deformations of the KdV theory given by the equation (1.1).

3.1. Parity argument for the traveling wave solutions

Consider a class of traveling wave solutions $u \equiv \bar{u}(x - \frac{\omega}{k}t + \delta)$ of the equation (1.1) which, of course, includes in it the one-soliton solutions as its particular cases. For a fixed value of the time t , we define the shifted space coordinate as

$$\bar{x} = x - \frac{\omega}{k}t + \delta \tag{3.1}$$

and introduce the space parity transformation

$$P_{\bar{x}} : \quad \bar{x} \rightarrow -\bar{x}. \tag{3.2}$$

The only hypothesis that we are making here is that the traveling wave solution is invariant under parity, *i.e.* that

$$P_{\bar{x}}(\bar{u}) = \bar{u}. \tag{3.3}$$

Since $\bar{u} = \bar{w}_t = \bar{v}_x$, it turns out that $(\partial_x = \partial_{\bar{x}})$

$$P_{\bar{x}}(\bar{w}) = \bar{w}, \quad P_{\bar{x}}(\bar{v}) = -\bar{v} \quad \text{and so} \quad P_{\bar{x}}(\bar{w}_x) = -\bar{w}_x, \quad P_{\bar{x}}(\bar{v}_t) = -\bar{v}_t. \tag{3.4}$$

In consequence, we see that $\bar{u}_{x\bar{t}} = -\omega \bar{u}_{\bar{x}\bar{x}}$ and so $P_{\bar{x}}(\bar{u}_{x\bar{t}}) = \bar{u}_{x\bar{t}}$. Thus, the anomaly X given in (2.6) and evaluated on such type of solutions, satisfies

$$P_{\bar{x}}(X) = X. \tag{3.5}$$

Note that the anomalies coefficients, γ 's given in (2.20), always involve an odd number of x -derivatives of the field u , and so

$$P_{\tilde{x}} \left(\gamma^{(-2n-1)} \right) = -\gamma^{(-2n-1)}. \tag{3.6}$$

Thus we see that

$$\int_{-\infty}^{\infty} dx X \gamma^{(-2n-1)} = \int_{-\infty}^{\infty} d\tilde{x} X \gamma^{(-2n-1)} = 0 \tag{3.7}$$

and so the charges (2.24) are exactly conserved for all traveling wave solutions, and so also for the one-soliton solutions.

3.2. Parity argument for multi-soliton fields

Now we consider the space–time parity transformation around a given point (x_{Δ}, t_{Δ}) , of space–time

$$P : \quad (\tilde{x}, \tilde{t}) \rightarrow (-\tilde{x}, -\tilde{t}) ; \quad \tilde{x} = x - x_{\Delta} \quad \tilde{t} = t - t_{\Delta}. \tag{3.8}$$

Again we make the hypothesis that the u -field evaluated on the soliton solution we are considering, that now we denote u_s , is even under such a parity transformation, *i.e.*

$$P(u_s) = u_s, \tag{3.9}$$

where u_s in (3.9) is evaluated on that particular soliton solution. In addition, since $w_s \equiv \int dt u_s$, and $v_s \equiv \int dx u_s$ (see (1.2)), we see that

$$P(w_s) = -w_s, \quad \text{and} \quad P(v_s) = -v_s, \tag{3.10}$$

where w_s and v_s are respectively the fields w and v evaluated on that soliton solution. From (2.3), (2.5) and (2.6) we have that

$$P(G) = G, \quad P(Y) = -Y, \quad P(X) = X. \tag{3.11}$$

Next we consider the following order 2 automorphism (*i.e.* $\sigma^2 = 1$) of the $\mathfrak{sl}(2)$ loop algebra

$$\sigma(T) = e^{i\pi d} T e^{-i\pi d}, \tag{3.12}$$

where d is the grading operator defined in (C.11). Then,

$$\sigma(b_{2n+1}) = -b_{2n+1}, \quad \sigma(F_{2n+1}) = -F_{2n+1}, \quad \sigma(F_{2n}) = F_{2n}. \tag{3.13}$$

We also consider the combination of these two operations, the parity and the automorphism:

$$\Omega \equiv P \sigma. \tag{3.14}$$

One can check that the Lax operators (2.1) and (2.2), when evaluated on soliton solutions satisfying (3.9) and (3.10), satisfy

$$\Omega(A_{\mu}) = -A_{\mu}, \quad \mu = x, t, \tag{3.15}$$

and so the curvature (2.4) satisfies

$$\Omega(F_{tx}) = F_{tx}. \tag{3.16}$$

Let us rewrite (2.12) as

$$a_x = \sum_{n=-1}^{\infty} \hat{a}_x^{(-2n-1)}; \quad \text{where } \hat{a}_x^{(1)} = -b_1; \quad \text{and}$$

$$\hat{a}_x^{(-2n-1)} = a_x^{(-2n-1)} b_{-2n-1}. \tag{3.17}$$

Since $\Omega(b_1) = -b_1$, it follows that

$$\Omega([b_1, \mathcal{F}_{-n}]) = -[b_1, \Omega(\mathcal{F}_{-n})], \tag{3.18}$$

where \mathcal{F}_{-n} is defined in (2.9), and so

$$(1 + \Omega)([b_1, \mathcal{F}_{-n}]) = [b_1, (1 - \Omega)(\mathcal{F}_{-n})]. \tag{3.19}$$

From (2.11) we see that $\hat{a}_x^{(0)} = [b_1, \mathcal{F}_{-1}]$, and so

$$(1 + \Omega)\hat{a}_x^{(0)} = [b_1, (1 - \Omega)(\mathcal{F}_{-1})]. \tag{3.20}$$

Since $\hat{a}_x^{(0)}$ lies in the kernel (of the adjoint action of b_1) and Ω maps the kernel into the kernel, it follows that the l.h.s. of (3.20) also lies in the kernel. But the r.h.s. of the same equation clearly lies in the image. Therefore, we conclude that both sides vanish. Since $(1 - \Omega)(\mathcal{F}_{-1})$ cannot be a non zero element of the kernel we conclude that

$$(1 - \Omega)(\mathcal{F}_{-1}) = 0; \quad \text{or} \quad \Omega(\mathcal{F}_{-1}) = \mathcal{F}_{-1} \tag{3.21}$$

Using (3.21), we get from the third line of (2.11) that

$$(1 + \Omega)\hat{a}_x^{(-1)} = [b_1, (1 - \Omega)(\mathcal{F}_{-2})]. \tag{3.22}$$

Since $\hat{a}_x^{(-1)}$ lies in the kernel, we get that the l.h.s. of (3.22) lies in the kernel, and its r.h.s. lies in the image. Therefore both side have to vanish and so similarly to the previous case we conclude that

$$\Omega(\mathcal{F}_{-2}) = \mathcal{F}_{-2}. \tag{3.23}$$

Continuing this process we conclude that $\Omega(\mathcal{F}_{-n}) = \mathcal{F}_{-n}$, for any positive integer n , and so the group element g defined in (2.9) satisfies

$$\Omega(g) = g. \tag{3.24}$$

Indeed from (D.1) we observed that

$$P(\zeta_n) = (-1)^n \zeta_n \quad \text{and so} \quad \Omega(\zeta_n F_{-n}) = \zeta_n F_{-n}. \tag{3.25}$$

Thus we see that

$$\Omega(g F_0 g^{-1}) = g F_0 g^{-1} \tag{3.26}$$

and so from (2.18) and (2.19) we get that

$$P(\gamma^{(-2n-1)}) = -\gamma^{(-2n-1)}. \tag{3.27}$$

Thus we see that all the anomalies appearing in (2.27) satisfy (see (3.11))

$$P(X \gamma^{(-2n-1)}) = -X \gamma^{(-2n-1)}. \tag{3.28}$$

In consequence, integrating the r.h.s. of this expression over a rectangle with centre at the origin of the (\tilde{x}, \tilde{t}) coordinates (see (3.8)) we get

$$\int_{-\tilde{t}}^{\tilde{t}} dt \int_{-}^{\tilde{x}} dx X \gamma^{(-2n-1)} = 0 \tag{3.29}$$

Sending \tilde{x} to infinity we find that the charges (2.24) satisfy the mirror type symmetry (see (2.27))

$$Q^{(-2n-1)}(\tilde{t}) = Q^{(-2n-1)}(-\tilde{t}). \tag{3.30}$$

Thus, for any soliton solution satisfying the property (3.9), all the charges $Q^{(-2n-1)}$ are quasi-conserved, and the sector defined by such soliton solutions constitutes a quasi-integrable sector of the theory.

4. The analytical Hirota soliton solutions

Next we investigate the existence of analytical soliton solutions for the deformed KdV equation (1.7), or equivalently (1.1). To do this we introduce the Hirota τ -function as

$$q = \beta \ln \tau \tag{4.1}$$

for some parameter β to be appropriately chosen later. Hirota’s bilinear equations for KdV-type equations as well for the RLW and mRLW equations have been discussed in the literature [20, 31,33]. Here we follow an approach that does not make use of the Hirota’s operators D_x and D_t . Putting (4.1) into (1.7) one gets the following Hirota’s equation

$$\begin{aligned} & -\tau^2 \left[2\beta\varepsilon_2\tau_{tt}\tau_{xx} + (4\beta - 2\varepsilon_1)\tau_{xt}^2 + \tau_t(-2\varepsilon_1\tau_{xxt} - (\varepsilon_1 - 1)\tau_{xxx} + \tau_x) \right. \\ & \left. - 2\varepsilon_1\tau_x\tau_{xtt} - \varepsilon_1\tau_{tt}\tau_{xx} - 3(\varepsilon_1 - 1)\tau_{xt}\tau_{xx} - 3\varepsilon_1\tau_x\tau_{xxt} + \tau_t^2 + 3\tau_x\tau_{xxt} \right] \\ & + 2\tau \left[(\beta\varepsilon_2 - \varepsilon_1)\tau_{xx}\tau_t^2 + (\beta\varepsilon_2 - \varepsilon_1)\tau_{tt}\tau_x^2 \right. \\ & \left. + \tau_x\tau_t(4(\beta - \varepsilon_1)\tau_{xt} - 3(\varepsilon_1 - 1)\tau_{xx}) - 3(\varepsilon_1 - 1)\tau_x^2\tau_{xt} \right] \\ & - 2\tau_t\tau_x^2 [(\beta(\varepsilon_2 + 2) - 3\varepsilon_1)\tau_t - 3(\varepsilon_1 - 1)\tau_x] \\ & + \tau^3(-\varepsilon_1\tau_{xxtt} - \varepsilon_1\tau_{xxxt} + \tau_{tt} + \tau_{xt} + \tau_{xxt}) = 0. \end{aligned} \tag{4.2}$$

Next we take the one-soliton ansatz

$$\tau = 1 + \eta e^{kx - \omega t + \delta} \tag{4.3}$$

and insert it into (4.2). We consider the expansion of the equation in powers of η . In the lowest order (η^0) the equation is automatically satisfied. In the next order (η^1) we find that the equation is satisfied if either $\omega = 0$ or

$$\omega = \frac{k + (1 - \varepsilon_1)k^3}{1 - \varepsilon_1k^2}. \tag{4.4}$$

The order η^2 terms show that the equation is satisfied if

$$\frac{2k^4 [(\varepsilon_1 - 1)k^2 - 1] [\beta(\varepsilon_2 + 2)((\varepsilon_1 - 1)k^2 - 1) + 3]}{(\varepsilon_1k^2 - 1)^2} = 0. \tag{4.5}$$

If we do not want k to depend on ε 's, but do allow β to depend upon k , then we can take

$$\beta = \frac{3}{(2 + \varepsilon_2) (1 + (1 - \varepsilon_1) k^2)}. \tag{4.6}$$

With this choice one can check that all the higher order terms, in powers of η , vanish automatically, showing that the truncation of the series leads to an exact solution. Thus, we see that we have found a family of one-soliton solutions, parameterized by ε_1 and ε_2 , and given by

$$u = \frac{6}{\alpha} \frac{k^2}{(2 + \varepsilon_2) (1 - \varepsilon_1 k^2)} \frac{1}{\left[\cosh\left(\frac{\Gamma}{2}\right)\right]^2} \tag{4.7}$$

with

$$\Gamma = kx - \omega t + \delta; \quad \omega = \frac{k + (1 - \varepsilon_1) k^3}{1 - \varepsilon_1 k^2}, \tag{4.8}$$

where we have absorbed the η parameter into δ , by writing $\eta = e^{\tilde{\delta}}$ and shifting $\delta + \tilde{\delta} \rightarrow \delta$. This is the solution given in (1.12). Note that this solution, written in terms of the q field, takes the form:

$$\begin{aligned} q_{\text{type 1}} &= \frac{3}{(2 + \varepsilon_2) (1 + (1 - \varepsilon_1) k^2)} \ln(1 + e^\Gamma) \\ &= \frac{3}{(2 + \varepsilon_2) (1 + (1 - \varepsilon_1) k^2)} \left[\ln 2 + \frac{\Gamma}{2} + \ln \cosh\left(\frac{\Gamma}{2}\right) \right]. \end{aligned} \tag{4.9}$$

Note also that, as we have stated in the introduction, for $\varepsilon_1 = 1$ and $\varepsilon_2 = 0$ we get the RLW one-soliton solution, and for $\varepsilon_1 = \varepsilon_2 = 1$, we get the mRLW one-soliton. In between we get a whole new family of one-soliton solutions. Note also that our dispersion relation (the relation between ω and k) is somewhat unusual, as it depends on ε_1 . However, if we insist on keeping the usual relation, *i.e.* $\omega = \frac{k}{1-k^2}$ we can find a one-soliton solution, even for $\varepsilon_1 \neq 1$, by solving the equation (1.7) directly. This calculation is presented in the Appendix A and the obtained solution then takes the form

$$q_{\text{type 2}} = \frac{3 [1 + (\varepsilon_1 - 1) k^2]}{(2 + \varepsilon_2)} \left[\ln \left(\cosh \left(\frac{\xi}{2\sqrt{1 + (\varepsilon_1 - 1) k^2}} \right) \right) + b\xi + c \right], \tag{4.10}$$

where $\xi = kx - \omega t + \delta$, b and c are arbitrary parameters, and $\omega = \frac{k}{1-k^2}$. We have to assume $1 + (\varepsilon_1 - 1) k^2 > 0$, in order for the solution to be real. Note that in the limit $\varepsilon_1 \rightarrow 1$, both solutions, (4.9) and (4.10), become the same and are of the form

$$q = \frac{3}{(2 + \varepsilon_2)} \left[\ln \left(\cosh \left(\frac{\xi}{2} \right) \right) + b\xi + c \right]. \tag{4.11}$$

We have also checked that applying the above Hirota-type procedure for a two-soliton type ansatz, *i.e.* by expanding τ up to a second order in η , leads to a solution only for the case $\varepsilon_2 = \varepsilon_1 = 1$ and $\beta = 1$. In such a case, the Hirota's equation (4.2) becomes

$$\tau_t^2 + \tau_x \tau_t - 2\tau_{xt} \tau_t + 2\tau_{xt}^2 - 2\tau_x \tau_{xtt} + \tau_{tt} \tau_{xx} - \tau (\tau_{tt} + \tau_{xt} - \tau_{xxt}) = 0. \tag{4.12}$$

Its two-soliton solution is given by

$$\tau = 1 + e^{\Gamma_1} + e^{\Gamma_2} + A_{12} e^{\Gamma_1 + \Gamma_2}; \quad \text{for} \quad \varepsilon_2 = \varepsilon_1 = \beta = 1 \tag{4.13}$$

with

$$\Gamma_i = k_i x - \omega_i t + \delta_i \quad \omega_i = \frac{k_i}{1 - k_i^2} \quad i = 1, 2 \quad (4.14)$$

and

$$A_{12} = -\frac{(k_1 - k_2)^2 [k_1^3 k_2 - k_1^2 (k_2^2 - 1) + k_1 k_2 (k_2^2 - 4) + k_2^2 - 3]}{(k_1 + k_2)^2 [k_1^3 k_2 + k_1^2 (k_2^2 - 1) + k_1 k_2 (k_2^2 - 4) - k_2^2 + 3]}. \quad (4.15)$$

This is precisely the two-soliton solution given in (1.14)–(1.16), and first found in [19]. One can check that by replacing ω_i given in (1.15) and inserting them into (1.16), one finds that A_{12} given in (4.15) is the same as the one in (1.16).

5. The analytical quasi-integrability of the mRLW theory

In subsection 3.1 we have shown that if the field u evaluated on a one-soliton is even under the space parity (3.2) (see (3.3)), then the anomalies vanish (see (3.7)), and so all the charges $Q^{(-2n-1)}$, introduced in (2.24), are exactly conserved. Note that the family of one-soliton solutions (4.7) are even under the parity (3.2), and so this infinity of charges are exactly conserved for such one-soliton solutions.

Let us now analyse the two-soliton solution (4.13)–(4.15). Denoting

$$A_{12} = e^\Delta \quad (5.1)$$

we can write the two-soliton tau-function (4.13) as

$$\tau = 1 + e^{\Gamma_1} + e^{\Gamma_2} + e^{\Gamma_1 + \Gamma_2 + \Delta} = 2e^{z_+} \left[\cosh z_+ + e^{-\Delta/2} \cosh z_- \right], \quad (5.2)$$

where we have defined

$$z_+ = \frac{1}{2} (\Gamma_1 + \Gamma_2 + \Delta) \quad \text{and} \quad z_- = \frac{1}{2} (\Gamma_1 - \Gamma_2). \quad (5.3)$$

Note that if $k_1 = k_2$ we see that $A_{12} = 0$ and so the two-soliton solution reduces to a one-soliton solution. Therefore, for truly two-soliton solutions, *i.e.* when $k_1 \neq k_2$, z_+ and z_- can be considered independent space–time variables, *i.e.* they are linearly independent combinations of x and t . Then we have that

$$\partial_x z_\pm = \frac{1}{2} (k_1 \pm k_2) \equiv k_\pm \quad \text{and} \quad \partial_t z_\pm = -\frac{1}{2} (\omega_1 \pm \omega_2) \equiv -\omega_\pm. \quad (5.4)$$

From (1.6) and (4.1), for $\beta = 1$, we find that the two-soliton solution (4.13)–(4.15) for the u -field can be rewritten as (see (5.2))

$$\begin{aligned} u &= -\frac{8}{\alpha} \frac{(\tau \tau_{xt} - \tau_x \tau_t)}{\tau^2} \\ &= \frac{8}{\alpha} \frac{e^{-\Delta/2}}{[\cosh z_+ + e^{-\Delta/2} \cosh z_-]^2} \left[e^{\Delta/2} k_+ \omega_+ + e^{-\Delta/2} k_- \omega_- \right. \\ &\quad \left. + (k_+ \omega_+ + k_- \omega_-) \cosh z_+ \cosh z_- - (k_- \omega_+ + k_+ \omega_-) \sinh z_+ \sinh z_- \right]. \end{aligned} \quad (5.5)$$

To find the point of space–time around which we perform our parity transformation, we consider the linear relation among z_\pm and x and t , *i.e.*

$$\begin{pmatrix} z_+ \\ z_- \end{pmatrix} = \begin{pmatrix} k_+ & -\omega_+ \\ k_- & -\omega_- \end{pmatrix} \begin{pmatrix} x \\ t \end{pmatrix} + \begin{pmatrix} \frac{\Delta}{2} + \delta_+ \\ \delta_- \end{pmatrix}, \tag{5.6}$$

where we have denoted $\delta_{\pm} = (\delta_1 \pm \delta_2) / 2$. Then

$$\begin{pmatrix} \tilde{x} \\ \tilde{t} \end{pmatrix} \equiv \begin{pmatrix} x - x_{\Delta} \\ t - t_{\Delta} \end{pmatrix} = \begin{pmatrix} k_+ & -\omega_+ \\ k_- & -\omega_- \end{pmatrix}^{-1} \begin{pmatrix} z_+ \\ z_- \end{pmatrix} \tag{5.7}$$

with

$$x_{\Delta} = \frac{\omega_+ \delta_- - \omega_- (\frac{\Delta}{2} + \delta_+)}{k_+ \omega_- - k_- \omega_+}; \quad t_{\Delta} = \frac{k_+ \delta_- - k_- (\frac{\Delta}{2} + \delta_+)}{k_+ \omega_- - k_- \omega_+}. \tag{5.8}$$

Thus, the space–time parity transformation

$$P : \quad (z_+, z_-) \rightarrow (-z_+, -z_-) \tag{5.9}$$

is of the same form as (3.8), and from (5.5) we see that the field u evaluated on the two-soliton solution of the mRLW model is invariant under this parity, and so it satisfies the hypothesis made in (3.9), *i.e.* that $P(u) = u$. So, as shown in subsection 3.2, all the charges $Q^{(-2n-1)}$, defined in (2.24), satisfy the mirror symmetry described in (3.30), and they are asymptotically conserved in the scattering of two solitons.

We have thus presented an analytical proof of the quasi-integrability of the mRLW theory. It is worth adding that this is the first analytical proof of the quasi-integrability of a (non-integrable) field theory in 1 + 1 dimensions (as in this case we have an analytical form of a two-soliton solution).

6. The parity versus dynamics argument

We shall now check whether the dynamics of the deformed model (1.1) favours or not the correct parity property of the field u (3.9), so as to make the infinite set of charges $Q^{(-2n-1)}$, defined in (2.24), quasi-conserved. The plan of our approach is to write a given solution of (1.1) as a perturbative expansion around an exact solution of the integrable KdV equation. In addition, if that exact solution does satisfy the desired property (3.9) under the parity, we want to understand how the higher terms in the expansion behave under this parity. Our studies will show that the dynamics of the deformed model (1.1) does indeed favour the parity property (3.9) in a quite non-trivial and interesting way.

To make the perturbative expansion simpler we parameterise our two deformation parameters as

$$\varepsilon_1 = \varepsilon \sin \xi; \quad \varepsilon_2 = \varepsilon \cos \xi. \tag{6.1}$$

Then (1.1) becomes

$$u_t + u_x + \left[\frac{\alpha}{2} u^2 + u_{xx} + \varepsilon \left(\cos \xi \frac{\alpha}{4} w_x v_t - \sin \xi (u_{xt} + u_{xx}) \right) \right]_x = 0. \tag{6.2}$$

Next we expand the field u as

$$u = u^{(0)} + \varepsilon u^{(1)} + \varepsilon^2 u^{(2)} + \dots, \tag{6.3}$$

where each $u^{(i)}$, in general, depends upon ξ . Thus, we find that

$$\begin{aligned}
 w &= \int dt u = \int dt u^{(0)} + \varepsilon \int dt u^{(1)} + \varepsilon^2 \int dt u^{(2)} + \dots \\
 &\equiv w^{(0)} + \varepsilon w^{(1)} + \varepsilon^2 w^{(2)} + \dots
 \end{aligned} \tag{6.4}$$

and

$$\begin{aligned}
 v &= \int dx u = \int dx u^{(0)} + \varepsilon \int dx u^{(1)} + \varepsilon^2 \int dx u^{(2)} + \dots \\
 &\equiv v^{(0)} + \varepsilon v^{(1)} + \varepsilon^2 v^{(2)} + \dots
 \end{aligned} \tag{6.5}$$

Next we split all fields into their eigen-function parts under the parity (3.8). Thus

$$u^{(i,\pm)} = \frac{1}{2} (1 \pm P) u^{(i)} \tag{6.6}$$

and the same for w and v . Of course, the ε^0 part of (6.2) is the KdV equation, *i.e.*

$$u_t^{(0)} + u_x^{(0)} + \left[\frac{\alpha}{2} \left(u^{(0)} \right)^2 + u_{xx}^{(0)} \right]_x = 0 \tag{6.7}$$

The next order – ε^1 equation is then given by

$$u_t^{(1)} + u_x^{(1)} + \left[\alpha u^{(0)} u^{(1)} + u_{xx}^{(1)} + \cos \xi \frac{\alpha}{4} w_x^{(0)} v_t^{(0)} - \sin \xi \left(u_{xt}^{(0)} + u_{xx}^{(0)} \right) \right]_x = 0. \tag{6.8}$$

Next we split this equation into its even and odd parts under the parity. The odd part is given by

$$\begin{aligned}
 u_t^{(1,+)} + u_x^{(1,+)} + \left[\alpha \left(u^{(0,+)} u^{(1,+)} + u^{(0,-)} u^{(1,-)} \right) + u_{xx}^{(1,+)} \right. \\
 \left. + \cos \xi \frac{\alpha}{4} \left(w_x^{(0,+)} v_t^{(0,+)} + w_x^{(0,-)} v_t^{(0,-)} \right) - \sin \xi \left(u_{xt}^{(0,+)} + u_{xx}^{(0,+)} \right) \right]_x = 0
 \end{aligned} \tag{6.9}$$

and the even part takes the form

$$\begin{aligned}
 u_t^{(1,-)} + u_x^{(1,-)} + \left[\alpha \left(u^{(0,+)} u^{(1,-)} + u^{(0,-)} u^{(1,+)} \right) + u_{xx}^{(1,-)} \right. \\
 \left. + \cos \xi \frac{\alpha}{4} \left(w_x^{(0,+)} v_t^{(0,-)} + w_x^{(0,-)} v_t^{(0,+)} \right) - \sin \xi \left(u_{xt}^{(0,-)} + u_{xx}^{(0,-)} \right) \right]_x = 0.
 \end{aligned} \tag{6.10}$$

Let us now suppose that the zeroth order solution is even under the parity, *i.e.* that

$$P \left(u^{(0)} \right) = u^{(0)}; \quad \text{and so} \quad u^{(0,-)} = 0. \tag{6.11}$$

In this case we find that

$$w^{(0,+)} = v^{(0,+)} = 0 \tag{6.12}$$

In consequence, we see that (6.9) becomes

$$\begin{aligned}
 u_t^{(1,+)} + u_x^{(1,+)} + \left[\alpha u^{(0,+)} u^{(1,+)} + u_{xx}^{(1,+)} \right]_x \\
 = - \left[\cos \xi \frac{\alpha}{4} w_x^{(0,-)} v_t^{(0,-)} - \sin \xi \left(u_{xt}^{(0,+)} + u_{xx}^{(0,+)} \right) \right]_x \\
 u_t^{(1,-)} + u_x^{(1,-)} + \left[\alpha u^{(0,+)} u^{(1,-)} + u_{xx}^{(1,-)} \right]_x = 0.
 \end{aligned} \tag{6.13}$$

Note that $u^{(1,+)}$ satisfies an inhomogeneous equation and $u^{(1,-)}$ a homogeneous one. Thus, we can have a solution for which

$$u^{(1,-)} = 0 \quad \text{and so} \quad w^{(1,+)} = v^{(1,+)} = 0. \tag{6.14}$$

The order ε^2 part of (6.2) takes the form

$$u_t^{(2)} + u_x^{(2)} + \left[\frac{\alpha}{2} \left((u^{(1)})^2 + 2u^{(0)} u^{(2)} \right) + u_{xx}^{(2)} + \cos \xi \frac{\alpha}{4} \left(w_x^{(0)} v_t^{(1)} + w_x^{(1)} v_t^{(0)} \right) - \sin \xi \left(u_{xt}^{(1)} + u_{xx}^{(1)} \right) \right]_x = 0. \tag{6.15}$$

Splitting (6.15) into its even and odd parts one gets that the odd part becomes

$$u_t^{(2,+)} + u_x^{(2,+)} + \left[\frac{\alpha}{2} \left((u^{(1,+)})^2 + (u^{(1,-)})^2 + 2 \left(u^{(0,+)} u^{(2,+)} + u^{(0,-)} u^{(2,-)} \right) \right) + u_{xx}^{(2,+)} + \cos \xi \frac{\alpha}{4} \left(w_x^{(0,+)} v_t^{(1,+)} + w_x^{(1,+)} v_t^{(0,+)} + w_x^{(0,-)} v_t^{(1,-)} + w_x^{(1,-)} v_t^{(0,-)} \right) - \sin \xi \left(u_{xt}^{(1,+)} + u_{xx}^{(1,+)} \right) \right]_x = 0 \tag{6.16}$$

while the even part takes the form

$$u_t^{(2,-)} + u_x^{(2,-)} + \left[\frac{\alpha}{2} \left(2u^{(1,+)} u^{(1,-)} + 2 \left(u^{(0,+)} u^{(2,-)} + u^{(0,-)} u^{(2,+)} \right) \right) + u_{xx}^{(2,-)} + \cos \xi \frac{\alpha}{4} \left(w_x^{(0,+)} v_t^{(1,-)} + w_x^{(1,+)} v_t^{(0,-)} + w_x^{(0,-)} v_t^{(1,+)} + w_x^{(1,-)} v_t^{(0,+)} \right) - \sin \xi \left(u_{xt}^{(1,-)} + u_{xx}^{(1,-)} \right) \right]_x = 0. \tag{6.17}$$

Note that if (6.11), (6.12) and (6.14) are all satisfied, then these equations become

$$u_t^{(2,+)} + u_x^{(2,+)} + \left[\alpha u^{(0,+)} u^{(2,+)} + u_{xx}^{(2,+)} \right]_x = - \left[\frac{\alpha}{2} \left(u^{(1,+)} \right)^2 + \cos \xi \frac{\alpha}{4} \left(w_x^{(0,-)} v_t^{(1,-)} + w_x^{(1,-)} v_t^{(0,-)} \right) - \sin \xi \left(u_{xt}^{(1,+)} + u_{xx}^{(1,+)} \right) \right]_x, \\ u_t^{(2,-)} + u_x^{(2,-)} + \left[\alpha u^{(0,+)} u^{(2,-)} + u_{xx}^{(2,-)} \right]_x = 0. \tag{6.18}$$

So, again $u^{(2,+)}$ satisfies an inhomogeneous equation and $u^{(2,-)}$ a homogeneous one. So again, we can have a solution where

$$u^{(2,-)} = 0 \quad \text{and so} \quad w^{(2,+)} = v^{(2,+)} = 0. \tag{6.19}$$

Continuing that process it seems that we can always have a solution in which the u field is even under our parity. Such a solution of the perturbed model fits into the scheme presented in subsection 3.2 and so all the charges $Q^{(-2n-1)}$ (from their infinite set) are quasi-conserved, *i.e.* they satisfy the property (3.30). In this sense, the dynamics of the perturbed model favours the even u field, since as we saw there cannot exist pure odd u field solution. Of course, there can exist mixed solutions, *i.e.* us with an even plus a (perhaps small) odd parts.

We do not understand yet how one could fine tune the perturbed solution to be purely even under the parity. Note that the condition of making the odd part of the solution vanish at each order of perturbation, does not work like an initial condition. So, we cannot prepare the solution at a given initial time t_0 and guarantee that it will evolve in time keeping its evenness. The conditions discussed above are imposed at all times. However, as our analysis has shown that a purely odd solution cannot exist, perhaps there might be a mechanism making the odd part of the solution small (maybe its appearance is not energetically favorable). We have not found such a mechanism yet. However, our numerical simulations, which we describe below, show that if we start with a seed configuration which is even under the parity, the numerical evolution of the configuration, under the perturbed dynamics, essentially keeps the configuration even. This is a very intriguing property of our quasi-integrable field theories, and we shall describe it, in more detail, in the next sections.

7. The exact Hirota’s soliton solutions for the KdV equation

In section 6 we have shown that the dynamics of the perturbed equation (1.1) favours the even property of the u field under the parity. Since our discussion involved a perturbative expansion around an exact solution of the KdV equation (1.3), in this section we discuss the properties of the exact multi-soliton solutions under the space–time parity transformation (3.8). We show that the 1-soliton, 2-soliton and some special 3-soliton solutions of the KdV theory are even under a parity transformation and so can be used as a seed in the expansion discussed in section 6. In order to do this, we present these solutions, using the Hirota’s method, in a notation that is useful for our purposes.

We introduce the Hirota’s τ -function for the KdV equation (1.3) as

$$u = \frac{12}{\alpha} \partial_x^2 \ln \tau. \tag{7.1}$$

Putting this expression into (1.3) we get the Hirota’s equation for KdV in the form

$$2\tau_x^3 + 8\tau_{xxx} \tau_x^2 - \left[6\tau_{xx}^2 + \tau(2\tau_{xt} + 3\tau_{xx} + 5\tau_{xxx}) \right] \tau_x + \tau_t \left(2\tau_x^2 - \tau\tau_{xx} \right) + \tau [2\tau_{xx}\tau_{xxx} + \tau(\tau_{xt} + \tau_{xx} + \tau_{xxx})] = 0. \tag{7.2}$$

As it well known, the N -soliton solutions of this equation can be obtained by the Hirota’s ansatz, which takes the form

$$\tau_{N\text{-sol}} = 1 + \sum_{i=1}^N e^{\Gamma_i} + \sum_{i>j=1}^N e^{\Gamma_i+\Gamma_j+\Delta_{ij}} + \sum_{i>j>k=1}^N e^{\Gamma_i+\Gamma_j+\Gamma_k+\Delta_{ij}+\Delta_{ik}+\Delta_{jk}} + \dots + e^{\Gamma_1+\Gamma_2+\dots+\Gamma_N+\sum_{i>j=1}^N \Delta_{ij}} \tag{7.3}$$

with

$$\Gamma_i = k_i x - \omega_i t + \delta_i ; \quad \omega_i = k_i \left(1 + k_i^2 \right) ; \quad e^{\Delta_{ij}} = \frac{(k_i - k_j)^2}{(k_i + k_j)^2} \tag{7.4}$$

Then, the 1-soliton solution reads as

$$u_{\text{one-sol}} = \frac{3}{\alpha} \frac{k^2}{\cosh^2 [(kx - \omega t + \delta) / 2]}. \tag{7.5}$$

So we see that the KdV one-soliton is invariant under the space parity $P_{\bar{x}}$, defined in (3.2).

The 2-soliton solution can be cast in the form

$$u_{\text{two-sol}} = \frac{12}{\alpha} \frac{1}{\left(e^{\Delta/2} \cosh(z_+) + \cosh(z_-) \right)^2} \left[e^{\Delta/2} \left(k_-^2 + k_+^2 \right) \cosh(z_-) \cosh(z_+) + k_-^2 \cosh^2(z_-) - \left(k_- \sinh(z_-) + k_+ e^{\Delta/2} \sinh(z_+) \right)^2 + k_+^2 e^{\Delta} \cosh^2(z_+) \right], \quad (7.6)$$

where

$$z_+ = \frac{1}{2} (\Gamma_1 + \Gamma_2 + \Delta), \quad z_- = \frac{1}{2} (\Gamma_1 - \Gamma_2), \quad k_{\pm} \equiv (k_1 \pm k_2) / 2. \quad (7.7)$$

Thus, the KdV 2-soliton solution is even under the parity transformation $P : (z_+, z_-) \rightarrow (-z_+, -z_-)$, which is of the same form as (3.8).

In order to study the parity properties of the 3-soliton solutions we introduce the quantities

$$z_+ = z_-^{(1)} + z_-^{(2)} + z_-^{(3)}, \quad z_-^{(i)} = k^{(i)} x - \omega^{(i)} t + \Delta^{(i)}, \quad i = 1, 2, 3, \quad (7.8)$$

where

$$\begin{aligned} k^{(1)} &= \frac{1}{2} (-k_1 + k_2 + k_3), & \omega^{(1)} &= \frac{1}{2} (-\omega_1 + \omega_2 + \omega_3), & \Delta^{(1)} &= \frac{1}{2} (-\delta_1 + \delta_2 + \delta_3 + \Delta_{23}), \\ k^{(2)} &= \frac{1}{2} (k_1 - k_2 + k_3), & \omega^{(2)} &= \frac{1}{2} (\omega_1 - \omega_2 + \omega_3), & \Delta^{(2)} &= \frac{1}{2} (\delta_1 - \delta_2 + \delta_3 + \Delta_{13}), \\ k^{(3)} &= \frac{1}{2} (k_1 + k_2 - k_3), & \omega^{(3)} &= \frac{1}{2} (\omega_1 + \omega_2 - \omega_3), & \Delta^{(3)} &= \frac{1}{2} (\delta_1 + \delta_2 - \delta_3 + \Delta_{12}). \end{aligned}$$

The 3-soliton τ -function can be written as $\ln \tau_{\text{three-sol}} = \ln 2 + z_+ + \ln F$, with

$$F = \cosh z_+ + e^{-(\Delta_{12} + \Delta_{13})/2} \cosh z_-^{(1)} + e^{-(\Delta_{12} + \Delta_{23})/2} \cosh z_-^{(2)} + e^{-(\Delta_{13} + \Delta_{23})/2} \cosh z_-^{(3)}. \quad (7.9)$$

Thus, for the 3-soliton solution we get:

$$u_{\text{three-sol}} = \frac{12}{\alpha} \partial_x^2 \ln F = \frac{12}{\alpha} \left[\frac{\partial_x^2 F}{F} - \left(\frac{\partial_x F}{F} \right)^2 \right] \quad (7.10)$$

with

$$\begin{aligned} \partial_x F &= \left(k^{(1)} + k^{(2)} + k^{(3)} \right) \sinh z_+ + k^{(1)} e^{-(\Delta_{12} + \Delta_{13})/2} \sinh z_-^{(1)} \\ &\quad + k^{(2)} e^{-(\Delta_{12} + \Delta_{23})/2} \sinh z_-^{(2)} + k^{(3)} e^{-(\Delta_{13} + \Delta_{23})/2} \sinh z_-^{(3)} \end{aligned} \quad (7.11)$$

and

$$\begin{aligned} \partial_x^2 F &= \left(k^{(1)} + k^{(2)} + k^{(3)} \right)^2 \cosh z_+ + \left(k^{(1)} \right)^2 e^{-(\Delta_{12} + \Delta_{13})/2} \cosh z_-^{(1)} \\ &\quad + \left(k^{(2)} \right)^2 e^{-(\Delta_{12} + \Delta_{23})/2} \cosh z_-^{(2)} + \left(k^{(3)} \right)^2 e^{-(\Delta_{13} + \Delta_{23})/2} \cosh z_-^{(3)}. \end{aligned} \quad (7.12)$$

One way of implementing the parity argument for the KdV three-soliton solution is to have the inversion (change of signs) of all three $z_-^{(i)}$'s, i.e. $z_-^{(i)} \rightarrow -z_-^{(i)}$, for $i = 1, 2, 3$, which would

also imply $z_+ \rightarrow -z_+$. However, we are performing our calculations in two dimensions and the three $z_-^{(i)}$'s cannot be linearly independent. Each $z_-^{(i)}$ defines a straight line in the (x, t) plane, and if they are to be simultaneously inverted, we need these three lines to cross at the same point, and so all three $z_-^{(i)}$'s should vanish at this point. So, there should exist a point (x_Δ, t_Δ) such that

$$\begin{pmatrix} k^{(1)} & -\omega^{(1)} & \Delta^{(1)} \\ k^{(2)} & -\omega^{(2)} & \Delta^{(2)} \\ k^{(3)} & -\omega^{(3)} & \Delta^{(3)} \end{pmatrix} \begin{pmatrix} x_\Delta \\ t_\Delta \\ 1 \end{pmatrix} = 0. \tag{7.13}$$

For this to happen, the determinant of this 3×3 matrix should vanish, and this implies

$$\begin{aligned} &(\Delta^{(1)} + \Delta^{(2)}) (k_1^3 k_2 - k_1 k_2^3) + (\Delta^{(1)} + \Delta^{(3)}) (k_3^3 k_1 - k_3 k_1^3) \\ &+ (\Delta^{(2)} + \Delta^{(3)}) (k_2^3 k_3 - k_2 k_3^3) = 0. \end{aligned}$$

Note that if we choose any pair of k_i 's to be equal, we reduce the three-soliton solution to a two-soliton solution. So, we need $k_1 \neq k_2 \neq k_3$. One way of satisfying this is to choose $\delta_i, i = 1, 2, 3$, in such a way that

$$\Delta^{(1)} = \Delta^{(2)} = \Delta^{(3)} = 0. \tag{7.14}$$

Note that in such a case the $z_-^{(i)}$'s, given in (7.8), (and so z_+) become homogeneous in x and t , i.e.

$$z_-^{(i)} = k^{(i)} x - \omega^{(i)} t \quad i = 1, 2, 3 \tag{7.15}$$

Therefore, the parity transformation

$$P : \quad (x, t) \rightarrow (-x, -t) \tag{7.16}$$

is sufficient to have what we want, i.e. $z_-^{(i)} \rightarrow -z_-^{(i)}$, and so $z_+ \rightarrow -z_+$.

The condition (7.14) leads to the linear system

$$\begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} \delta_1 \\ \delta_2 \\ \delta_3 \end{pmatrix} = \begin{pmatrix} \Delta_{23} \\ \Delta_{13} \\ \Delta_{12} \end{pmatrix} \tag{7.17}$$

and so

$$\begin{aligned} \delta_1 &= -\frac{1}{2} (\Delta_{13} + \Delta_{12}) = -\frac{1}{2} \ln \left[\frac{(k_1 - k_3)^2 (k_1 - k_2)^2}{(k_1 + k_3)^2 (k_1 + k_2)^2} \right], \\ \delta_2 &= -\frac{1}{2} (\Delta_{23} + \Delta_{12}) = -\frac{1}{2} \ln \left[\frac{(k_2 - k_3)^2 (k_1 - k_2)^2}{(k_2 + k_3)^2 (k_1 + k_2)^2} \right], \\ \delta_3 &= -\frac{1}{2} (\Delta_{23} + \Delta_{13}) = -\frac{1}{2} \ln \left[\frac{(k_2 - k_3)^2 (k_1 - k_3)^2}{(k_2 + k_3)^2 (k_1 + k_3)^2} \right]. \end{aligned} \tag{7.18}$$

Denoting

$$\Gamma_i = \tilde{\Gamma}_i + \delta_i \quad \text{with} \quad \tilde{\Gamma}_i = k_i x - \omega_i t \tag{7.19}$$

we find that the corresponding τ -function for the 3-soliton solution takes the form

$$\begin{aligned} \tau_{\text{three-sol}} = & 1 + e^{\tilde{\Gamma}_1 + \tilde{\Gamma}_2 + \tilde{\Gamma}_3} + e^{-(\Delta_{13} + \Delta_{12})/2} \left(e^{\tilde{\Gamma}_1} + e^{\tilde{\Gamma}_2 + \tilde{\Gamma}_3} \right) \\ & + e^{-(\Delta_{23} + \Delta_{12})/2} \left(e^{\tilde{\Gamma}_2} + e^{\tilde{\Gamma}_1 + \tilde{\Gamma}_3} \right) + e^{-(\Delta_{23} + \Delta_{13})/2} \left(e^{\tilde{\Gamma}_3} + e^{\tilde{\Gamma}_1 + \tilde{\Gamma}_2} \right). \end{aligned} \quad (7.20)$$

One can check that in all these cases the three soliton collide simultaneously at the origin $x = 0$ at $t = 0$. The conclusion we can draw from this is that, for the KdV three-soliton solution in which the three solitons collide at the same point in space–time, the field u is even under the parity (7.16), when evaluated on such a solution.

8. Numerical analysis of quasi-conserved charges

In this section we discuss results of some of our numerical simulations of equation (1.7) performed for various values of ε_1 and ε_2 . We particularly focus on investigating the corresponding first non-trivial quasi-conserved charge $Q^{(-3)}$, defined in equation (2.24). The numerical scheme used to approximate equation (1.7) is very similar to the algorithm used to solve the mRLW equation [6]. We have adjusted this scheme appropriately, which is discussed in appendix B, for the simulations presented in this section.

We have used several different initial conditions depending on the values of ε_1 and ε_2 used for each simulation, and we discuss these initial conditions in more detail in the subsections below. We have performed the numerical experiments for two-soliton simulations for a range of values for Γ_1 and Γ_2 but, for consistency, all the plots presented in this section regarding two-soliton simulations are obtained using the same values regardless of the initial conditions used, and similarly for all the three-soliton simulations (Γ_1 , Γ_2 and Γ_3) presented in this section. For all the simulations presented in this section we have used a grid spacing $h = 0.1$ and time level $\tau = 0.001$. More details on the variables used for the simulations discussed in this section can be found in appendix B.2.

8.1. Quasi-conserved charges of the mRLW equation

In this subsection we discuss the results of our investigations of the quasi-conserved charges of the mRLW equation. To perform the simulations of the mRLW equation, we have set $\varepsilon_1 = \varepsilon_2 = 1$ in equation (1.7).

8.1.1. Two-soliton solutions of the mRLW equation

In Fig. 1 we have plotted the scattering of two solitons described by equation (1.14) at various values of t , and in Fig. 2a we have plotted the time dependence of $\partial_t Q^{(-3)}$ for this interaction. Since the mRLW equation possesses an analytical form of the two-soliton solution, in Fig. 2 we have in fact plotted the values $\partial_t Q^{(-3)}$ for both the analytical and numerical simulation. To better illustrate how small the discrepancy is, we have added an insert of the region near the global maximum on a much smaller scale. We note that there is hardly any discrepancy between the analytical and numerical values.

Now, comparing Figs. 1 and 2a, we see that as $t \rightarrow \pm\infty$ the charge is conserved except for during the collision, where it first decreases and then increases back to the same value as before the collision. Fig. 2b presents the plots of $\int_0^t dt' \partial_{t'} Q^{(-3)}$, and we see that the two lumps exactly cancel each other out, that is,

$$\lim_{t \rightarrow -\infty} Q^{(-3)} = \lim_{t \rightarrow \infty} Q^{(-3)}. \quad (8.1)$$




Fig. 1. The spatial-dependence of the u field for two solitons described by mRLW equation, interacting with each other at different points in time, as given in equations (1.6) and (1.14).




Fig. 2. The time-dependence of the analytical and numerically (represented by the red line and the green line, respectively) obtained values of the quantity $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ for the two-soliton simulation of the mRLW equation. (For interpretation of the colors in the figure(s), the reader is referred to the web version of this article.)

Thus, we can conclude that these quantities are indeed quasi-conserved, as has been shown analytically in section 5. The agreement of the analytical and numerical results is a good test of the numerical code that we are using.

8.1.2. Three-soliton solutions of the mRLW equation

In Fig. 3 we present a selection of plots (at different values of t) of the u field obtained in a typical numerical simulation of a superposition of three one-soliton solutions. This was done in this way since we do not have analytic three-soliton solutions of the mRLW equation. However, as was discussed in [6], such a field is a very good approximation to a solution and the solitons of the numerical three-soliton configuration behave as integrable (or quasi-integrable) solitons. So here we have tested this by looking at the behaviour of corresponding charges. Fig. 4 shows the corresponding time dependence of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$. As $Q^{(-3)}$ changes only during the collision, and has the same value in the asymptotic regions (see Fig. 4b), we conclude that our charge is also quasi-conserved for the three-soliton configuration.




Fig. 3. Snapshots at different points in time of the numerical three-soliton solution of the mRLW equation (equation (1.7) for $\varepsilon_1 = \varepsilon_2 = 1$), for the case when the solitons collide all around the point $x = 0$.




Fig. 4. The time-dependence of the obtained values of the quantity $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ for the three-soliton simulation of the mRLW equation.

In section 7 we have shown that the u -field for the exact Hirota three-soliton of the KdV equation is even under the parity transformation (7.16), when the three solitons collide all at the same point. As we have discussed in section 6, one should therefore expect that the deformed three-soliton solution should keep that parity property, and so to have an infinity of quasi-conserved charges given by (2.24) (see section 3.1). The numerical results presented in Fig. 4 show that the charge $Q^{(-3)}$ is indeed quasi-conserved for the mRLW equation, and so it confirms this expectation.

8.2. Quasi-conserved charges of the RLW equation

In this subsection we discuss similar topics for the RLW equation. This model, is described by equation (1.4), in which the conventional re-scaling of the u -field used in the literature sets $\alpha = 12$. The model possesses the same exact one-soliton solution with the same dispersion relation as the mRLW equation, but it does not possess an analytical two-soliton solutions [7].




Fig. 5. The field u , at selected values of time, seen in our simulation of a two-soliton configuration of the RLW equation.




Fig. 6. The time-dependence of the obtained values of the quantity $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ for the two-soliton simulation of the RLW equation.

However, since we are simulating equation (1.7) with $\varepsilon_1 = 1$ and $\varepsilon_2 = 0$, we must perform the appropriate rescaling of q (i.e., $q \rightarrow \frac{3}{2}q$) in order to obtain the analytical one-soliton solution. Thus, to construct the two- and three numerical systems we have used, as initial conditions for the numerical simulation, a linear superposition of the following analytical single-soliton solutions

$$q = \frac{3}{2} \sum_{i=1}^n \ln(1 + e^{\Gamma_i}) \tag{8.2}$$

where $n = 2$ corresponds to the two-soliton simulation and $n = 3$ to the three-soliton simulation.

8.2.1. Two-soliton solutions of the RLW equation

The results of the two-soliton simulation for the RLW equation are shown in Fig. 5. Clearly, the figure shows some radiation (see the inserts in Figs. 5e and 5f), and this agrees with the results presented in [24]. This is expected since the initial conditions used for this simulation do not solve the RLW equation analytically. Note that the amplitudes of these solitons are bigger compared to the solitons of the mRLW equation due to the aforementioned rescaling of q . Furthermore, in




Fig. 7. The field u , for a few values of time, found in our simulation of the three-soliton system of the RLW equation.




Fig. 8. The time-dependence of the quantity $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ found in our three-soliton simulation of the RLW equation.

Fig. 6 we have presented the plots of the time dependence of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen in this simulation. The shapes of the curves in these plots are very similar to the ones found in Fig. 2. We note that the small observed emission of radiation does not seem to have a visible effect on the quasi-conservation of the charge.

8.2.2. Three-soliton solutions of the RLW equation

Next we have looked at the three solitons systems of the RLW equation. In Fig. 7 we present the plots of the fields u seen at various times in their evolution, and in Fig. 8 the plots of the corresponding $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$.

Note that the amplitudes of the solitons are again larger when compared with the solitons of the mRLW equation (see Fig. 3). This is due to the factor of $3/2$, mentioned before. The three solitons behave in a very similar way to the solitons shown in Fig. 3 except that, around the time they collide, a small bit of radiation is emitted (see the inserts in Figs. 7d, 7e and 7f). Just as for the two-soliton simulation, this emission of radiation does not seem to have a visible effect on the quasi-conservation of the charge.




Fig. 9. The time-evolution of two-soliton systems found in our simulation of equation (1.7) with $\varepsilon_1 = 1$, $\varepsilon_2 = 0.5$ (red curve) and $\varepsilon_1 = 1$, $\varepsilon_2 = 1.5$ (green curve).

8.3. Intermediate models: quasi-conserved charges for $\varepsilon_1 = 1$ and $\varepsilon_2 \neq 1$

So far we have discussed the properties of solutions of the mRLW and RLW equations. However, as we have shown in section 4, our basic equation (1.7) possesses one-soliton solutions, whose analytical form is given by equation (1.8), for any real value of ε_2 as long as $\varepsilon_1 = 1$. So in this section we investigate the question of quasi-integrability of two- and three-soliton systems by performing simulations with the initial conditions taken from linear superpositions of such analytical single-soliton solutions, that is, from

$$q = \frac{3}{(2 + \varepsilon_2)} \sum_{i=1}^n \ln(1 + e^{\Gamma_i}), \quad (8.3)$$

where $n = 2$ corresponds to the two-soliton simulation and $n = 3$ to the three-soliton simulation. We present the results of such two- and three-soliton simulations of equation (1.7), for various values of $\varepsilon_2 \neq 1$ while keeping $\varepsilon_1 = 1$ constant.

8.3.1. Two-soliton configurations

As before, first we present our results for two solitons. Fig. 9 shows the time-evolution of the u field for a simulation with $\varepsilon_2 = 0.5$ and a simulation with $\varepsilon_2 = 1.5$. Figs. 10a and 11a show the time dependence of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen these simulations. The system for $\varepsilon_2 = 0.5$ emits a small amount of (visible) radiation during the interaction of the soliton fields (similarly to the two-soliton simulation for the RLW equation (see subsection 8.2.1)). This radiation is difficult to spot on the scale shown in Fig. 9, and so we have added inserts in plots 9e and 9f to show more clearly this radiation emitted during the interaction. Looking at Fig. 10a, we see that $Q^{(-3)}$ changes significantly only during the interaction of the two soliton fields. Furthermore, Fig. 11a shows that the quantity $Q^{(-3)}$ satisfies




Fig. 10. The time-dependence of the obtained values of the quantity $\partial_t Q^{(-3)}$ of the simulation of equation (1.7), for two-soliton solutions, with different values of $\epsilon_1 = 1, \epsilon_2 \neq 1$.




Fig. 11. The time-dependence of the obtained values of $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen in our simulations of equation (1.7), for two-soliton solutions, for $\epsilon_1 = 1$, and $\epsilon_2 = 0.7$ and $\epsilon_2 = 1.3$.

$$\lim_{t \rightarrow -\infty} Q^{(-3)} = \lim_{t \rightarrow \infty} Q^{(-3)}, \tag{8.4}$$

which is again what we saw for the RLW and mRLW equation.

The other plots in Figs. 10 and 11 show the time dependence of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ for simulations with various other values of $\epsilon_2 \neq 1$. They show similar patterns to the ones described for $\epsilon_2 = 0.5$ and $\epsilon_2 = 1.5$. Note that as $|1 - \epsilon_2|$ becomes smaller, the values of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ converge closer to the values displayed in Fig. 2.

8.3.2. Three-soliton configurations

Next, we also looked at some three solitons systems which were constructed using equation (8.3), with $n = 3$, as the initial conditions. Fig. 12 shows the time evolution seen in two simulations for the values $\epsilon_1 = 1, \epsilon_2 = 0.5$ and $\epsilon_1 = 1, \epsilon_2 = 1.5$. These plots show that the three solitons interact with each other in a way very similar to the mRLW and RLW equation (see subsections 8.1.2 and 8.2.2). Unlike for the two-soliton simulation, we now see slightly more radiation emitted in the $\epsilon_2 = 1.5$ case than in the $\epsilon_2 = 0.5$ case (see the insert in Fig. 12f). Figs. 13 and 14 present the plots of time dependence of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen in our simulations. Fig. 13 clearly shows that $Q^{(-3)}$ only changes during the interaction of the soliton fields, and Fig. 14 demonstrates that this quantity is indeed truly quasi-conserved in the sense that

$$\lim_{t \rightarrow -\infty} Q^{(-3)} = \lim_{t \rightarrow \infty} Q^{(-3)}. \tag{8.5}$$




Fig. 12. The time-evolution of two three-soliton systems seen in our simulations of the equation (1.7) with $\varepsilon_1 = 1$, $\varepsilon_2 = 0.5$ (red curve) and $\varepsilon_1 = 1$, $\varepsilon_2 = 1.5$ (green curve).




Fig. 13. The time-dependence of $\partial_t Q^{(-3)}$ for a three-soliton system as determined by the simulation of equation (1.7) with $\varepsilon_1 = 1$ and different values of $\varepsilon_2 \neq 1$.




Fig. 14. The time-dependence of the quantity $\int_0^t dt' \partial_{t'} Q^{(-3)}$ as determined in the simulation of equation (1.7), for three-soliton solutions, with $\varepsilon_1 = 1$ and different values of $\varepsilon_2 \neq 1$.




Fig. 15. The u field configurations seen in our time-evolution of the numerical time-evolution of the equation (1.7) for $\varepsilon_1 = 1$ and $\varepsilon_2 = 1.5$ at several values of time t .

8.4. Intermediate models: quasi-conserved charges for various values of ε_1 and ε_2

We also would like to consider other values of ε_1 and ε_2 . However, we have no analytical expressions of one-soliton solutions when $\varepsilon_1 \neq 1$. Therefore, we have used the analytical two-soliton solutions of the mRLW equation as initial conditions for the two-soliton configurations, and the linear superposition of three single-soliton solutions of the mRLW equation for the three-soliton simulations.

In the next subsection we use the same values as those used in subsection 8.3 (for $\varepsilon_1 = 1$ and $\varepsilon_2 \neq 1$) to simulate the two-soliton configurations. We have repeated these simulations to study the effect of using the ‘wrong’ initial conditions. This was achieved by starting our simulations with the initial conditions corresponding to $\varepsilon_1 = 1$ and ε_2 also equal to 1 and then performing the simulations for different values of ε_2 . Then in the subsequent subsections we investigate the quasi-integrability for various values of $\varepsilon_1 \neq 1$ while keeping $\varepsilon_2 = 1$.

8.4.1. Two-soliton configurations for $\varepsilon_1 = 1$ and $\varepsilon_2 \neq 1$

Fig. 15 presents various plots of the simulation with $\varepsilon_1 = 1$ and $\varepsilon_2 = 1.5$, and Fig. 16 shows the plots of the time-dependence of the corresponding quantities $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$. Fig. 15a shows the initial condition at $t = 0$. Fig. 15b shows these two soliton fields after they have evolved for 5 units of time. Looking at that figure, we note that both solitons have moved to the right and have left some radiation behind immediately after the start of the simulation, while their amplitudes have increased slightly. This is because the true soliton of such a system is probably significantly different the one described by the initial conditions used in this simulation. This implies that the field u changes from the form given by the initial conditions and settles to a true solution, and in doing so the two solitons leave behind some radiation. Then Fig. 15c shows the largest soliton field after it has interacted with the radiation emitted by the smallest soliton field. Fig. 15d shows the solution at $t = 15$ when the largest soliton starts to interact with the smallest soliton. Fig. 15e shows the actual soliton fields interacting with each other.




Fig. 16. The time dependence of the quantity $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ obtained in two-soliton simulation of equation (1.7) with $\varepsilon_1 = 1$ and $\varepsilon_2 = 1.5$.




Fig. 17. The time-dependence of the obtained values of $\partial_t Q^{(-3)}$ seen in the simulation of equation (1.7) for $\varepsilon_1 = 1$ and $\varepsilon_2 \neq 1$, with the analytical two-soliton solutions of the mRLW equation used as initial conditions.

Finally, Fig. 15f shows the field when the solitons have moved away from each other again. These changes are all seen in Fig. 16a; from the beginning (at $t = 0$ to $t \simeq 2$) the initial conditions settle down to the true solution of the equation leaving behind some radiation, from about $t \simeq 6$ to $t \simeq 10$ the largest soliton interacts with the radiation left behind by the smallest soliton. Finally, from $t \simeq 16$ to $t \simeq 23$ the two soliton fields interact, after which the solitons keep moving unhindered. Notice also that Fig. 16b shows that, after the initial conditions settle down to the true solution, all the lumps cancel each other out.

For other values of $\varepsilon_1 = 1$ and $\varepsilon_2 \neq 1$ we have observed a very similar behaviour to the one described above. To illustrate this better, Fig. 17 presents three plots comparing $\partial_t Q^{(-3)}$ for $\varepsilon_2 = 0.5$ and $\varepsilon_2 = 1.5$ (see Fig. 17a), $\varepsilon_2 = 0.7$ and $\varepsilon_2 = 1.3$ (see Fig. 17b), and $\varepsilon_2 = 0.9$ and $\varepsilon_2 = 1.1$ (see Fig. 17c). We see that the larger $|1 - \varepsilon_2|$ becomes, the more intense are the changes of $Q^{(-3)}$ when the solitons readjust their form from the one described by the initial conditions and when the larger soliton interacts with the radiation left behind by the smaller soliton. This is expected since the larger the value of $|1 - \varepsilon_2|$ is, the larger is the difference between the initial conditions and the true solutions of the equation, which results in a faster rate of change of the charge $Q^{(-3)}$. Furthermore, this implies that the radiation left behind by the soliton is larger, and so the interaction between the larger soliton and the radiation due to the smaller soliton is more intense. However, when the two actual soliton fields collide, we see that as $|1 - \varepsilon_2|$ becomes smaller, the peak of $\partial_t Q^{(-3)}$ converges closer to the peak shown in Fig. 2a, both in terms of amplitude as well as in terms of its location on the horizontal axis. The horizontal movement is due to the fact that when the solitons emit radiation, their amplitudes increase/decrease (depending on whether $\varepsilon_2 < 1$ or $\varepsilon_2 > 1$) which causes their velocities to increase/decrease as well.




Fig. 18. The time-evolution of two two-soliton systems studied in the equation (1.7) with $\epsilon_1 = 1$ and $\epsilon_2 = 0.5$ (red curve), and $\epsilon_1 = 1$ and $\epsilon_2 = 1.5$ (green curve).




Fig. 19. The time-dependence of $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen in the simulation of equation (1.7) with different values of $\epsilon_1 = 1$ and $\epsilon_2 \neq 1$.

To illustrate better these observed properties, Fig. 18 presents again the various plots shown in Fig. 15 corresponding to $\epsilon_1 = 1$ and $\epsilon_2 = 1.5$ but it also shows the results of the simulation corresponding to $\epsilon_1 = 1$ and $\epsilon_2 = 0.5$, in order to compare the two simulations. This figure also illustrates the fact that if $\epsilon_2 > 1$, then the solitons’ amplitudes and velocities increase as a result of the emission of this radiation. On the other hand, if $\epsilon_2 < 1$, the solitons’ amplitudes and velocities decrease. Fig. 19 presents the corresponding values of $\int_0^t dt' \partial_{t'} Q^{(-3)}$. Overall, we see that after the fields have settled from their form given by the initial conditions to the values corresponding to a solution, all the lumps in Fig. 17 cancel each other out.

8.4.2. Two-soliton configurations for $\epsilon_1 \neq 1$ and $\epsilon_2 = 1$

Armed with the observation of the last subsection we present here results of some simulations for $\epsilon_1 \neq 1$ and $\epsilon_2 = 1$ where, just as in the previous subsection, the initial conditions were constructed from the analytical two-soliton solution of the mRLW equation.

Fig. 20 shows the u fields seen in the two-soliton simulation corresponding to $\epsilon_1 = 0.5$ (represented by the red curve) and $\epsilon_1 = 1.5$ (represented by the green curve). We observe that when




Fig. 20. The time-evolution of two two-soliton systems of equation (1.7) for $\epsilon_1 = 0.5, \epsilon_2 = 1$ (red curve) and $\epsilon_1 = 1.5, \epsilon_2 = 1$, (green curve) in which the initial conditions were taken from solitons of the mRLW equation.




Fig. 21. The time-dependence of the determined values of the quantity $\partial_t Q^{(-3)}$ see in the simulations of equation (1.7) for different values of $\epsilon_1 \neq 1, \epsilon_2 = 1$.




Fig. 22. The time-dependence of $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen in the simulations of equation (1.7) for different values of $\epsilon_1 \neq 1, \epsilon_2 = 1$.




Fig. 23. The time-evolution of two three-soliton systems seen in our simulations of equation (1.7) for $\epsilon_1 = 0.5$ and $\epsilon_2 = 1$ (red curve), and $\epsilon_1 = 1.5$ and $\epsilon_2 = 1$ (green curve).




Fig. 24. The time-dependence of $\partial_t Q^{(-3)}$ seen in the simulations of equation (1.7) for two values of $\epsilon_1 \neq 1$ and $\epsilon_2 = 1$.

$\epsilon_1 < 1$ the radiation emitted by the solitons has initially a positive amplitude, and the solitons' amplitudes and velocities increases as a result of emission of this radiation. On the other hand, if $\epsilon_1 > 1$, the radiation emitted by the solitons has initially a negative amplitude, and the solitons' amplitudes and velocities decrease. Comparing Fig. 20 with Fig. 18, we see that changing ϵ_1 (while keeping $\epsilon_2 = 1$) results in more radiation emitted than when changing ϵ_2 (while keeping $\epsilon_1 = 1$) by the same amount. Figs. 21 and 22 present the time dependence of $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$ observed in these simulations, and for various other simulations with other values of $\epsilon_1 \neq 1$.

8.4.3. Three-soliton configurations for $\epsilon_1 \neq 1$ and $\epsilon_2 = 1$

The simulations of three soliton systems corresponding to $\epsilon_1 = 0.5$, $\epsilon_2 = 1$ and $\epsilon_1 = 1.5$, $\epsilon_2 = 1$ are shown in Fig. 23, and Figs. 24 and 25 present the time dependence of the corresponding $\partial_t Q^{(-3)}$ and $\int_0^t dt' \partial_{t'} Q^{(-3)}$. These simulations are run with the linear superposition of three analytical one-soliton solutions of the mRLW equation as initial conditions, just as in




Fig. 25. The time-dependence of the determined values of $\int_0^t dt' \partial_{t'} Q^{(-3)}$ seen in the simulations of equation (1.7) for 3 sets of values of $\varepsilon_1 \neq 1$ and $\varepsilon_2 = 1$.

subsubsection 8.1.2. These simulations show similar patterns to those seen for the two-soliton simulations.

9. Conclusions

We have performed a detailed study of the quasi-integrability properties of various deformations of the KdV equation which include among them the RLW and mRLW equations. The charges were constructed by introducing an anomalous zero-curvature condition as explained in section 2. Note that only the time component A_t of the Lax pair was deformed away from the KdV potentials. The space component A_x was not deformed and so the charges $Q^{(-2n-1)}$, that we presented in (2.24), are the same as the exact conserved charges of the KdV equation. The difference lies in the anomalies $\alpha^{(-2n-1)}$, also introduced in (2.24), which vanish for KdV but not for its deformations. However, we have also shown that the deformations of the KdV theory considered in this paper possess a very interesting property. Even though the charges are not conserved and do vary during the scatterings of solitons, they all return, after the scattering, to the values they had before the scattering. It is this property, the asymptotic conservation of the charges, that defines what we call a quasi-integrability of the theory. The mechanisms underlying such an asymptotic conservation of the charges are still not fully understood. Our results are clearly consistent with the property of asymptotic stability of these models but we have tried to relate them to some specific properties of the soliton fields. From these studies we have learned that the quasi-integrability goes hand in hand with some special properties of the soliton solutions under a specific space–time parity transformation, as explained in section 3, and that these properties lead to the asymptotic vanishing of the anomalies. In addition, another intriguing property that correlates with quasi-integrability is that if a soliton solution of the exactly integrable theory possesses the correct behaviour under this parity, then the dynamics of the deformed theories seem to preserve this behaviour under a perturbative expansion around the integrable theory. In section 6 we have presented the details of such an expansion and showed its compatibility with the expected parity behaviour of the solutions.

Our analysis of the deformations of the KdV theory present two novel ideas in the study of the quasi-integrability. First, we believe that it is the first time that the quasi-integrability ideas were tested for the scattering of three solitons, where the implementation of the parity transformation is much more involved. When the three solitons scatter in separated pairs, it is expected that the quasi-integrability of two-soliton scattering would be preserved, since the third soliton is away and not interacting with the other two. However, when all three solitons collide together at the same time and in a given position in space, the three body interactions come into play, and one cannot rely on the two body interactions to analyze the quasi-integrability. We have

shown in section 7 that the exact Hirota three-soliton solutions of the KdV equation do have the correct parity properties when their solitons all collide together at the same point in space. Using the results of section 6 one could therefore expect that deformed three-soliton solutions would possess the same property. That is exactly what our numerical simulations have shown: the charges are indeed asymptotically conserved when the three solitons of various deformations of the KdV theory collide together. This provides a strong support for our working definition of quasi-integrability.

The other key result of the present paper is that it presents the first example of an analytical and not only numerical, demonstration, of the quasi-integrability of a truly non-integrable theory. This was discussed in section 5 where we have showed that the two-soliton sector of the mRLW theory was analytically quasi-integrable. The proof of this result is valid for all charges from their infinite set. Our numerical simulations have confirmed it for the case of the first non-trivial charge. For next charges the results were also supportive; however, as the next charges involved expressions which contained terms with more derivatives, our numerical results had more numerical errors and so were less reliable. Clearly, better numerical techniques have to be developed to prove our claims more decisively (*i.e.* to get the same accuracy as we had for $Q^{(-3)}$). This work lies beyond the present paper.

We must emphasise, however, that the strongest support for our working definition of quasi-integrability comes from the results of our numerical simulations. In all simulations reported in section 8 it has been shown that the charge $Q^{(-3)}$ is quasi-conserved, *i.e.* that it returns, after the scattering of two or three solitons, to the value it had before the scattering. This quasi-conservation has been tested to a very good precision. In addition, our simulations have shown what had been observed before in other theories; namely, that the charge only varies in time when the solitons come together and interact with each other. The charge remains constant when the solitons are far apart. Another important and intriguing result of our simulations is related to the observation that our simulations have shown that the interaction of radiation with the solitons also seems to respect quasi-integrability.

For our simulations we have used the initial fields constructed from various combinations of one soliton solutions of our equations. At the beginning of our study we did not have an expression for one soliton field when ε_1 was different from one. So we used slightly incorrect fields as seeds for these simulations. In these cases we observed a small amount of radiation sent out while the initial configuration gradually settled to a proper solution of the deformed theory. The radiation emitted by each soliton interacted nonlinearly with the other solitons, and we observed that such a radiation-soliton interaction made the charge $Q^{(-3)}$ to vary quite reasonably in time. However, when the radiation and the soliton have got separated, the charge returned to its original value. So, we have observed that the radiation-soliton interaction also seems to respect quasi-integrability, but at this stage we do not have any clear idea why this property holds too. Our analytical parity arguments do not apply to such an interaction. All our results have confirmed our expectations as to the behaviour of the fields – supporting our claims of quasi-integrability of our models.

We have recently found analytical expressions for 2 different one soliton solutions of the models when $\varepsilon_1 \neq 1$, which become the same, and agree with the known expression, in the limit of $\varepsilon_1 \rightarrow 1$. We expect that simulations with such fields would give even stronger results but they are clearly outside the scope of this paper. As the two solutions have different dispersion relations it would be interesting to study the effects of their difference on the properties of various superpositions of the fields and their scattering properties. As our results, started with slightly incorrect initial fields, were in good agreement with our quasi-integrability expectations (as the radiation ef-

fects were small and restricted themselves to the initial re-adjustments) we expect the effects also to be very small, and though very important, they are clearly not likely to be of great help in what we have tried to demonstrate in this paper. This is clearly one of the topics to study in future work.

The numerical methods used in this paper involved the specially constructed numerical programs which were based on the original program of [12] and have been appropriately adapted to the numerical study of the equation (1.7). More details are given in appendix B. The results of our numerical simulations supported very well our expectations. This was particularly true for the lowest charge but it was also true to higher charges. However, expressions for higher charges involved more derivatives of the fields and so our expressions were more prone to numerical errors. Thus, although their behaviour supported our claims, the reliability of these results was not as sound as for the ones we have included in this paper.

Acknowledgements

LAF and WJZ want to thank the Royal Society for its grant which supported their research. All three authors want to thank FAPESP and Durham University for their joint ‘Collaboration grant’ (SPRINT) that supported their reciprocal visits to São Carlos and Durham which have made the research described in this paper possible. LAF is partially supported by CNPq-Brazil. The authors are very grateful to the Referee for the very careful reading of the manuscript and for very valuable comments that have improved its readability. In particular we are grateful to him/her for drawing our attention to the concept of asymptotic stability.

Appendix A. A more conventional solution to (1.7) for the case $\varepsilon_1 \neq 1$

We have stated in the Introduction that when $\varepsilon_1 \neq 1$ a one soliton solution with the usual dispersion relation ($\omega = \frac{k}{1-k^2}$) cannot be found by the Hirota method. However, one can construct it using a more direct method. Since, we are interested in traveling wave solutions we use the ansatz

$$q = f(\xi); \quad \xi = kx - \omega t \quad (\text{A.1})$$

in the equation (1.7), which then leads to the following ordinary differential equation

$$\left(\omega^2 - k\omega\right) f_{\xi\xi\xi} - \left(4k^2\omega^2 + 2\varepsilon_2 k^2\omega^2\right) f_{\xi\xi}^2 - \left(\omega k^3 - \varepsilon_1 \omega k^3 + \varepsilon_1 \omega^2 k^2\right) f_{\xi\xi\xi\xi} = 0 \quad (\text{A.2})$$

where the subindices ξ denote the derivatives w.r.t. ξ . Since, we want to keep the same dispersion relation as in the RLW and mRLW equations, we choose ω to have the form $\omega = \frac{k}{1-k^2}$. Putting this into (A.2), dropping an overall factor ω , and denoting $f_{\xi\xi\xi} \equiv g$ we get the equation

$$\frac{k^3}{1-k^2} g - (4 + 2\varepsilon_2) \frac{k^3}{1-k^2} g^2 - \left[\varepsilon_1 \frac{k^3}{1-k^2} + (1 - \varepsilon_1) k^3 \right] g_{\xi\xi} = 0. \quad (\text{A.3})$$

Dividing (A.3) by the factor $\frac{k^3}{1-k^2}$ we note that it can be written as

$$g - G g^2 - H g_{\xi\xi} = 0 \quad \text{with} \quad G = 2(2 + \varepsilon_2); \quad H = 1 + (\varepsilon_1 - 1) k^2. \quad (\text{A.4})$$

Note that by rescaling the function g and the variable ξ we can change the coefficients G and H as we wish. In addition, the second derivative of \cosh^{-2} produce terms proportional to \cosh^{-2} and \cosh^{-4} , which can be cancelled by the g and g^2 in the equation. So, we take the ansatz

$$g = \frac{R}{\cosh^2(\alpha\xi)}. \tag{A.5}$$

Substituting it into (A.4) we get the equation

$$\frac{R}{2\cosh^4(\alpha\xi)} \left[1 + 8\alpha^2 H - 2GR + (1 - 4\alpha^2 H) \cosh(2\alpha\xi) \right] = 0, \tag{A.6}$$

which implies that

$$\alpha = \frac{1}{2\sqrt{H}}; \quad \text{and} \quad R = \frac{3}{2G}. \tag{A.7}$$

Thus

$$g = \frac{3}{4(2 + \varepsilon_2)} \cosh^{-2} \left(\frac{\xi}{2\sqrt{1 + (\varepsilon_1 - 1)k^2}} \right) \tag{A.8}$$

Integrating this expression twice we find that

$$q = \frac{3[1 + (\varepsilon_1 - 1)k^2]}{(2 + \varepsilon_2)} \left[\ln \left(\cosh \left(\frac{\xi}{2\sqrt{1 + (\varepsilon_1 - 1)k^2}} \right) \right) + b\xi + c \right] \tag{A.9}$$

with b and c being arbitrary constants. One needs the condition

$$1 + (\varepsilon_1 - 1)k^2 > 0 \tag{A.10}$$

for this solution to be real.

Appendix B. Numerical methods

B.1. Numerical approximation of the deformations of the KdV equation

In order to perform numerical simulations of equation (1.7) we follow the techniques discussed by J.C. Eilbeck and G.R. McGuire [12]. The equation necessitates the introduction of implicit methods. Hence first of all we introduce a new field $p(x, t)$ by

$$p = q_t, \tag{B.1}$$

so that equation (1.7) can be written as

$$p_t + p_x - 4p_x^2 - 2\varepsilon_2 q_{xx} p_t + p_{xxx} - \varepsilon_1 (p_{xxt} + p_{xxx}) = 0. \tag{B.2}$$

We see that the main difference in approximating this equation compared to the numerical scheme used in [6] is an extra term proportional to p_{xxx} .

In order to adjust our scheme to this extra term, we first discretize in both x and t by taking a finite set of points x_0, x_1, \dots, x_N and t_0, t_1, \dots, t_K , where h and τ denote the step size in space and time. Furthermore, we denote the grid points as $(ih, m\tau) \equiv (i, m)$, where $i = 0, 1, 2, \dots, N$ and $m = 0, 1, 2, \dots, K$, and we employ the notation $p_i^m \equiv p(ih, m\tau)$ and $q_i^m \equiv q(ih, m\tau)$. Finally, let v_i^m denote our approximation to p_i^m and let w_i^m denote our approximation to q_i^m .

Next we introduce the following central finite difference operators by their actions on v_i^m :

$$\delta_x^2 v_i^m = (v_{i+1}^m - 2v_i^m + v_{i-1}^m)/h^2, \tag{B.3}$$

$$H_x v_i^m = (v_{i+1}^m - v_{i-1}^m)/2h, \tag{B.4}$$

$$H_t v_i^m = (v_i^{m+1} - v_i^{m-1})/2\tau, \tag{B.5}$$

and similarly on w_i^m . Applying these operators to equation (B.2) in a straightforward manner we get

$$H_t v_i^m + H_x v_i^m - 4(H_x v_i^m)^2 - 2\varepsilon_2 \delta_x^2 w_i^m H_t v_i^m + \delta_x^2 H_x v_i^m - \varepsilon_1 (\delta_x^2 H_t v_i^m + \delta_x^2 H_x v_i^m) = 0, \tag{B.6}$$

which can be rewritten as

$$\begin{aligned} & \frac{v_i^{m+1}}{2\tau} - \varepsilon_1 \frac{v_{i+1}^{m+1} - 2v_i^{m+1} + v_{i-1}^{m+1}}{2h^2\tau} - \varepsilon_2 \frac{w_{i+1}^m v_i^{m+1} - 2w_i^m v_i^{m+1} + w_{i-1}^m v_i^{m+1}}{h^2\tau} \\ &= \frac{v_i^{m-1}}{2\tau} - \frac{v_{i+1}^m - v_{i-1}^m}{2h} - \varepsilon_1 \frac{v_{i+1}^{m-1} - 2v_i^{m-1} + v_{i-1}^{m-1}}{2h^2\tau} \\ & \quad - \varepsilon_2 \frac{w_{i+1}^m v_i^{m-1} - 2w_i^m v_i^{m-1} + w_{i-1}^m v_i^{m-1}}{h^2\tau} \\ & \quad + \frac{(v_{i+1}^m)^2 - 2v_{i+1}^m v_{i-1}^m + (v_{i-1}^m)^2}{h^2} \\ & \quad - (1 - \varepsilon_1) \frac{v_{i+2}^m - 2v_{i+1}^m + 2v_{i-1}^m - v_{i-2}^m}{2h^3}. \end{aligned} \tag{B.7}$$

Next we introduce the following matrices and vectors

$$A \equiv \begin{pmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & \frac{1}{2h^2\tau} & a_2^m & \frac{1}{2h^2\tau} & 0 & \vdots & \vdots \\ 0 & 0 & \frac{1}{2h^2\tau} & a_3^m & \frac{1}{2h^2\tau} & & \\ \vdots & & & \ddots & \ddots & \ddots & 0 & 0 \\ & & & & \frac{1}{2h^2\tau} & a_{N-2}^m & \frac{1}{2h^2\tau} & 0 \\ 0 & 0 & \dots & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & \dots & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \tag{B.8}$$

$$B \equiv \begin{pmatrix} v_0^{m+1} \\ v_1^{m+1} \\ v_2^{m+1} \\ \vdots \\ v_{N-1}^{m+1} \\ v_N^{m+1} \end{pmatrix}, \quad C \equiv \begin{pmatrix} c_0^m \\ c_1^m \\ c_2^m \\ \vdots \\ c_{N-1}^m \\ c_N^m \end{pmatrix}, \tag{B.9}$$

where

$$a_i^m \equiv \frac{1}{2\tau} - \varepsilon_2 \frac{w_{i+1}^m - 2w_i^m + w_{i-1}^m}{h^2\tau} + \varepsilon_1 \frac{v_i^{m+1}}{h^2\tau}, \tag{B.10}$$

$$\begin{aligned}
 c_i^m \equiv & \frac{v_i^{m-1}}{2\tau} - \frac{v_{i+1}^m - v_{i-1}^m}{2h} - \varepsilon_1 \frac{v_{i+1}^{m-1} - 2v_i^{m-1} + v_{i-1}^{m-1}}{2h^2\tau} \\
 & - \varepsilon_2 \frac{w_{i+1}^m v_i^{m-1} - 2w_i^m v_{i-1}^{m-1} + w_{i-1}^m v_i^{m-1}}{h^2\tau} \\
 & + \frac{(v_{i+1}^m)^2 - 2v_{i+1}^m v_{i-1}^m + (v_{i-1}^m)^2}{h^2} \\
 & - (1 - \varepsilon_1) \frac{v_{i+2}^m - 2v_{i+1}^m + 2v_{i-1}^m - v_{i-2}^m}{2h^3}, \quad i = 2, 3, \dots, N - 2,
 \end{aligned} \tag{B.11}$$

with the following boundary conditions

$$c_0^m \equiv v_0^{m+1}, \quad c_1^m \equiv v_1^{m+1}, \tag{B.12}$$

$$c_{N-1}^m \equiv v_{N-1}^{m+1}, \quad c_N^m \equiv v_N^{m+1}. \tag{B.13}$$

Note that we need 2 elements per boundary, which is a consequence of the term proportional to p_{xxx} . Having introduced this notation, our problem reduces to solving the following matrix equation

$$AB = C, \tag{B.14}$$

where we need to solve for vector B . In our algorithm we have used the well-known LU decomposition method for this problem, see for instance the book [37] for more information.

Once we have solved for B , we have the values of v_i^m at the next time level. We then solve equation (B.1) to determine all the values of w_i^m at the next time level

$$v_i^m = \frac{w_i^{m+1} - w_i^{m-1}}{2\tau} \implies w_i^{m+1} = 2\tau v_i^m + w_i^{m-1}. \tag{B.15}$$

Repeating this procedure for many time levels allows us to determine the numerical time evolution of a system.

It is not too difficult to verify that this scheme is both second-order accurate in τ and in h . Furthermore, this scheme is an extension of the scheme discussed in [6], and so that gives us an indication that we can trust the results of our simulations.

B.2. Summary of parameters used to produce figures

Tables 1, 2 and 3 show all the parameters used to produce the figures shown in this paper.

Appendix C. The $\mathfrak{sl}(2)$ loop algebra

We use the $\mathfrak{sl}(2)$ finite algebra with commutation relations

$$[T_3, T_{\pm}] = \pm T_{\pm} \quad \text{and} \quad [T_+, T_-] = 2T_3, \tag{C.1}$$

which is satisfied by the basis

$$T_i = \frac{1}{2}\sigma_i \quad \text{and} \quad T_{\pm} = T_1 \pm iT_2, \tag{C.2}$$

where σ_i , are the Pauli matrices. To proceed further, we take the following basis for the corresponding loop algebra

Table 1

This table summarizes the variables used to produce Figs. 1 to 8.

	Figs. 1 and 2	Figs. 3 and 4	Figs. 5 and 6	Figs. 7 and 8
x_0	-50	-200	-50	-200
x_N	250	200	250	200
h	0.1	0.1	0.1	0.1
t_0	0	0	0	0
t_K	40	70	40	70
τ	0.001	0.001	0.001	0.001
ω_1	5.00	0.80	5.00	0.80
δ_1	0.00	27.71	0.00	27.71
ω_2	3.00	3.07	3.00	3.07
δ_2	-40.00	106.28	-40.00	106.28
ω_3	N/A	4.28	N/A	4.28
δ_3	N/A	148.34	N/A	148.34

Table 2

This table summarizes the variables used to produce Figs. 9 to 19.

	Figs. 9, 10 and 11	Figs. 12, 13 and 14	Figs. 15, 16, 17, 18 and 19
x_0	-50	-200	-50
x_N	300	200	300
h	0.1	0.1	0.1
t_0	0	0	0
t_K	40	70	40
τ	0.001	0.001	0.001
ω_1	5.00	0.80	5.00
δ_1	0.00	27.71	0.00
ω_2	3.00	3.07	3.00
δ_2	-40.00	106.28	-40.00
ω_3	N/A	4.28	N/A
δ_3	N/A	148.34	N/A

Table 3

This table summarizes the variables used to produce Figs. 20 to 25.

	Figs. 20, 21 and 22	Figs. 23, 24 and 25
x_0	-50	-200
x_N	300	300
h	0.1	0.1
t_0	0	0
t_K	40	70
τ	0.001	0.001
ω_1	5.00	0.80
δ_1	0.00	27.71
ω_2	3.00	3.07
δ_2	-40.00	106.28
ω_3	N/A	4.28
δ_3	N/A	148.34

$$b_{2m+1} = \lambda^m (T_+ + \lambda T_-), \quad (\text{C.3})$$

and

$$F_{2m+1} = \lambda^m (T_+ - \lambda T_-) \quad \text{and} \quad F_{2m} = 2\lambda^m T_3, \quad (\text{C.4})$$

where $m \in \mathbb{Z}$, and λ is the so-called spectral parameter. The algebra is then given by

$$[b_{2m+1}, b_{2n+1}] = 0, \tag{C.5}$$

$$[F_{2m+1}, F_{2n+1}] = 0, \tag{C.6}$$

$$[F_{2m}, F_{2n}] = 0, \tag{C.7}$$

$$[b_{2m+1}, F_{2n+1}] = -2F_{2(m+n+1)}, \tag{C.8}$$

$$[b_{2m+1}, F_{2n}] = -2F_{2(m+n)+1}, \tag{C.9}$$

$$[F_{2m+1}, F_{2n}] = -2b_{2(m+n)+1}. \tag{C.10}$$

We introduce the grading operator as

$$d = T_3 + 2\lambda \frac{d}{d\lambda} \tag{C.11}$$

such that

$$[d, b_{2m+1}] = (2m + 1) b_{2m+1} \quad \text{and} \quad [d, F_m] = m F_m. \tag{C.12}$$

The important ingredient in such procedure is that the generator b_1 in A_x given in (2.1), is a semi-simple element of the algebra in the sense that its adjoint action on the loop algebra, splits it into kernel and image with a trivial intersection, i.e.

$$\mathcal{G} = \text{Ker} + \text{Im}; \quad [b_1, \text{Ker}] = 0; \quad \text{Im} = [b_1, \mathcal{G}]; \quad \text{Ker} \cap \text{Im} = 0 \tag{C.13}$$

and we have that

$$\text{Ker} = \{b_{2n+1} \mid n \in \mathbb{Z}\}; \quad \text{Im} = \{F_n \mid n \in \mathbb{Z}\} \tag{C.14}$$

Appendix D. The parameters of the gauge transformation (2.7)

We give here the explicit expressions for the parameters of the group element performing the gauge transformation (2.7):

$$\begin{aligned} \zeta_1 &= 0 \\ \zeta_2 &= -\frac{1}{24}\alpha u \\ \zeta_3 &= \frac{1}{48}\alpha u^{(1,0)} \\ &= \partial_x \left[\frac{1}{48}\alpha u \right] \\ \zeta_4 &= \frac{1}{288} \left[-\alpha^2 u^2 - 3\alpha u^{(2,0)} \right] \\ \zeta_5 &= \frac{1}{192} \left[\alpha^2 u u^{(1,0)} + \alpha u^{(3,0)} \right] \\ &= \partial_x \left[\frac{1}{192} \left[\frac{\alpha^2}{2} u^2 + \alpha u_{xx} \right] \right] \\ \zeta_6 &= \frac{1}{10368} \left[-4\alpha^3 u^3 - 36\alpha^2 u^{(2,0)} u - 27\alpha^2 \left(u^{(1,0)} \right)^2 - 27\alpha u^{(4,0)} \right] \end{aligned} \tag{D.1}$$

$$\begin{aligned} \zeta_7 &= \frac{1}{20736} \left[22\alpha^3 u^2 u^{(1,0)} + 90\alpha^2 u^{(1,0)} u^{(2,0)} + 45\alpha^2 u u^{(3,0)} + 27\alpha u^{(5,0)} \right] \\ &= \partial_x \left[\frac{1}{20736} \left[\frac{22}{3} \alpha^3 u^3 + \frac{45}{2} \alpha^2 u_x^2 + 45\alpha^2 u u_{xx} + 27\alpha u_{xxxx} \right] \right] \\ \zeta_8 &= \frac{1}{41472} \left[-2\alpha^4 u^4 - 36\alpha^3 u^{(2,0)} u^2 - 53\alpha^3 \left(u^{(1,0)} \right)^2 u - 54\alpha^2 u^{(4,0)} u - 90\alpha^2 \left(u^{(2,0)} \right)^2 \right. \\ &\quad \left. - 135\alpha^2 u^{(1,0)} u^{(3,0)} - 27\alpha u^{(6,0)} \right] \end{aligned}$$

where we have used the notation

$$\partial_x^m \partial_t^n u \equiv u^{(m,n)} \tag{D.2}$$

The explicit expressions for the first few components of the time component a_t of transformed connection given in (2.17) are given by

$$\begin{aligned} a_t^{(-1)} &= -\frac{\alpha^2}{72} u^2 - \frac{G}{2} \\ a_t^{(-3)} &= \frac{\alpha^2}{288} \left(u^{(1,0)} \right)^2 + \frac{\alpha^2}{288} u^2 - \frac{\alpha}{24} G u \\ a_t^{(-5)} &= \frac{\alpha^4 u^4}{13824} + \frac{\alpha^3 u^{(2,0)} u^2}{1728} + \frac{\alpha^3 u^3}{1728} - \frac{1}{192} \alpha^2 G u^2 + \frac{\alpha^2 u^{(1,1)} u}{1152} + \frac{1}{576} \alpha^2 u^{(2,0)} u \\ &\quad - \frac{\alpha^2 \left(u^{(1,0)} \right)^2}{1152} + \frac{\alpha^2 \left(u^{(2,0)} \right)^2}{1152} - \frac{\alpha^2 u^{(0,1)} u^{(1,0)}}{1152} - \frac{1}{96} \alpha G u^{(2,0)} \\ a_t^{(-7)} &= \frac{\alpha^5 u^5}{62208} + \frac{5\alpha^4 u^{(2,0)} u^3}{20736} - \frac{5\alpha^4 \left(u^{(1,0)} \right)^2 u^2}{82944} + \frac{5\alpha^4 u^4}{55296} - \frac{5\alpha^3 G u^3}{6912} + \frac{\alpha^3 u^{(1,1)} u^2}{3456} \\ &\quad + \frac{5\alpha^3 u^{(2,0)} u^2}{6912} + \frac{\alpha^3 u^{(4,0)} u^2}{6912} + \frac{\alpha^3 \left(u^{(2,0)} \right)^2 u}{1728} - \frac{\alpha^3 u^{(0,1)} u^{(1,0)} u}{6912} - \frac{\alpha^3 u^{(1,0)} u^{(3,0)} u}{2304} \\ &\quad + \frac{\alpha^3 \left(u^{(1,0)} \right)^2 u^{(2,0)}}{2304} - \frac{5\alpha^2 G u^{(2,0)} u}{1152} - \frac{5\alpha^2 G \left(u^{(1,0)} \right)^2}{2304} + \frac{\alpha^2 u^{(3,1)} u}{4608} + \frac{\alpha^2 u^{(4,0)} u}{2304} \\ &\quad + \frac{\alpha^2 \left(u^{(2,0)} \right)^2}{4608} - \frac{\alpha^2 \left(u^{(3,0)} \right)^2}{4608} + \frac{\alpha^2 u^{(1,1)} u^{(2,0)}}{4608} - \frac{\alpha^2 u^{(1,0)} u^{(2,1)}}{4608} - \frac{\alpha^2 u^{(0,1)} u^{(3,0)}}{4608} \\ &\quad - \frac{\alpha^2 u^{(1,0)} u^{(3,0)}}{2304} + \frac{\alpha^2 u^{(2,0)} u^{(4,0)}}{2304} - \frac{\alpha G u^{(4,0)}}{384} \end{aligned}$$

and

$$\begin{aligned} c_t^{(2)} &= 0 \\ c_t^{(1)} &= 0 \\ c_t^{(0)} &= 0 \\ c_t^{(-1)} &= -\frac{\alpha^2}{36} u^2 - \frac{\alpha}{12} u^{(2,0)} - \frac{\alpha}{12} u + \frac{G}{2} \\ &= \frac{X}{2} \\ c_t^{(-2)} &= \frac{\alpha^2}{24} u u^{(1,0)} + \frac{\alpha}{24} u^{(0,1)} + \frac{\alpha}{24} u^{(1,0)} + \frac{\alpha}{24} u^{(3,0)} \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{4} X_x \\
c_t^{(-3)} &= -\frac{1}{432} \alpha^3 u^3 - \frac{1}{36} \alpha^2 u^{(2,0)} u - \frac{1}{48} \alpha^2 \left(u^{(1,0)} \right)^2 - \frac{1}{144} \alpha^2 u^2 + \frac{1}{24} \alpha G u \\
&\quad - \frac{1}{48} \alpha u^{(1,1)} - \frac{1}{48} \alpha u^{(2,0)} - \frac{1}{48} \alpha u^{(4,0)} \\
&= \frac{\alpha}{24} u X + \frac{1}{8} X_{xx}
\end{aligned} \tag{D.3}$$

References

- [1] H. Aratyn, L.A. Ferreira, J.F. Gomes, A.H. Zimerman, The conserved charges and integrability of the conformal affine Toda models, *Mod. Phys. Lett. A* 9 (1994) 2783, arXiv:hep-th/9308086.
- [2] A. Arnaudon, On a Lagrangian reduction and a deformation of completely integrable systems, *J. Nonlinear Sci.* 26 (2016) 1133–1160, <https://doi.org/10.1007/s00332-016-9300-2>.
- [3] V.H. Aurichio, L.A. Ferreira, Quasi-integrable deformations of the Bullough–Dodd model, *J. High Energy Phys.* 1503 (2015) 152, [https://doi.org/10.1007/JHEP03\(2015\)152](https://doi.org/10.1007/JHEP03(2015)152), arXiv:1501.01821 [hep-th].
- [4] T.B. Benjamin, The stability of solitary waves, *Proc. R. Soc. Lond. A* 328 (1972) 153–183.
- [5] T.B. Benjamin, J.L. Bona, J.J. Mahoney, Model equations for long waves in nonlinear dispersive systems, *Philos. Trans. R. Soc. A* 272 (1972) 47–78.
- [6] F. ter Braak, W.J. Zakrzewski, Aspects of modified regularized long-wave equation, arXiv:1707.00669 [nlin.PS], 2017.
- [7] A. Bryan, A. Stuart, Solitons and the regularized long wave equation: a nonexistence theorem, *Chaos Solitons Fractals* 7 (11) (1996) 1881–1886.
- [8] F. Calogero, in: V.E. Zakharov (Ed.), *What Is Integrability?*, Springer-Verlag, 1991.
- [9] L.A. Dickey, *Soliton Equations and Hamiltonian Systems*, Advanced Series in Mathematical Physics, vol. 12, World Scientific Publishing Co. Pte. Ltd., 1991.
- [10] V.G. Drinfeld, V.V. Sokolov, Lie algebras and equations of Korteweg–de Vries type, *J. Sov. Math.* 30 (1984) 1975; Equations of Korteweg–de Vries type and simple Lie algebras, *Sov. Math. Dokl.* 23 (1981) 457.
- [11] M. Dunajski, Integrable systems, [www.damtp.cam.ac.uk](http://www.damtp.cam.ac.uk/user/md327/ISlecture_notes_2012.pdf), [online]. Available at: http://www.damtp.cam.ac.uk/user/md327/ISlecture_notes_2012.pdf, 2017. (Accessed 16 May 2017).
- [12] J.C. Eilbeck, G.R. McGuire, Numerical study of the regularized long-wave equation I: numerical methods, *J. Comput. Phys.* 19 (1) (1975) 43–57.
- [13] J.C. Eilbeck, G.R. McGuire, Numerical study of the regularized long-wave equation II: interaction of solitary waves, *J. Comput. Phys.* 23 (1) (1977) 63–73.
- [14] L.A. Ferreira, W.J. Zakrzewski, The concept of quasi-integrability: a concrete example, *J. High Energy Phys.* 1105 (2011) 130, [https://doi.org/10.1007/JHEP05\(2011\)130](https://doi.org/10.1007/JHEP05(2011)130), arXiv:1011.2176 [hep-th].
- [15] L.A. Ferreira, G. Luchini, W.J. Zakrzewski, The concept of quasi-integrability for modified non-linear Schrödinger models, *J. High Energy Phys.* 1209 (2012) 103, [https://doi.org/10.1007/JHEP09\(2012\)103](https://doi.org/10.1007/JHEP09(2012)103), arXiv:1206.5808 [hep-th].
- [16] L.A. Ferreira, W.J. Zakrzewski, Numerical and analytical tests of quasi-integrability in modified sine-Gordon models, *J. High Energy Phys.* 1401 (058) (2014), [https://doi.org/10.1007/JHEP01\(2014\)058](https://doi.org/10.1007/JHEP01(2014)058), arXiv:1308.4412 [hep-th].
- [17] L.A. Ferreira, W.J. Zakrzewski, Breather-like structures in modified sine-Gordon models, *Nonlinearity* 29 (2016) 1622–1644, arXiv:1404.5812 [hep-th].
- [18] L.A. Ferreira, P. Klimas, W.J. Zakrzewski, Quasi-integrable deformations of the $SU(3)$ affine Toda theory, *J. High Energy Phys.* 1605 (065) (2016), [https://doi.org/10.1007/JHEP05\(2016\)065](https://doi.org/10.1007/JHEP05(2016)065), arXiv:1602.02003 [hep-th].
- [19] J.D. Gibbon, J.C. Eilbeck, R.K. Dodd, A modified regularized long-wave equation with an exact two-soliton solution, *J. Phys. A* 9 (1976) L127–L130.
- [20] J. Hietarinta, A search for bilinear equations passing Hirota’s three-soliton condition. I. KdV-type bilinear equations, *J. Math. Phys.* 28 (8) (1987) 1732–1742.
- [21] Yuri S. Kivshar, Boris A. Malomed, Dynamics of solitons in nearly integrable systems, *Rev. Mod. Phys.* 61 (4) (1989) 763–915.

- [22] Y. Kodama, A.V. Mikhailov, Obstacles to asymptotic integrability, in: A.S. Fokas, I.M. Gelfand (Eds.), *Algebraic Aspects of Integrable Systems*, Birkhäuser, 1997, pp. 173–204.
- [23] D.J. Korteweg, G. de Vries, On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves, *Philos. Mag.* 39 (1895) 422–443.
- [24] S. Kutluay, A. Esen, A finite difference solution of the regularized long-wave equation, *Math. Probl. Eng.* 1 (2006).
- [25] P.D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Commun. Pure Appl. Math.* 21 (1968) 467–490;
V.E. Zakharov, A.B. Shabat, *Zh. Exp. Teor. Fiz.* 61 (1971) 118–134, *Sov. Phys. JETP* 34 (1972) 62–69, English transl.
- [26] Y. Martel, Asymptotic N-soliton-like solutions of the subcritical and critical gKdV equations, *Amer. Math. J.* 127 (2005) 1103–1140.
- [27] Y. Martel, F. Merle, Inelastic interaction of nearly equal solitons for the BBM equation, *Discrete Contin. Dyn.* 27 (2010) 487–532.
- [28] Y. Martel, F. Merle, T. Mizumachi, Description of the inelastic collision of two solitary waves for the BBM equation, *Arch. Ration. Mech. Anal.* 196 (2010) 517–574, <https://doi.org/10.1007/s00205-009-0244-7>.
- [29] Judith R. Miller, Michael I. Weinstein, Asymptotic stability of solitary waves for the regularized long-wave equation, *Commun. Pure Appl. Math.* XLIX (1996) 399–441.
- [30] Tetsu Mizumachi, Asymptotic stability of solitary wave solutions to the regularized long-wave equation, *J. Differ. Equ.* 200 (2004) 312–341, <https://doi.org/10.1016/j.jde.2004.01.006>.
- [31] M. Musette, F. Lambert, J.C. Decuyper, Soliton and antisoliton resonant interactions, *J. Phys. A, Math. Gen.* 20 (18) (1987) 6223–6235.
- [32] D.I. Olive, N. Turok, Local conserved densities and zero curvature conditions for Toda lattice field theories, *Nucl. Phys. B* 257 (1985) 277;
The Toda lattice field theory hierarchies and zero curvature conditions in Kac–Moody algebras, *Nucl. Phys. B* 265 (1986) 469.
- [33] A. Parker, On exact solutions of the regularized long-wave equation: a direct approach to partially integrable equations. I. Solitary wave and solitons, *J. Math. Phys.* 36 (7) (1995) 3498–3505.
- [34] Robert L. Pego, Michael I. Weinstein, On asymptotic stability of solitary waves, *Phys. Lett. A* 162 (1992) 263–268.
- [35] Robert L. Pego, Michael I. Weinstein, Asymptotic stability of solitary waves, *Commun. Math. Phys.* 164 (1994) 305–349.
- [36] D.H. Peregrine, Calculations of the development of an undular bore, *J. Fluid Mech.* 25 (1966) 321–330.
- [37] G. Schay, *A Concise Introduction to Linear Algebra*, Birkhäuser, Boston, 2012.
- [38] Terence Tao, Why are solitons stable?, *Bull. Am. Math. Soc.* 46 (1) (2009) 1–33.
- [39] N.J. Zabusky, M.D. Kruskal, Interaction of ‘Solitons’ in a collisionless plasma and the recurrence of initial states, *Phys. Rev. Lett.* 15 (1965) 240, <https://doi.org/10.1103/PhysRevLett.15.240>.