


First Observation of a Three-Resonance Structure in $e^+e^- \rightarrow$ Nonopen Charm HadronsM. Ablikim *et al.**
(BESIII Collaboration) (Received 19 July 2023; revised 10 March 2024; accepted 29 March 2024; published 9 May 2024)

We report the measurement of the inclusive cross sections for $e^+e^- \rightarrow$ nOCH (where nOCH denotes non-open charm hadrons) with improved precision at center-of-mass (c.m.) energies from 3.645 to 3.871 GeV. We observe three resonances: $\mathcal{R}(3760)$, $\mathcal{R}(3780)$, and $\mathcal{R}(3810)$ with significances of 8.1σ , 13.7σ , and 8.8σ , respectively. The $\mathcal{R}(3810)$ state is observed for the first time, while the $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states are observed for the first time in the nOCH cross sections. Two sets of resonance parameters describe the energy-dependent line shape of the cross sections well. In set I [set II], the $\mathcal{R}(3810)$ state has mass $(3805.7 \pm 1.1 \pm 2.7)$ [(3805.7 \pm 1.1 \pm 2.7)] MeV/ c^2 , total width $(11.6 \pm 2.9 \pm 1.9)$ [(11.5 \pm 2.8 \pm 1.9)] MeV, and an electronic width multiplied by the nOCH decay branching fraction of $(10.9 \pm 3.8 \pm 2.5)$ [(11.0 \pm 3.4 \pm 2.5)] eV. In addition, we measure the branching fractions $\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{nOCH}] = (25.2 \pm 16.1 \pm 30.4)\%$ [(6.4 \pm 4.8 \pm 7.7)%] and $\mathcal{B}[\mathcal{R}(3780) \rightarrow \text{nOCH}] = (12.3 \pm 6.6 \pm 8.3)\%$ [(10.4 \pm 4.8 \pm 7.0)%] for the first time. The $\mathcal{R}(3760)$ state can be interpreted as an open-charm (OC) molecular state, but containing a simple four-quark state component. The $\mathcal{R}(3810)$ state can be interpreted as a hadrocharmonium state.

DOI: [10.1103/PhysRevLett.132.191902](https://doi.org/10.1103/PhysRevLett.132.191902)

Until two decades ago, it was widely believed that hadron resonances with masses higher than the open-charm (OC) pair thresholds decay entirely to OC final states via the strong interaction. However, in July, 2003, BES reported the observation of seven events of hadron resonance(s) in this mass regime decaying to nOCH [1–4]. This discovery overturned the understanding of resonance decays and opened up a new era in hadron spectroscopy. In this Letter, we denote these resonances as X_{aboveOC} [3], which encompasses both supposed pure quark-antiquark $c\bar{c}$ states, i.e., $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$, and nonpure quark-antiquark $c\bar{c}$ states (hereafter referred to as ‘non- $c\bar{c}$ ’), such as four-quark states, OC-pair molecular states, hadrocharmonium states, and hybrid charmonium states [5–8]. Quantum chromodynamics (QCD) expects that the non- $c\bar{c}$ states exist in nature, and thus a discovery of these systems would be an important validation of the QCD predictions. The first nOCH final-state decay of a X_{aboveOC} resonance to be observed after the 2002 discoveries was $J/\psi\pi^+\pi^-$ [1,2,9], seen by BES-II. This final state can originate from a $c\bar{c}$ state, a non- $c\bar{c}$ state, or a combination of both [4]. This discovery stimulated strong interest in using $J/\psi\pi^+\pi^-$ or similar final states as golden

channels to probe other nOC decays from the X_{aboveOC} , and led to the discovery of several exotic states [10–20].

Subsequent studies of the $\psi(3770)$ resonance showed that it has a branching fraction of about 15% into nOCH final states [21–24]. This fact indicates the contribution of some undiscovered states [25] with masses around 3.773 GeV/ c^2 , which predominantly decay into nOCH. In 2008, the BES-II experiment observed for the first time a double-peaked structure named $\mathcal{R}_s(3770)$ in $e^+e^- \rightarrow$ hadrons at c.m. energies around 3.76 GeV [26], which is composed of two states labeled $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$. The existence of the $\mathcal{R}(3760)$ state was confirmed in $e^+e^- \rightarrow J/\psi X$ by BESIII [4]. It is seen that the study of the inclusive nOCH decays of X_{aboveOC} both helps the understanding of known states, and is a sensitive probe for undiscovered resonances, particularly non- $c\bar{c}$ states.

In this Letter, we report a measurement of the cross sections for $e^+e^- \rightarrow$ nOCH at c.m. energies from 3.645 to 3.871 GeV, studies of $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ production and decays, and a search for an additional resonant state in this energy region. The data samples used in this analysis were collected at 42 c.m. energies in 2010 and correspond to a total integrated luminosity of 75.5 pb $^{-1}$.

The BESIII detector [27] response is studied using Monte Carlo (MC) samples. The simulations are performed using a GEANT4-based [28] software package. Simulated samples for $q\bar{q}$ vector states (i.e. $u\bar{u}$, $d\bar{d}$, $s\bar{s}$, and $c\bar{c}$) and their decays to hadrons are generated using the MC event generators KKMC [29], EVTGEN [30], and LUNDCHARM [31]. Background sources are estimated with MC samples

*Full author list given at the end of the Letter.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP³.

generated using KKMC, and the MC event generators BABAYAGA [32] and TWOGAM [33].

We select the inclusive nOCH events from the events with charged particles or charged and neutral particles. In order to reject background contributions from $e^+e^- \rightarrow (\gamma)e^+e^-$ and $e^+e^- \rightarrow (\gamma)\mu^+\mu^-$, we require the events to have more than two charged tracks (N_{CTrk}), and impose the following selection criteria for each track: (i) the distance (R_{xy}) of the point of the closest approach to the beam pipe must satisfy the condition $R_{xy} \leq 1.0$ cm; (ii) the polar angle θ must satisfy the condition $|\cos\theta| < 0.93$; (iii) the momentum p must be less than $E_b + 0.02E_b\sqrt{1 + E_b^2}$, where E_b is the beam energy; (iv) the time-of-flight t_{TOF} must satisfy $2.0 < t_{\text{TOF}} < 20.0$ ns and $|t_{\text{TOF}} - t_p| < 2.0$ ns, where t_p is the expected time-of-flight of protons; (v) the energy E_{EMC} deposited in the electromagnetic calorimeter (EMC) must be less than 1 GeV; (vi) the penetration depth in the muon-chamber system must be less than 30 cm.

For the selection of photons, we require the deposited energy of a neutral cluster in the EMC to be greater than 25 MeV in the barrel and 50 MeV in the end caps. We do not apply any requirements on the number of photons in the event. To suppress electronic noise and showers unrelated to the event, we impose the condition that the difference between the EMC time and the event start time be within [0,700] ns. To reduce the beam-associated events (beam interactions with gas or material), we demand that at least one charged track or photon must point into each hemisphere of $\cos\theta < 0$ and $\cos\theta > 0$. In addition, for each event, we require the total energy ($E_{\text{EMC}}^{\text{tot}}$) deposited in the EMC by the charged and neutral particles to be greater than $0.28E_b$.

Some beam-associated background sources still survive this selection. These background sources are produced at random z positions [21], while genuine nOCH events are produced around $z = 0$, where z is the distance to the interaction point along the BESIII axis. To distinguish the nOCH events from the background sources, we calculate the averaged z (Z_{AVRG}) of the charged tracks in each event. Figure 1 shows the distribution of the averaged Z_{AVRG} of the accepted events from the data sample collected at $\sqrt{s} = 3.779$ GeV. Using a double-Gaussian function to describe the signal shape plus a second-order Chebychev function to parameterize the background shape, we fit the Z_{AVRG} distribution of event vertices to extract the number of nOCH candidates, $N_{\text{had}}^{\text{fit}}$ at each energy point.

The background to the $N_{\text{had}}^{\text{fit}}$ distribution comes from various sources, e.g., (i) $e^+e^- \rightarrow (\gamma)e^+e^-$, (ii) $e^+e^- \rightarrow (\gamma)\mu^+\mu^-$, (iii) $e^+e^- \rightarrow \gamma\gamma$, (iv) $e^+e^- \rightarrow (\gamma)\tau^+\tau^-$, (v) $e^+e^- \rightarrow (\gamma)e^+e^-\ell^+\ell^-$ ($\ell = e, \mu$ or τ), (vi) $e^+e^- \rightarrow (\gamma)e^+e^-X_{\text{had}}$ (where X_{had} denotes hadrons), and (vii) $e^+e^- \rightarrow (\gamma)D\bar{D}$. The total amount of background N_b at each c.m. energy \sqrt{s} is determined by $N_b = \sum_{i=1}^7 N_{b,i}$ with $N_{b,i} = \mathcal{L} \times \sigma_{b,i} \times \eta_i$, where \mathcal{L} is the




FIG. 1. Z_{AVRG} distribution of the vertices of the selected events from the data sample collected at $\sqrt{s} = 3.779$ GeV, where the dots with error bar represent data, the solid line in blue is the best fit, and the dashed line in red is the fitted background shape.

integrated luminosity of the data sample, $\sigma_{b,i}$ is the cross section for the i th background source and η_i is the probability of misidentifying a candidate from the i th background source as a nOCH event, which is determined by analyzing the large background MC samples. The cross sections σ_b for sources (i), (ii), and (iii) are taken from BABAYAGA. For source (iv), σ_b is calculated using the formulae given in Refs. [34,35]. The cross sections for sources (v) and (vi) are taken from TWOGAM, while that for source (vii) is taken from the observed cross sections $\sigma_{D\bar{D}}^o(s)$ for $e^+e^- \rightarrow D\bar{D}$ determined using the same data samples. For example, at $\sqrt{s} = 3.7731$ GeV, $N_{\text{had}}^{\text{fit}} = (35235 \pm 225)$, $N_{b,\ell^+\ell^-\gamma\gamma} = (1206 \pm 7)$, $N_{b,e^+e^-\ell^+\ell^-,e^+e^-X_{\text{had}}} = (341 \pm 6)$, and $N_{b,D\bar{D}} = (9458 \pm 186)$, where the uncertainties on the first two backgrounds arise from the statistical uncertainties on \mathcal{L} , $\sigma_{b,i}$, and η_i , and that on the third is due to the statistical uncertainty on $\sigma_{D\bar{D}}^o(s)$. To provide the most conservative signal significance for the resonance searches that are discussed below, we directly subtract N_b from $N_{\text{had}}^{\text{fit}}$ and allow the statistical uncertainty of $N_{b,D\bar{D}}$ to fully contribute to the uncertainty in the difference between $N_{\text{had}}^{\text{fit}}$ and N_b . This procedure yields $N_{\text{nOCH}}^o = (24230 \pm 292)$, where the uncertainty is statistical and includes the contribution from the background estimates.

We determine the detection efficiency using MC simulated events for the four components of the process $e^+e^- \rightarrow \text{nOCH}$: (i) $e^+e^- \rightarrow$ light-hadron (LH) continuum processes including lower-mass resonances (LMRs) with masses below 2 GeV (CPLMRs), (ii) $J/\psi \rightarrow$ hadrons, (iii) $\psi(3686) \rightarrow$ hadrons, and (iv) $\mathcal{R}s(3770) \rightarrow \text{nOCH}$ [26]. We generate simulated samples for these events with the KKMC package. These events include initial-state radiation (ISR) and final-state radiation (FSR) processes. For the subsequent decays of the J/ψ , $\psi(3686)$ and $\mathcal{R}s(3770)$, we use EVTGEN to generate the known final states, given in Ref. [36], and use LUNDCHARM to generate the remaining unknown final states. The resulting selection




FIG. 2. The efficiency ϵ (top) versus the c.m energy; the ISR correction factor $[1 + \delta(s)]$ (bottom) versus the c.m energy.

efficiency is determined using $\epsilon = \sum_1^4 w_i \epsilon_i$, where w_i is the number of nOCH events simulated for the i th component over the total number of MC simulated nOCH events for all four components, and ϵ_i is the corresponding efficiency for selection of the nOCH events from the i th component. Figure 2 (top) shows the efficiencies determined at the 42 energy points.

At $\sqrt{s} = 3.7731$ GeV, the efficiency is $\epsilon = 62.11\%$. The integrated luminosity corresponding to the data collected at this energy is $\mathcal{L} = (1831.63 \pm 4.49) \text{ nb}^{-1}$. Dividing the number $N_{\text{nOCH}}^0 = (24230 \pm 292)$ by both the integrated luminosity and the efficiency yields the observed nOCH cross section $\sigma_{\text{nOCH}}^0 = (21.299 \pm 0.262) \text{ nb}$, where the uncertainty arises from the statistical uncertainty on the N_{OCH}^0 , the size of the MC samples, and the statistical uncertainty of the luminosity measurement. Similarly, we measure $\sigma_{\text{nOCH}}^0(s)$ at the remaining 41 energy points.

Table I summarizes the systematic uncertainties assigned to the $\sigma_{\text{nOCH}}^0(s)$ measurements. To determine the systematic uncertainties due to the choice of the selection criteria, we vary each criterion from its baseline value to an alternative setting, as given in Table I, and assign the resulting change in $\sigma_{\text{nOCH}}^0(s)$ as the systematic uncertainty. Removing the requirement that tracks must point into different hemispheres in z changes $\sigma_{\text{nOCH}}^0(s)$ by 0.80%, which is taken as the corresponding uncertainty associated with this possible source of bias. To estimate the uncertainty associated with the fit to the Z_{AVRG} distribution, we reperform the fit changing the background shape from a second-order to

TABLE I. Systematic uncertainties on $\sigma_{\text{nOCH}}^0(s)$.

Source	Variation range	Uncertainty (%)
R_{xy} cut	[0.9, 1.1]	0.20
$\cos \theta$ cut	[0.875, 0.930]	0.40
$p < E_b + 0.02E_b\sqrt{1 + E_b}$	[nominal value, ∞]	0.01
t_{TOF} cut and $t_{\text{TOF}} - t_p$	[0, ∞] and [2, ∞]	0.09
E_{EMC} cut	[1, ∞]	0.39
Penetration depth	[30, ∞]	0.04
$E_{\text{EMC}}^{\text{tot}}$ cut	[0, 0.28]	0.34
N_{CTRk} cut	[2, 3]	1.45
Different hemisphere in z		0.80
Fitting Z_{AVRG} distribution		0.83
N_b		0.75
MC signal model		1.70
Integrated luminosity		1.00
Total		2.89

third-order, and then from a third-order to fourth-order Chebychev function. These result in 0.83% change in the cross sections, which is assigned as the corresponding systematic uncertainty. The uncertainty on the number of background sources is dominated by the estimation of the two-photon contribution, which induces a 0.75% uncertainty of the $\sigma_{\text{nOCH}}^0(s)$.

The choice of the MC generator for $e^+e^- \rightarrow \text{nOCH}$ impacts the selection efficiency. We recalculate the efficiency with different MC packages and take the 1.70% variation observed as the associated uncertainty. There is an uncertainty on $\sigma_{\text{nOCH}}^0(s)$ of 1.00% arising from the corresponding uncertainty on the luminosity. Adding these contributions in quadrature yields a total systematic uncertainty of 2.89% on $\sigma_{\text{nOCH}}^0(s)$. This total does not include an energy-dependent uncertainty on $\sigma_{\text{nOCH}}^0(s)$, caused by the 1.43% uncertainty on the $\sigma_{DD}^0(s)$ shape, which is accounted for when considering the systematic uncertainties on the fitted parameter values, as discussed below.

To investigate whether the $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states decay to nOCH and if a new resonance \mathcal{R} exists in this energy region, we perform a least- χ^2 fit to the nOCH cross sections. The dressed nOCH cross section is modeled by

$$\sigma_{\text{nOCH}}^{\text{D}}(s') = \sigma_{\text{LH}}^{\text{D}}(s') + \sigma_{J/\psi}^{\text{D}}(s') + \sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s'), \quad (1)$$

where $s' = s(1 - x)$, x is the radiative-photon energy fraction, $\sigma_{\text{LH}}^{\text{D}}(s')$, $\sigma_{J/\psi}^{\text{D}}(s')$ and $\sigma_{\mathcal{R}_{\text{Sup}3680}}^{\text{D}}(s')$ are the cross sections of $e^+e^- \rightarrow \text{LH}$, $J/\psi \rightarrow \text{hadrons}$, and $\mathcal{R}_{\text{Sup}3680} \rightarrow \text{nOCH}$, respectively. Here $\mathcal{R}_{\text{Sup}3680}$ indicates the states with masses above 3.680 GeV/ c^2 . The cross sections for CPLMRs are taken to be $\sigma_{\text{LH}}^{\text{D}}(s') = f\sigma_{\mu^+\mu^-}^{\text{B}}(s') + \sigma_{\text{LMRs}}^{\text{D}}(s')$, where $\sigma_{\mu^+\mu^-}^{\text{B}}(s')$ is the Born cross section for continuum $e^+e^- \rightarrow \mu^+\mu^-$ production, f is a free parameter, $f\sigma_{\mu^+\mu^-}^{\text{B}}(s')$ gives the cross section for continuum $e^+e^- \rightarrow \text{hadrons}$

production [24], and $\sigma_{\text{LMRS}}^{\text{D}}(s')$ is the cross section for the production of LMRs decaying into LH, which is determined using the zeroth-order cross sections [36] multiplied by the vacuum-polarization correction factor $(1/|1 - \Pi(s)|^2)$ [37,38] at energies below 2 GeV. The cross sections for J/ψ and $\mathcal{R}_{\text{Sup3680}}$ decaying into nOCH are, respectively, taken as $\sigma_{J/\psi}^{\text{D}}(s') = |A_{J/\psi}(s')|^2$ and $\sigma_{\mathcal{R}_{\text{Sup3680}}}^{\text{D}}(s') = |A_{\psi(3686)}(s') + \sum_1^3 A_k e^{i\phi_k}(s')|^2$, where $k = 1, 2, 3$ are for the $\mathcal{R}(3760)$ state, $\mathcal{R}(3780)$ state, and \mathcal{R} , respectively, and ϕ_k are relative phases.

In the above formulations, A_S are the generic decay amplitudes for these states, which are parameterized by $A_S(s') = \sqrt{12\pi\Gamma_S^{ee}\Gamma_S^{\text{tot}}\mathcal{B}(S \rightarrow \text{nOCH})}/[(s' - M_S^2) + i(\Gamma_S^{\text{tot}}M_S)]$, in which S stands for the J/ψ , $\psi(3686)$, $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states, as well as resonance \mathcal{R} ; M_S , Γ_S^{tot} , and Γ_S^{ee} are, respectively, the mass, total, and electronic widths of the S , and $\mathcal{B}(S \rightarrow \text{nOCH})$ is the decay branching fraction of the S . For the $\mathcal{R}(3780)$ state, the total width is set to be energy dependent, as in Ref. [24].

In the fit, the masses, the total widths, the electronic widths, and the hadronic decay branching fractions of J/ψ and $\psi(3686)$ resonances are fixed to those given in Ref. [36]. The $\mathcal{R}(3780)$ mass and total width are fixed to $(3781.0 \pm 1.3 \pm 0.5) \text{ MeV}/c^2$ and $(19.3 \pm 3.1 \pm 0.1) \text{ MeV}$ [26], respectively. The remaining parameters are left as free parameters in the fit.

The observed cross section is described by $\sigma_{\text{nOCH}}^{\text{O}}(s) = \int_0^{1-(4m_\pi^2/s)} dx \sigma_{\text{nOCH}}^{\text{D}}[s(1-x)]\mathcal{F}(x, s)$, where $\mathcal{F}(x, s)$ is the sampling function [24,37], and m_π is the pion mass. The cross section as a function of energy is determined by a fit to $\sigma_{\text{nOCH}}^{\text{O}}(s)$ at the 42 energy points. Using $\sigma_{\text{nOCH}}^{\text{D}}(s)$ obtained from the parameter values, we determine the ISR correction factors $[1 + \delta(s)] = \sigma_{\text{nOCH}}^{\text{O}}(s)/\sigma_{\text{nOCH}}^{\text{D}}(s)$ at the 42 energy points, which are shown in Fig. 2 (bottom). Dividing the observed nOCH cross section $\sigma_{\text{nOCH}}^{\text{O}}(s)$ by $[1 + \delta(s)]$ yields the observed dressed nOCH cross section $\sigma_{\text{nOCH}}^{\text{d}}(s)$, where the lower case superscripts are used to distinguish the measured from the predicted (upper case superscripts) quantities.

The circles with error bars in Fig. 3 show the $\sigma_{\text{nOCH}}^{\text{d}}(s)$ measurements, where the uncertainties are statistical. Using the same fit procedure described above, we fit $\sigma_{\text{nOCH}}^{\text{d}}(s)$ with the function $\sigma_{\text{nOCH}}^{\text{D}}(s)$ given in Eq. (1), with x fixed to zero. The fit converges at two sets of acceptable parameter values. We denote these as Result I and Result II. Table II summarizes the parameter values, where the first uncertainties are from the fit to the $\sigma_{\text{nOCH}}^{\text{d}}(s)$, and the second are systematic. The two sets of results have the fit quality of χ^2/ndof of 22.1/31, and 22.0/31, respectively. As the mass of the fitted resonance is close to 3810 MeV, we denote it as $\mathcal{R}(3810)$. The cross section described by Result I is superimposed on the fit results in Fig. 3. Also shown are the fit results including one contribution to $\sigma_{\mathcal{R}_{\text{Sup3680}}}^{\text{D}}(s)$ included at a time. The measured mass and total width of




FIG. 3. The dressed cross sections for $e^+e^- \rightarrow \text{nOCH}$, showing also the Result-I fit (line 1) and the contributions to this fit including separate contributions to $\sigma_{\mathcal{R}_{\text{Sup3680}}}^{\text{D}}(s)$ (lines 2–6).

the $\mathcal{R}(3760)$ state are consistent within 1.8σ and 0.9σ , respectively, with those measured by the BES-II experiment [26].

To estimate the systematic uncertainties of the fitted parameter in Table II, we vary the values of the $\sigma_{\text{nOCH}}^{\text{d}}(s)$, the values of the $\sigma_{D\bar{D}}^{\text{O}}(s)$, and the fixed parameters by $\pm 1\sigma$, refit the $\sigma_{\text{nOCH}}^{\text{d}}(s)$, and take the difference between the refitted parameter value and the baseline fit result as the corresponding systematic uncertainty. The estimation of the

TABLE II. Results of the fit to the cross sections for $e^+e^- \rightarrow \text{nOCH}$ showing the values of the mass M_k [MeV/c^2], total width Γ_k^{tot} [MeV], the product of electronic width (Γ_k^{ee}) and nOCH branching fraction (\mathcal{B}_k) $\Gamma_k^{ee}\mathcal{B}_k$ [eV], and relative phase ϕ_k [degree], where k represents $\mathcal{R}(3760)$, $\mathcal{R}(3780)$, and $\mathcal{R}(3810)$. \mathcal{B}_k is the branching fractions [%] for $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ decays into nOCH and OC.

Parameters	Result I	Result II
$M_{\mathcal{R}(3760)}$	$3739.9 \pm 4.2 \pm 2.6$	$3739.7 \pm 3.9 \pm 2.6$
$\Gamma_{\mathcal{R}(3760)}^{\text{tot}}$	$23.9 \pm 8.2 \pm 4.8$	$22.5 \pm 8.3 \pm 4.5$
$\Gamma_{\mathcal{R}(3760)}^{ee}\mathcal{B}_{\mathcal{R}(3760)}$	$46.8 \pm 29.9 \pm 25.1$	$11.9 \pm 9.0 \pm 6.4$
$\phi_{\mathcal{R}(3760)}$	$228 \pm 52 \pm 58$	$113 \pm 51 \pm 29$
$\Gamma_{\mathcal{R}(3780)}^{ee}\mathcal{B}_{\mathcal{R}(3780)}$	$29.9 \pm 16.1 \pm 3.7$	$25.3 \pm 11.6 \pm 3.1$
$\phi_{\mathcal{R}(3780)}$	$82 \pm 126 \pm 17$	$250 \pm 119 \pm 52$
$M_{\mathcal{R}(3810)}$	$3805.7 \pm 1.1 \pm 2.7$	$3805.7 \pm 1.1 \pm 2.7$
$\Gamma_{\mathcal{R}(3810)}^{\text{tot}}$	$11.6 \pm 2.9 \pm 1.9$	$11.5 \pm 2.8 \pm 1.9$
$\Gamma_{\mathcal{R}(3810)}^{ee}\mathcal{B}_{\mathcal{R}(3810)}$	$10.9 \pm 3.8 \pm 2.5$	$11.0 \pm 3.4 \pm 2.5$
$\phi_{\mathcal{R}(3810)}$	$52 \pm 149 \pm 25$	$215 \pm 148 \pm 103$
f	$2.28 \pm 0.01 \pm 0.07$	$2.28 \pm 0.01 \pm 0.07$
$\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{nOCH}]$	$25.2 \pm 16.1 \pm 30.4$	$6.4 \pm 4.8 \pm 7.7$
$\mathcal{B}[\mathcal{R}(3780) \rightarrow \text{nOCH}]$	$12.3 \pm 6.6 \pm 8.3$	$10.4 \pm 4.8 \pm 7.0$
$\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{OC}]$	$74.8 \pm 16.1 \pm 30.4$	$93.6 \pm 4.8 \pm 7.7$
$\mathcal{B}[\mathcal{R}(3780) \rightarrow \text{OC}]$	$87.7 \pm 6.6 \pm 8.3$	$89.6 \pm 4.8 \pm 7.0$

systematic uncertainty due to the uncertainty of the c.m. energies is similar to that described in Ref. [4]. Adding these uncertainties in quadrature yields the total systematic uncertainty.

Dividing the measured values for $\Gamma_{\mathcal{R}(3760)}^{ee} \mathcal{B}_{\mathcal{R}(3760)}$ and $\Gamma_{\mathcal{R}(3780)}^{ee} \mathcal{B}_{\mathcal{R}(3780)}$ in Table II by $\Gamma_{\mathcal{R}(3760)}^{ee} = (186 \pm 201 \pm 8) \text{ eV}$ [26] and $\Gamma_{\mathcal{R}(3780)}^{ee} = (243 \pm 160 \pm 9) \text{ eV}$ [26], respectively, yields the nOCH branching fractions for the decays $\mathcal{R}(3760) \rightarrow \text{nOCH}$ and $\mathcal{R}(3780) \rightarrow \text{nOCH}$, which are shown in Table II. Their corresponding OC branching fractions are also presented in Table II. These nOCH branching fractions of $\mathcal{R}(3780)$ decays are in good agreement with $\mathcal{B}[\psi(3770) \rightarrow \text{non-}D\bar{D}] = (15.1 \pm 5.6 \pm 1.8)\%$ [21–24] measured by the BES-II experiment.

By removing the $\mathcal{R}(3810)$ from the $\sigma_{\mathcal{R}_{\text{Sup}3680}}^D(s)$ as discussed above, the χ^2/ndof of the fit changes from 22.1/31 to 116.1/36, indicating that the significance of the $\mathcal{R}(3810)$ signal is 8.8σ . Similarly, the significance of the $\mathcal{R}(3760)$ signal is determined to be 8.1σ by comparing the difference of χ^2/ndof relative to the number degrees of freedom with and without including the $\mathcal{R}(3760)$ component in the fit. A similar procedure is applied to determine the $\mathcal{R}(3780)$ significance, which is 13.7σ . These significances include the systematic uncertainties.

The charmonium model [39] predicts only the 1^3D_1 state existing in the c.m. energy range from 3.733 to 3.870 GeV, which is generally assumed to be the $\psi(3770)$, and so the $\mathcal{R}(3760)$ and $\mathcal{R}(3810)$ states are presumably two non- $c\bar{c}$ states. The $\mathcal{R}(3760)$ state can be explained as a p -wave resonance of a four-quark ($c\bar{c}q\bar{q}$) state [40,41]. It can be thought of either as an OC molecular state, or as a four-quark bound state [41]. Reference [42] interprets $\mathcal{R}(3760)$ as a possible molecular OC threshold resonance. The most salient features of the $\mathcal{R}(3760)$ state is that its mass, which is $(3739.9 \pm 4.2 \pm 2.6) \text{ MeV}/c^2$ for Result I [$(3739.7 \pm 3.9 \pm 2.6) \text{ MeV}/c^2$ for Result II], is just at the D^+D^- threshold $(3739.3 \pm 0.1) \text{ MeV}/c^2$, and its branching fraction is $\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{OC}] = (74.8 \pm 16.1 \pm 30.4)\%$ [$\mathcal{B}\{\mathcal{R}(3760) \rightarrow \text{OC}\} = (93.6 \pm 4.8 \pm 7.7)\%$]. These experimental facts can lead one naturally to interpret the $\mathcal{R}(3760)$ state as an OC pair molecular state, but containing a simple four-quark state component. As no signal for the decay $\mathcal{R}(3810) \rightarrow D\bar{D}$ is observed in the cross sections for $e^+e^- \rightarrow D\bar{D}$ at c.m. energies around 3.810 GeV [43,44], and the $\mathcal{R}(3810)$ mass $(3805.7 \pm 1.1 \pm 2.7) \text{ MeV}/c^2$ for Result I [$(3805.7 \pm 1.1 \pm 2.7) \text{ MeV}/c^2$ for Result II] is exactly at the $h_c\pi^+\pi^-$ threshold $(3804.5 \pm 0.1) \text{ MeV}/c^2$, the $\mathcal{R}(3810)$ state can be interpreted as a hadrocharmonium resonance [8].

In summary, we have measured the inclusive nOCH cross sections of $e^+e^- \rightarrow \text{nOCH}$ with improved precision at c.m. energies from 3.645 to 3.871 GeV. We observe three

resonances: $\mathcal{R}(3760)$, $\mathcal{R}(3780)$, and $\mathcal{R}(3810)$ in the energy-dependent line shape of the nOCH cross sections with significances of 8.1σ , 13.7σ , and 8.8σ , respectively. The $\mathcal{R}(3760)$ and $\mathcal{R}(3780)$ states are observed for the first time in the nOCH cross sections, while the $\mathcal{R}(3810)$ state is observed for the first time with mass $(3805.7 \pm 1.1 \pm 2.7) [(3805.7 \pm 1.1 \pm 2.7)] \text{ MeV}/c^2$, total width $(11.6 \pm 2.9 \pm 1.9) [(11.5 \pm 2.8 \pm 1.9)] \text{ MeV}$, and the product of the electronic width and the nOCH decay branching fraction $(10.9 \pm 3.8 \pm 2.5) [(11.0 \pm 3.4 \pm 2.5)] \text{ eV}$ for Result I [Result II]. In addition, for the first time, we measure $\mathcal{B}[\mathcal{R}(3760) \rightarrow \text{nOCH}] = (25.2 \pm 16.1 \pm 30.4)\%$ [$(6.4 \pm 4.8 \pm 7.7)\%$] and $\mathcal{B}[\mathcal{R}(3780) \rightarrow \text{nOCH}] = (12.3 \pm 6.6 \pm 8.3)\%$ [$(10.4 \pm 4.8 \pm 7.0)\%$]. The $\mathcal{R}(3760)$ state can be interpreted as an OC pair molecular state, but containing a simple four-quark state component. The $\mathcal{R}(3810)$ state can be interpreted as a hadrocharmonium state [8].

The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts No. 2009CB825204, No. 2020YFA0406300, No. 2020YFA0406400; National Natural Science Foundation of China (NSFC) under Contracts No. 10935007, No. 11635010, No. 11735014, No. 11835012, No. 11935015, No. 11935016, No. 11935018, No. 11961141012, No. 12022510, No. 12025502, No. 12035009, No. 12035013, No. 12061131003, No. 12192260, No. 12192261, No. 12192262, No. 12192263, No. 12192264, No. 12192265, No. 12221005, No. 12225509, No. 12235017; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); CAS Key Research Program of Frontier Sciences under Contracts No. QYZDJ-SSW-SLH003, No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; The CAS Research Program under Code No. Y41G1010Y1; The CAS Other Research Program under Code No. Y129360; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; ERC under Contract No. 758462; European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement under Contract No. 894790; German Research Foundation DFG under Contracts No. 443159800, No. 455635585, Collaborative Research Center CRC 1044, FOR5327, GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia; National Science Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation of Thailand under

Contract No. B16F640076; Polish National Science Centre under Contract No. 2019/35/O/ST2/02907; The Swedish Research Council; U.S. Department of Energy under Contract No. DE-FG02-05ER41374.

- [1] J.Z. Bai *et al.* (BES Collaboration), [arXiv:hep-ex/0307028v1](#); W.G. Li, G. Rong, and D.G. Cassel, in *Proceedings of Tenth International Conference on Hadron Spectroscopy, Aschaffenburg, Germany, 2003*, pp. 495, 592, 937; J.Z. Bai *et al.* (BES Collaboration), *High Energy Physics and Nuclear Physics* **28**, 325 (2004).
- [2] G. Rong, in *Proceedings of the Symposium of 30 Years of BES Physics, Beijing, China, 2019* (World Scientific, Singapore, 2019), p. 48.
- [3] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **102**, 112009 (2020).
- [4] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **127**, 082002 (2021).
- [5] F.E. Close and P.R. Page, *Phys. Lett. B* **578**, 119 (2004).
- [6] M. B. Voloshin and L. B. Okun, *JETP Lett.* **23**, 369 (1976).
- [7] N. Brambilla *et al.*, *Eur. Phys. J. C* **71**, 1534 (2011).
- [8] S. Dubynskiy and M. B. Voloshin, *Phys. Lett. B* **666**, 344 (2008).
- [9] J.Z. Bai *et al.* (BES Collaboration), *Phys. Lett. B* **605**, 63 (2005).
- [10] S. K. Choi *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **91**, 262001 (2003).
- [11] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **95**, 142001 (2005).
- [12] Q. He *et al.* (CELO Collaboration), *Phys. Rev. D* **74**, 091104(R) (2006).
- [13] T. E. Coan *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **96**, 162003 (2006).
- [14] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **110**, 252001 (2013).
- [15] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **112**, 092001 (2014).
- [16] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **111**, 242001 (2013).
- [17] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **99**, 142002 (2007).
- [18] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **98**, 212001 (2007).
- [19] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **118**, 092001 (2017).
- [20] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **118**, 092002 (2017).
- [21] M. Ablikim *et al.* (BES Collaboration), *Phys. Lett. B* **641**, 145 (2006).
- [22] M. Ablikim *et al.* (BES Collaboration), *Phys. Rev. D* **76**, 122002 (2007).
- [23] M. Ablikim *et al.* (BES Collaboration), *Phys. Lett. B* **659**, 74 (2008).
- [24] M. Ablikim *et al.* (BES Collaboration), *Phys. Rev. Lett.* **97**, 121801 (2006).
- [25] G. Rong, *Chin. Phys. C* **34**, 788 (2010).
- [26] M. Ablikim *et al.* (BES Collaboration), *Phys. Rev. Lett.* **101**, 102004 (2008).
- [27] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
- [28] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [29] S. Jadach, B. F. L. Ward, and Z. Was, *Comput. Phys. Commun.* **130**, 260 (2000).
- [30] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001); R. G. Ping, *Chin. Phys. C* **32**, 599 (2008).
- [31] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang, and Y. S. Zhu, *Phys. Rev. D* **62**, 034003 (2000).
- [32] G. Balossini, C.M. Carloni Calame, G. Montagna, O. Nicosini, and F. Piccinini, *Nucl. Phys.* **B758**, 227 (2006); G. Balossini, C. Bignamini, C. M. C. Calame, G. Montagna, O. Nicosini, and F. Piccinini, *Phys. Lett. B* **663**, 209 (2008).
- [33] S. Nova, A. Olchevski, and T. Todorov, TWOGAM, a Monte Carlo event generator for two photon physics, DELPHI Note 90-35 PROG 152.
- [34] J.Z. Bai *et al.* (BES Collaboration), *Phys. Rev. D* **53**, 20 (1996).
- [35] H. Burkhardt and B. Pietrzyk, *Phys. Lett. B* **513**, 46 (2001).
- [36] R. L. Workman *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [37] E. A. Kuraev and V. S. Fadin, *Yad. Fiz.* **41**, 733 (1985); *Sov. J. Nucl. Phys.* **41**, 466 (1985).
- [38] D. Zhang, G. Rong, and J. C. Chen, *Phys. Rev. D* **74**, 054012 (2006).
- [39] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T.-M. Yan, *Phys. Rev. D* **17**, 3090 (1978); E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T. M. Yan, *Phys. Rev. D* **21**, 203 (1980).
- [40] A. De Rujula, Howard Georgi, and S. L. Glashow, *Phys. Rev. Lett.* **38**, 317 (1977).
- [41] Rafe Hyam Schindler, Stanford Linear Accelerator Center, Stanford University, Stanford, California, Report No. 219, p. 98, May 1979.
- [42] S. Dubynskiy and M. B. Voloshin, *Phys. Rev. D* **78**, 116014 (2008).
- [43] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. D* **76**, 111105(R) (2007).
- [44] G. Pakhlova *et al.* (Belle Collaboration), *Phys. Rev. D* **77**, 011103(R) (2008).

M. Ablikim,¹ M. N. Achasov,^{5,b} P. Adlarson,⁷⁵ X. C. Ai,⁸¹ R. Aliberti,³⁶ A. Amoroso,^{74a,74c} M. R. An,⁴⁰ Q. An,^{71,58} Y. Bai,⁵⁷ O. Bakina,³⁷ I. Balossino,^{30a} Y. Ban,^{47,g} V. Batozskaya,^{1,45} K. Begzsuren,³³ N. Berger,³⁶ M. Berlowski,⁴⁵ M. Bertani,^{29a} D. Bettoni,^{30a} F. Bianchi,^{74a,74c} E. Bianco,^{74a,74c} A. Bortone,^{74a,74c} I. Boyko,³⁷ R. A. Briere,⁶ A. Brueggemann,⁶⁸ H. Cai,⁷⁶

X. Cai,^{1,58} A. Calcaterra,^{29a} G. F. Cao,^{1,63} N. Cao,^{1,63} S. A. Cetin,^{62a} J. F. Chang,^{1,58} T. T. Chang,⁷⁷ W. L. Chang,^{1,63} G. R. Che,⁴⁴ G. Chelkov,^{37,a} C. Chen,⁴⁴ Chao Chen,⁵⁵ G. Chen,¹ H. S. Chen,^{1,63} M. L. Chen,^{1,58,63} S. J. Chen,⁴³ S. M. Chen,⁶¹ T. Chen,^{1,63} X. R. Chen,^{32,63} X. T. Chen,^{1,63} Y. B. Chen,^{1,58} Y. Q. Chen,³⁵ Z. J. Chen,^{26,h} W. S. Cheng,^{74c} S. K. Choi,¹¹ X. Chu,⁴⁴ G. Cibinetto,^{30a} S. C. Coen,⁴ F. Cossio,^{74c} J. J. Cui,⁵⁰ H. L. Dai,^{1,58} J. P. Dai,⁷⁹ A. Dbeyssi,¹⁹ R. E. de Boer,⁴ D. Dedovich,³⁷ Z. Y. Deng,¹ A. Denig,³⁶ I. Denysenko,³⁷ M. Destefanis,^{74a,74c} F. De Mori,^{74a,74c} B. Ding,^{66,1} X. X. Ding,^{47,g} Y. Ding,⁴¹ Y. Ding,³⁵ J. Dong,^{1,58} L. Y. Dong,^{1,63} M. Y. Dong,^{1,58,63} X. Dong,⁷⁶ M. C. Du,¹ S. X. Du,⁸¹ Z. H. Duan,⁴³ P. Egorov,^{37,a} Y. L. Fan,⁷⁶ J. Fang,^{1,58} S. S. Fang,^{1,63} W. X. Fang,¹ Y. Fang,¹ R. Farinelli,^{30a} L. Fava,^{74b,74c} F. Feldbauer,⁴ G. Felici,^{29a} C. Q. Feng,^{71,58} J. H. Feng,⁵⁹ K. Fischer,⁶⁹ M. Fritsch,⁴ C. Fritsch,⁶⁸ C. D. Fu,¹ J. L. Fu,⁶³ Y. W. Fu,¹ H. Gao,⁶³ Y. N. Gao,^{47,g} Yang Gao,^{71,58} S. Garbolino,^{74c} I. Garzia,^{30a,30b} P. T. Ge,⁷⁶ Z. W. Ge,⁴³ C. Geng,⁵⁹ E. M. Gersabeck,⁶⁷ A. Gilman,⁶⁹ K. Goetzen,¹⁴ L. Gong,⁴¹ W. X. Gong,^{1,58} W. Gradl,³⁶ S. Gramigna,^{30a,30b} M. Greco,^{74a,74c} M. H. Gu,^{1,58} Y. T. Gu,¹⁶ C. Y. Guan,^{1,63} Z. L. Guan,²³ A. Q. Guo,^{32,63} L. B. Guo,⁴² M. J. Guo,⁵⁰ R. P. Guo,⁴⁹ Y. P. Guo,^{13,f} A. Guskov,^{37,a} T. T. Han,⁵⁰ W. Y. Han,⁴⁰ X. Q. Hao,²⁰ F. A. Harris,⁶⁵ K. K. He,⁵⁵ K. L. He,^{1,63} F. H. H. Heinsius,⁴ C. H. Heinz,³⁶ Y. K. Heng,^{1,58,63} C. Herold,⁶⁰ T. Holtmann,⁴ P. C. Hong,^{13,f} G. Y. Hou,^{1,63} X. T. Hou,^{1,63} Y. R. Hou,⁶³ Z. L. Hou,¹ H. M. Hu,^{1,63} J. F. Hu,^{56,i} T. Hu,^{1,58,63} Y. Hu,¹ G. S. Huang,^{71,58} K. X. Huang,⁵⁹ L. Q. Huang,^{32,63} X. T. Huang,⁵⁰ Y. P. Huang,¹ T. Hussain,⁷³ N. Hüsken,^{28,36} W. Imoehl,²⁸ M. Irshad,^{71,58} J. Jackson,²⁸ S. Jaeger,⁴ S. Janchiv,³³ J. H. Jeong,¹¹ Q. Ji,¹ Q. P. Ji,²⁰ X. B. Ji,^{1,63} X. L. Ji,^{1,58} Y. Y. Ji,⁵⁰ X. Q. Jia,⁵⁰ Z. K. Jia,^{71,58} H. J. Jiang,⁷⁶ L. L. Jiang,¹ P. C. Jiang,^{47,g} S. S. Jiang,⁴⁰ T. J. Jiang,¹⁷ X. S. Jiang,^{1,58,63} Y. Jiang,⁶³ J. B. Jiao,⁵⁰ Z. Jiao,²⁴ S. Jin,⁴³ Y. Jin,⁶⁶ M. Q. Jing,^{1,63} T. Johansson,⁷⁵ X. Kui,¹ S. Kabana,³⁴ N. Kalantar-Nayestanaki,⁶⁴ X. L. Kang,¹⁰ X. S. Kang,⁴¹ R. Kappert,⁶⁴ M. Kavatsyuk,⁶⁴ B. C. Ke,⁸¹ A. Khoukaz,⁶⁸ R. Kiuchi,¹ R. Kliemt,¹⁴ O. B. Kolcu,^{62a} B. Kopf,⁴ M. K. Kuessner,⁴ A. Kupsc,^{45,75} W. Kühn,³⁸ J. J. Lane,⁶⁷ P. Larin,¹⁹ A. Lavania,²⁷ L. Lavezzi,^{74a,74c} T. T. Lei,^{71,k} Z. H. Lei,^{71,58} H. Leithoff,³⁶ M. Lellmann,³⁶ T. Lenz,³⁶ C. Li,⁴⁴ C. Li,⁴⁸ C. H. Li,⁴⁰ Cheng Li,^{71,58} D. M. Li,⁸¹ F. Li,^{1,58} G. Li,¹ H. Li,^{71,58} H. B. Li,^{1,63} H. J. Li,²⁰ H. N. Li,^{56,i} Hui Li,⁴⁴ J. R. Li,⁶¹ J. S. Li,⁵⁹ J. W. Li,⁵⁰ K. L. Li,²⁰ Ke Li,¹ L. J. Li,^{1,63} L. K. Li,¹ Lei Li,³ M. H. Li,⁴⁴ P. R. Li,^{39,j,k} Q. X. Li,⁵⁰ S. X. Li,¹³ T. Li,⁵⁰ W. D. Li,^{1,63} W. G. Li,¹ X. H. Li,^{71,58} X. L. Li,⁵⁰ Xiaoyu Li,^{1,63} Y. G. Li,^{47,g} Z. J. Li,⁵⁹ Z. X. Li,¹⁶ C. Liang,⁴³ H. Liang,³⁵ H. Liang,^{1,63} H. Liang,^{71,58} Y. F. Liang,⁵⁴ Y. T. Liang,^{32,63} G. R. Liao,¹⁵ L. Z. Liao,⁵⁰ Y. P. Liao,^{1,63} J. Libby,²⁷ A. Limphirat,⁶⁰ D. X. Lin,^{32,63} T. Lin,¹ B. J. Liu,¹ B. X. Liu,⁷⁶ C. Liu,³⁵ C. X. Liu,¹ F. H. Liu,⁵³ Fang Liu,¹ Feng Liu,⁷ G. M. Liu,^{56,i} H. Liu,^{39,j,k} H. B. Liu,¹⁶ H. M. Liu,^{1,63} Huanhuan Liu,¹ Huihui Liu,²² J. B. Liu,^{71,58} J. L. Liu,⁷² J. Y. Liu,^{1,63} K. Liu,¹ K. Y. Liu,⁴¹ Ke Liu,²³ L. Liu,^{71,58} L. C. Liu,⁴⁴ Lu Liu,⁴⁴ M. H. Liu,^{13,f} P. L. Liu,¹ Q. Liu,⁶³ S. B. Liu,^{71,58} T. Liu,^{13,f} W. K. Liu,⁴⁴ W. M. Liu,^{71,58} X. Liu,^{39,j,k} Y. Liu,^{39,j,k} Y. Liu,⁸¹ Y. B. Liu,⁴⁴ Z. A. Liu,^{1,58,63} Z. Q. Liu,⁵⁰ X. C. Lou,^{1,58,63} F. X. Lu,⁵⁹ H. J. Lu,²⁴ J. G. Lu,^{1,58} X. L. Lu,¹ Y. Lu,⁸ Y. P. Lu,^{1,58} Z. H. Lu,^{1,63} C. L. Luo,⁴² M. X. Luo,⁸⁰ T. Luo,^{13,f} X. L. Luo,^{1,58} X. R. Lyu,⁶³ Y. F. Lyu,⁴⁴ F. C. Ma,⁴¹ H. L. Ma,¹ J. L. Ma,^{1,63} L. L. Ma,⁵⁰ M. M. Ma,^{1,63} Q. M. Ma,¹ R. Q. Ma,^{1,63} R. T. Ma,⁶³ X. Y. Ma,^{1,58} Y. Ma,^{47,g} Y. M. Ma,³² F. E. Maas,¹⁹ M. Maggiora,^{74a,74c} S. Malde,⁶⁹ Q. A. Malik,⁷³ A. Mangoni,^{29b} Y. J. Mao,^{47,g} Z. P. Mao,¹ S. Marcello,^{74a,74c} Z. X. Meng,⁶⁶ J. G. Messchendorp,^{14,64} G. Mezzadri,^{30a} H. Miao,^{1,63} T. J. Min,⁴³ R. E. Mitchell,²⁸ X. H. Mo,^{1,58,63} N. Yu. Muchnoi,^{5,b} Y. Nefedov,³⁷ F. Nerling,^{19,d} I. B. Nikolaev,^{5,b} Z. Ning,^{1,58} S. Nisar,^{12,1} Y. Niu,⁵⁰ S. L. Olsen,⁶³ Q. Ouyang,^{1,58,63} S. Pacetti,^{29b,29c} X. Pan,⁵⁵ Y. Pan,⁵⁷ A. Pathak,³⁵ P. Patteri,^{29a} Y. P. Pei,^{71,58} M. Pelizaeus,⁴ H. P. Peng,^{71,58} K. Peters,^{14,d} J. L. Ping,⁴² R. G. Ping,^{1,63} S. Plura,³⁶ S. Pogodin,³⁷ V. Prasad,³⁴ F. Z. Qi,¹ H. Qi,^{71,58} H. R. Qi,⁶¹ M. Qi,⁴³ T. Y. Qi,^{13,f} S. Qian,^{1,58} W. B. Qian,⁶³ C. F. Qiao,⁶³ J. J. Qin,⁷² L. Q. Qin,¹⁵ X. P. Qin,^{13,f} X. S. Qin,⁵⁰ Z. H. Qin,^{1,58} J. F. Qiu,¹ S. Q. Qu,⁶¹ C. F. Redmer,³⁶ K. J. Ren,⁴⁰ A. Rivetti,^{74c} V. Rodin,⁶⁴ M. Rolo,^{74c} G. Rong,^{1,63} Ch. Rosner,¹⁹ S. N. Ruan,⁴⁴ N. Salone,⁴⁵ A. Sarantsev,^{37,c} Y. Schelhaas,³⁶ K. Schoenning,⁷⁵ M. Scodreggio,^{30a,30b} K. Y. Shan,^{13,f} W. Shan,²⁵ X. Y. Shan,^{71,58} J. F. Shangguan,⁵⁵ L. G. Shao,^{1,63} M. Shao,^{71,58} C. P. Shen,^{13,f} H. F. Shen,^{1,63} W. H. Shen,⁶³ X. Y. Shen,^{1,63} B. A. Shi,⁶³ H. C. Shi,^{71,58} J. L. Shi,¹³ J. Y. Shi,¹ Q. Q. Shi,⁵⁵ R. S. Shi,^{1,63} X. Shi,^{1,58} J. J. Song,²⁰ T. Z. Song,⁵⁹ W. M. Song,^{35,1} Y. J. Song,¹³ Y. X. Song,^{47,g} S. Sosio,^{74a,74c} S. Spataro,^{74a,74c} F. Stieler,³⁶ Y. J. Su,⁶³ G. B. Sun,⁷⁶ G. X. Sun,¹ H. Sun,⁶³ H. K. Sun,¹ J. F. Sun,²⁰ K. Sun,⁶¹ L. Sun,⁷⁶ S. S. Sun,^{1,63} T. Sun,^{1,63} W. Y. Sun,³⁵ Y. Sun,¹⁰ Y. J. Sun,^{71,58} Y. Z. Sun,¹ Z. T. Sun,⁵⁰ Y. X. Tan,^{71,58} C. J. Tang,⁵⁴ G. Y. Tang,¹ J. Tang,⁵⁹ Y. A. Tang,⁷⁶ L. Y. Tao,⁷² Q. T. Tao,^{26,h} M. Tat,⁶⁹ J. X. Teng,^{71,58} V. Thoren,⁷⁵ W. H. Tian,⁵⁹ W. H. Tian,⁵² Y. Tian,^{32,63} Z. F. Tian,⁷⁶ I. Uman,^{62b} S. J. Wang,⁵⁰ B. Wang,¹ B. L. Wang,⁶³ Bo Wang,^{71,58} C. W. Wang,⁴³ D. Y. Wang,^{47,g} F. Wang,⁷² H. J. Wang,^{39,j,k} H. P. Wang,^{1,63} J. P. Wang,⁵⁰ K. Wang,^{1,58} L. L. Wang,¹ M. Wang,⁵⁰ Meng Wang,^{1,63} S. Wang,^{39,j,k} S. Wang,^{13,f} T. Wang,^{13,f} T. J. Wang,⁴⁴ W. Wang,⁷² W. Wang,⁵⁹ W. P. Wang,^{71,58} X. Wang,^{47,g} X. F. Wang,^{39,j,k} X. J. Wang,⁴⁰ X. L. Wang,^{13,f} Y. Wang,⁶¹ Y. D. Wang,⁴⁶ Y. F. Wang,^{1,58,63} Y. H. Wang,⁴⁸ Y. N. Wang,⁴⁶ Y. Q. Wang,¹ Yaqian Wang,^{18,1} Yi Wang,⁶¹ Z. Wang,^{1,58} Z. L. Wang,⁷² Z. Y. Wang,^{1,63} Ziyi Wang,⁶³ D. Wei,⁷⁰ D. H. Wei,¹⁵ F. Weidner,⁶⁸

S. P. Wen,¹ C. W. Wenzel,⁴ U. W. Wiedner,⁴ G. Wilkinson,⁶⁹ M. Wolke,⁷⁵ L. Wollenberg,⁴ C. Wu,⁴⁰ J. F. Wu,^{1,63} L. H. Wu,¹ L. J. Wu,^{1,63} X. Wu,^{13,f} X. H. Wu,³⁵ Y. Wu,⁷¹ Y. J. Wu,³² Z. Wu,^{1,58} L. Xia,^{71,58} X. M. Xian,⁴⁰ T. Xiang,^{47,g} D. Xiao,^{39,j,k} G. Y. Xiao,⁴³ H. Xiao,^{13,f} S. Y. Xiao,¹ Y. L. Xiao,^{13,f} Z. J. Xiao,⁴² C. Xie,⁴³ X. H. Xie,^{47,g} Y. Xie,⁵⁰ Y. G. Xie,^{1,58} Y. H. Xie,⁷ Z. P. Xie,^{71,58} T. Y. Xing,^{1,63} C. F. Xu,^{1,63} C. J. Xu,⁵⁹ G. F. Xu,¹ H. Y. Xu,⁶⁶ Q. J. Xu,¹⁷ Q. N. Xu,³¹ W. Xu,^{1,63} W. L. Xu,⁶⁶ X. P. Xu,⁵⁵ Y. C. Xu,⁷⁸ Z. P. Xu,⁴³ Z. S. Xu,⁶³ F. Yan,^{13,f} L. Yan,^{13,f} W. B. Yan,^{71,58} W. C. Yan,⁸¹ X. Q. Yan,¹ H. J. Yang,^{51,e} H. L. Yang,³⁵ H. X. Yang,¹ Tao Yang,¹ Y. Yang,^{13,f} Y. F. Yang,⁴⁴ Y. X. Yang,^{1,63} Yifan Yang,^{1,63} Z. W. Yang,^{39,j,k} Z. P. Yao,⁵⁰ M. Ye,^{1,58} M. H. Ye,⁹ J. H. Yin,¹ Z. Y. You,⁵⁹ B. X. Yu,^{1,58,63} C. X. Yu,⁴⁴ G. Yu,^{1,63} J. S. Yu,^{26,h} T. Yu,⁷² X. D. Yu,^{47,g} C. Z. Yuan,^{1,63} L. Yuan,² S. C. Yuan,¹ X. Q. Yuan,¹ Y. Yuan,^{1,63} Z. Y. Yuan,⁵⁹ C. X. Yue,⁴⁰ A. A. Zafar,⁷³ F. R. Zeng,⁵⁰ X. Zeng,^{13,f} Y. Zeng,^{26,h} Y. J. Zeng,^{1,63} X. Y. Zhai,³⁵ Y. C. Zhai,⁵⁰ Y. H. Zhan,⁵⁹ A. Q. Zhang,^{1,63} B. L. Zhang,^{1,63} B. X. Zhang,¹ D. H. Zhang,⁴⁴ G. Y. Zhang,²⁰ H. Zhang,⁷¹ H. H. Zhang,⁵⁹ H. H. Zhang,³⁵ H. Q. Zhang,^{1,58,63} H. Y. Zhang,^{1,58} J. J. Zhang,⁵² J. L. Zhang,²¹ J. Q. Zhang,⁴² J. W. Zhang,^{1,58,63} J. X. Zhang,^{39,j,k} J. Y. Zhang,¹ J. Z. Zhang,^{1,63} Jianyu Zhang,⁶³ Jiawei Zhang,^{1,63} L. M. Zhang,⁶¹ L. Q. Zhang,⁵⁹ Lei Zhang,⁴³ P. Zhang,¹ Q. Y. Zhang,^{40,81} Shuihan Zhang,^{1,63} Shulei Zhang,^{26,h} X. D. Zhang,⁴⁶ X. M. Zhang,¹ X. Y. Zhang,⁵⁰ Xuyan Zhang,⁵⁵ Y. Zhang,⁷² Y. Zhang,⁶⁹ Y. T. Zhang,⁸¹ Y. H. Zhang,^{1,58} Yan Zhang,^{71,58} Yao Zhang,¹ Z. H. Zhang,¹ Z. L. Zhang,³⁵ Z. Y. Zhang,⁴⁴ Z. Y. Zhang,⁷⁶ G. Zhao,¹ J. Zhao,⁴⁰ J. Y. Zhao,^{1,63} J. Z. Zhao,^{1,58} Lei Zhao,^{71,58} Ling Zhao,¹ M. G. Zhao,⁴⁴ S. J. Zhao,⁸¹ Y. B. Zhao,^{1,58} Y. X. Zhao,^{32,63} Z. G. Zhao,^{71,58} A. Zhemchugov,^{37,a} B. Zheng,⁷² J. P. Zheng,^{1,58} W. J. Zheng,^{1,63} Y. H. Zheng,⁶³ B. Zhong,⁴² X. Zhong,⁵⁹ H. Zhou,⁵⁰ L. P. Zhou,^{1,63} X. Zhou,⁷⁶ X. K. Zhou,⁷ X. R. Zhou,^{71,58} X. Y. Zhou,⁴⁰ Y. Z. Zhou,^{13,f} J. Zhu,⁴⁴ K. Zhu,¹ K. J. Zhu,^{1,58,63} L. Zhu,³⁵ L. X. Zhu,⁶³ S. H. Zhu,⁷⁰ S. Q. Zhu,⁴³ T. J. Zhu,^{13,f} W. J. Zhu,^{13,f} Y. C. Zhu,^{71,58} Z. A. Zhu,^{1,63} J. H. Zou,¹ and J. Zu^{71,58}

(BESIII Collaboration)

¹*Institute of High Energy Physics, Beijing 100049, People's Republic of China*

²*Beihang University, Beijing 100191, People's Republic of China*

³*Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China*

⁴*Bochum Ruhr-University, D-44780 Bochum, Germany*

⁵*Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*

⁶*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

⁷*Central China Normal University, Wuhan 430079, People's Republic of China*

⁸*Central South University, Changsha 410083, People's Republic of China*

⁹*China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*

¹⁰*China University of Geosciences, Wuhan 430074, People's Republic of China*

¹¹*Chung-Ang University, Seoul, 06974, Republic of Korea*

¹²*COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*

¹³*Fudan University, Shanghai 200433, People's Republic of China*

¹⁴*GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*

¹⁵*Guangxi Normal University, Guilin 541004, People's Republic of China*

¹⁶*Guangxi University, Nanning 530004, People's Republic of China*

¹⁷*Hangzhou Normal University, Hangzhou 310036, People's Republic of China*

¹⁸*Hebei University, Baoding 071002, People's Republic of China*

¹⁹*Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*

²⁰*Henan Normal University, Xinxiang 453007, People's Republic of China*

²¹*Henan University, Kaifeng 475004, People's Republic of China*

²²*Henan University of Science and Technology, Luoyang 471003, People's Republic of China*

²³*Henan University of Technology, Zhengzhou 450001, People's Republic of China*

²⁴*Huangshan College, Huangshan 245000, People's Republic of China*

²⁵*Hunan Normal University, Changsha 410081, People's Republic of China*

²⁶*Hunan University, Changsha 410082, People's Republic of China*

²⁷*Indian Institute of Technology Madras, Chennai 600036, India*

²⁸*Indiana University, Bloomington, Indiana 47405, USA*

^{29a}*INFN Laboratori Nazionali di Frascati, INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*

^{29b}*INFN Sezione di Perugia, I-06100 Perugia, Italy*

^{29c}*University of Perugia, I-06100 Perugia, Italy*

^{30a}*INFN Sezione di Ferrara, INFN Sezione di Ferrara, I-44122 Ferrara, Italy*

- ^{30b}University of Ferrara, I-44122 Ferrara, Italy
- ³¹Inner Mongolia University, Hohhot 010021, People's Republic of China
- ³²Institute of Modern Physics, Lanzhou 730000, People's Republic of China
- ³³Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia
- ³⁴Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile
- ³⁵Jilin University, Changchun 130012, People's Republic of China
- ³⁶Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany
- ³⁷Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia
- ³⁸Justus-Liebig-Universität Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany
- ³⁹Lanzhou University, Lanzhou 730000, People's Republic of China
- ⁴⁰Liaoning Normal University, Dalian 116029, People's Republic of China
- ⁴¹Liaoning University, Shenyang 110036, People's Republic of China
- ⁴²Nanjing Normal University, Nanjing 210023, People's Republic of China
- ⁴³Nanjing University, Nanjing 210093, People's Republic of China
- ⁴⁴Nankai University, Tianjin 300071, People's Republic of China
- ⁴⁵National Centre for Nuclear Research, Warsaw 02-093, Poland
- ⁴⁶North China Electric Power University, Beijing 102206, People's Republic of China
- ⁴⁷Peking University, Beijing 100871, People's Republic of China
- ⁴⁸Qufu Normal University, Qufu 273165, People's Republic of China
- ⁴⁹Shandong Normal University, Jinan 250014, People's Republic of China
- ⁵⁰Shandong University, Jinan 250100, People's Republic of China
- ⁵¹Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China
- ⁵²Shanxi Normal University, Linfen 041004, People's Republic of China
- ⁵³Shanxi University, Taiyuan 030006, People's Republic of China
- ⁵⁴Sichuan University, Chengdu 610064, People's Republic of China
- ⁵⁵Soochow University, Suzhou 215006, People's Republic of China
- ⁵⁶South China Normal University, Guangzhou 510006, People's Republic of China
- ⁵⁷Southeast University, Nanjing 211100, People's Republic of China
- ⁵⁸State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China
- ⁵⁹Sun Yat-Sen University, Guangzhou 510275, People's Republic of China
- ⁶⁰Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand
- ⁶¹Tsinghua University, Beijing 100084, People's Republic of China
- ^{62a}Turkish Accelerator Center Particle Factory Group, Istinye University, 34010, Istanbul, Turkey
- ^{62b}Near East University, Nicosia, North Cyprus, 99138, Mersin 10, Turkey
- ⁶³University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China
- ⁶⁴University of Groningen, NL-9747 AA Groningen, The Netherlands
- ⁶⁵University of Hawaii, Honolulu, Hawaii 96822, USA
- ⁶⁶University of Jinan, Jinan 250022, People's Republic of China
- ⁶⁷University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom
- ⁶⁸University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany
- ⁶⁹University of Oxford, Keble Road, Oxford OX13RH, United Kingdom
- ⁷⁰University of Science and Technology Liaoning, Anshan 114051, People's Republic of China
- ⁷¹University of Science and Technology of China, Hefei 230026, People's Republic of China
- ⁷²University of South China, Hengyang 421001, People's Republic of China
- ⁷³University of the Punjab, Lahore-54590, Pakistan
- ^{74a}University of Turin and INFN, University of Turin, I-10125 Turin, Italy
- ^{74b}University of Eastern Piedmont, I-15121 Alessandria, Italy
- ^{74c}INFN, I-10125 Turin, Italy
- ⁷⁵Uppsala University, Box 516, SE-75120 Uppsala, Sweden
- ⁷⁶Wuhan University, Wuhan 430072, People's Republic of China
- ⁷⁷Xinyang Normal University, Xinyang 464000, People's Republic of China
- ⁷⁸Yantai University, Yantai 264005, People's Republic of China
- ⁷⁹Yunnan University, Kunming 650500, People's Republic of China
- ⁸⁰Zhejiang University, Hangzhou 310027, People's Republic of China
- ⁸¹Zhengzhou University, Zhengzhou 450001, People's Republic of China

^aAlso at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

^bAlso at the Novosibirsk State University, Novosibirsk, 630090, Russia.

^cAlso at the NRC "Kurchatov Institute," PNPI, 188300, Gatchina, Russia.

^dAlso at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

^eAlso at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China.

^fAlso at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China.

^gAlso at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China.

^hAlso at School of Physics and Electronics, Hunan University, Changsha 410082, China.

ⁱAlso at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China.

^jAlso at Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People's Republic of China.

^kAlso at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China.

^lAlso at the Department of Mathematical Sciences, IBA, Karachi 75270, Pakistan.