



Double-heavy axial-vector tetraquark $T_{bc;\bar{u}\bar{d}}^0$

S.S. Agaev^a, K. Azizi^{b,c,*}, H. Sundu^d

^a Institute for Physical Problems, Baku State University, Az-1148 Baku, Azerbaijan

^b Department of Physics, University of Tehran, North Karegar Avenue, Tehran 14395-547, Iran

^c Department of Physics, Doğuş University, Acibadem-Kadıköy, 34722 Istanbul, Turkey

^d Department of Physics, Kocaeli University, 41380 Izmit, Turkey

Received 31 October 2019; received in revised form 30 November 2019; accepted 7 December 2019

Available online 12 December 2019

Editor: Hong-Jian He

Abstract

The mass and coupling of the axial-vector tetraquark $T_{bc;\bar{u}\bar{d}}^0$ (in a short form T_{bc}^0) are calculated by means of the QCD two-point sum rule method. In computations we take into account contributions arising from various quark, gluon and mixed vacuum condensates up to dimension 10. The central value of the mass $m = (7105 \pm 155)$ MeV lies below the thresholds for the strong and electromagnetic decays of the T_{bc}^0 state, and hence it transforms to conventional mesons only through the weak decays. In the case of $m = 7260$ MeV the tetraquark T_{bc}^0 becomes the strong- and electromagnetic-interaction unstable particle. In the first case, we find the full width and mean lifetime of T_{bc}^0 using its semileptonic decays $T_{bc}^0 \rightarrow T_{cc;\bar{u}\bar{d}}^+ l \bar{\nu}_l$ ($l = e, \mu, \tau$), where the final-state tetraquark is a scalar state. We compute also partial widths of the nonleptonic weak decays $T_{bc}^0 \rightarrow T_{cc;\bar{u}\bar{d}}^+ \pi^-$ (K^-, D^-, D_s^-), and take into account their effects on the full width of T_{bc}^0 . In the context of the second scenario we calculate partial widths of S -wave strong decays $T_{bc}^0 \rightarrow B^{*-} D^+$ and $T_{bc}^0 \rightarrow \bar{B}^{*0} D^0$, and using these channels evaluate the full width of T_{bc}^0 . Predictions for $\Gamma_{\text{full}} = (3.98 \pm 0.51) \times 10^{-10}$ MeV and mean lifetime $\tau = 1.65_{-0.18}^{+0.25}$ ps of T_{bc}^0 obtained in the context of the first option, as well as the full width $\Gamma_{\text{full}} = (63.5 \pm 8.9)$ MeV extracted in the second scenario may be useful for experimental and theoretical exploration of double-heavy exotic mesons.

© 2019 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author at: Department of Physics, University of Tehran, North Karegar Avenue, Tehran, Iran.
E-mail address: kazem.azizi@ut.ac.ir (K. Azizi).

1. Introduction

During last two decades double-heavy tetraquarks as real candidates to stable four-quark states became objects of intensive studies. In the pioneering papers [1–3] it was demonstrated that a heavy Q and light q quarks may form the stable exotic mesons $QQ\bar{q}\bar{q}$ provided the ratio m_Q/m_q is large enough. These results were obtained in the context of a potential model with the additive pairwise interaction, but even models with relaxed restrictions on the confining potential led to the similar conclusions. Indeed, in accordance with Ref. [4] the isoscalar axial-vector tetraquark $T_{bb;\bar{u}\bar{d}}^-$ (or T_{bb}^-) turns to be strong-interaction stable state that lies below the $B\bar{B}^*$ threshold. It is worth noting that an only constraint imposed in Ref. [4] on the potential was its finiteness at close distances of two particles. Therefore, T_{bb}^- decays to conventional mesons only through weak processes and has a long lifetime, which is important for its experimental exploration. A situation with the tetraquarks $T_{bc;\bar{q}\bar{q}'}$ and $T_{cc;\bar{q}\bar{q}'}$ was not clear, because bc and cc diquarks might constitute both stable and unstable states.

In years followed after this progress, various models of high energy physics were used to investigate the double-heavy tetraquarks T_{QQ} [5–12]. An interest to these problems was renewed by results of the LHCb Collaboration which measured parameters of the doubly charmed baryon $\Xi_{cc}^{++} = ccu$ [13]. These parameters were used in Ref. [14] to evaluate the mass and analyze possible decay channels of T_{bb}^- . Predictions obtained there confirmed the stability of T_{bb}^- against the strong and electromagnetic decays to $B^-\bar{B}^{*0}$ and $B^-\bar{B}^0\gamma$, respectively. The strong-interaction stable nature of the tetraquarks T_{bb}^- , $T_{bb;\bar{u}\bar{s}}^-$, and $T_{bb;\bar{d}\bar{s}}^0$ was demonstrated in Ref. [15] by invoking heavy-quark symmetry relations. The mass and coupling of T_{bb}^- was evaluated in our work [16] as well, in which we estimated also its full width and mean lifetime using the semileptonic decay channel $T_{bb}^- \rightarrow Z_{bc;\bar{u}\bar{d}}^0 l\bar{\nu}_l$.

Another class of four-quark mesons, namely one that contains the heavy diquarks bc is on agenda of physicists as well. The scalar and axial-vector tetraquarks $bc\bar{u}\bar{d}$ are particles of special interest, because they may form strong-interaction stable compounds. But calculations performed in the context of different approaches lead to controversial results. Thus, the Bethe-Salpeter method predicts the mass of the scalar tetraquark $Z_{bc;\bar{u}\bar{d}}^0$ (in what follows Z_{bc}^0) at around 6.93 GeV, which is below the threshold 7145 MeV for S -wave strong decays to heavy mesons B^-D^+ and \bar{B}^0D^0 [17]. Recent analysis demonstrated that Z_{bc}^0 lies 11 MeV below this threshold [14], whereas the authors of Ref. [15] found the masses of the scalar and axial-vector tetraquarks $bc\bar{u}\bar{d}$ equal to 7229 MeV and 7272 MeV, respectively. These predictions make kinematically allowed their strong decays to ordinary B^-D^+/\bar{B}^0D^0 and B^*D mesons.

It is interesting that lattice calculations prove the strong-interaction stable nature of the axial-vector tetraquark $ud\bar{b}\bar{c}$, because its mass is below the $\bar{D}B^*$ threshold [18]. However, the authors could not decide would this exotic meson decay weakly or might transform also to the final state $\bar{D}B\gamma$. The stability of $J^P = 0^+$ and 1^+ isoscalar tetraquarks $bc\bar{u}\bar{d}$ was confirmed in Ref. [19], in which it was found that $J^P = 0^+$ state is a strong- and electromagnetic-interaction stable, whereas $J^P = 1^+$ may also transform through the electromagnetic interaction.

In the context of the QCD sum rule approach the spectroscopic parameters of the scalar tetraquark Z_{bc}^0 were calculated also in our work [16]. For the mass of Z_{bc}^0 our computations predicted $m_Z = (6660 \pm 150)$ MeV, which is considerably below the threshold 7145 MeV. The electromagnetic decay modes $Z_{bc}^0 \rightarrow \bar{B}^0D_1^0\gamma$ and $B^*D_0^*\gamma$ are also among forbidden processes, because relevant thresholds exceed 7600 MeV and are higher than the mass of Z_{bc}^0 . In other

words, in accordance with our results the scalar tetraquark Z_{bc}^0 is a strong- and electromagnetic-interaction stable, and transforms due to weak decays, which were used to find its full width and mean lifetime [20].

In the present article we study the axial-vector tetraquark $T_{bc;\bar{u}\bar{d}}^0$ (hereafter T_{bc}^0) by computing its spectroscopic parameters, full width and mean lifetime. The mass m and coupling f of T_{bc}^0 are evaluated in the framework of the QCD two-point sum rule method by taking into account vacuum expectation values of the local quark, gluon and mixed operators up to dimension ten. The mass of T_{bc}^0 extracted in the present work $m = (7105 \pm 155)$ MeV contains theoretical errors typical for sum rule computations, hence, there are two options to find its full width and estimate mean lifetime. Thus, the central value of the mass is lower than the thresholds 7190 MeV and 7286 MeV for strong S -wave decays of T_{bc}^0 to final states $B^{*-}D^+/\bar{B}^{*0}D^0$ and $B^-D^{*+}/\bar{B}^0D^{*0}$, respectively. This mass is also lower than the threshold 7145 MeV for the electromagnetic decays $D^+B^-\gamma/D^0\bar{B}^0\gamma$. Therefore, in this case the full width and lifetime of the exotic meson T_{bc}^0 should be determined from its weak decays. But considering the maximum theoretical prediction for $m = 7260$ MeV, one sees that it is higher than the threshold for strong decays $B^{*-}D^+/\bar{B}^{*0}D^0$ and electromagnetic transitions $D^+B^-\gamma/D^0\bar{B}^0\gamma$. Realization of this scenario means that the width of the tetraquark T_{bc}^0 is determined mainly by strong decays, because partial widths of weak and electromagnetic processes are very small and can be neglected.

To calculate the full width of the tetraquark T_{bc}^0 , we consider both scenarios. In the first case $m = 7105$ MeV we evaluate partial widths of the processes $T_{bc}^0 \rightarrow T_{cc;\bar{u}\bar{d}}^+ l\bar{\nu}_l$ ($l = e, \mu$ and τ) by treating the final-state tetraquark $T_{cc;\bar{u}\bar{d}}^+$ (in what follows T_{cc}^+) as a scalar particle. These decays run due to transition $b \rightarrow W^-c$. The differential rates of these semileptonic decays are determined by the weak form factors $G_i(q^2)$ ($i = 0, 1, 2, 3$), which are evaluated by employing the QCD three-point sum rule approach. Then, partial width of the processes $T_{bc}^0 \rightarrow T_{cc}^+ l\bar{\nu}_l$ can be found by integrating the relevant differential rates over the momentum transfer q^2 . The sum rule method does not encompass all kinematically allowed values of q^2 , therefore we introduce fit functions that coincide with sum rule predictions, and can be extrapolated to cover a whole integration region.

But a decay $b \rightarrow W^-c$ can be followed by transitions $W^- \rightarrow d\bar{u}, s\bar{u}, d\bar{c}$ and $s\bar{c}$ as well. Afterwards these quark pairs can form ordinary mesons through different mechanisms. Thus, in the hard-scattering picture a pair $d\bar{u}$, for example, can create conventional mesons with $q\bar{q}$ quarks appeared due to a gluon from one of d or \bar{u} quarks. These processes generate final states $T_{bc}^0 \rightarrow T_{cc}^+ M_1(d\bar{q})M_2(q\bar{u})$ which are suppressed relative to the semileptonic decays by the factor $\alpha_s^2 |V_{ud}|^2$. Alternatively, pairs of quarks $d\bar{u}, s\bar{u}, d\bar{c}$ and $s\bar{c}$ can form π^-, K^-, D^- and D_s^- mesons triggering the two-body nonleptonic decays $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-(K^-, D^-, D_s^-)$. Another class of the T_{bc}^0 tetraquark's weak decays is connected with possibility of direct combination of these quarks with ones from $T_{cc;\bar{u}\bar{d}}^+$ and creation of three-meson final states. The two-body and three-meson nonleptonic decays do not suppressed by additional factors relative to the semileptonic decays, and their contributions to full width of T_{bc}^0 may be considerable.

In the second scenario $m = 7260$ MeV, and this mass is above the threshold for strong decays to mesons $B^{*-}D^+/\bar{B}^{*0}D^0$, but is still below the threshold for other two possible decay modes to final states $B^-D^{*+}/\bar{B}^0D^{*0}$. Therefore, we calculate the partial width of the kinematically allowed strong S -wave decays $T_{bc}^0 \rightarrow B^{*-}D^+$ and $T_{bc}^0 \rightarrow \bar{B}^{*0}D^0$. To this end, we use again the QCD three-point sum rule method and evaluate the strong form factors $g_1(q^2)$ and $g_2(q^2)$. By

extrapolating these form factors to the corresponding mass shells we determine couplings of the vertices $T_{bc}^0 B^{*-} D^+$ and $T_{bc}^0 \bar{B}^{*0} D^0$, and calculate partial width of these decays. The full width of the tetraquark T_{bc}^0 is evaluated using these two dominant strong decay channels.

This article is organized in the following manner: In Section 2, from analysis of the two-point correlation function with an appropriate interpolating current, we derive sum rules to evaluate the spectroscopic parameters of the tetraquark T_{bc}^0 . In the next Section 3, using the parameters of T_{bc}^0 and ones of the final-state tetraquark, we calculate the partial width of its semileptonic decays $T_{bc}^0 \rightarrow T_{cc;\bar{u}\bar{d}}^+ l \bar{\nu}_l$ ($l = e, \mu, \tau$). To this end, we derive the sum rules for the weak form factors and by means of fit functions extrapolate them to the whole region, where an integration over q^2 should be carried out. In Section 4, we analyze the nonleptonic weak decays $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-$ (K^-, D^-, D_s^-) of the tetraquark T_{bc}^0 and find their partial widths. Here, we also calculate the full width of T_{bc}^0 in the first scenario, i.e., for $m = 7105$ MeV. The Sec. 5 is devoted to calculation of the partial widths of the strong processes $T_{bc}^0 \rightarrow B^{*-} D^+$ and $T_{bc}^0 \rightarrow \bar{B}^{*0} D^0$, where we also evaluate the full width of the tetraquark T_{bc}^0 if $m = 7260$ MeV. Section 6 is reserved for analysis of obtained results, and contains also our concluding notes.

2. Mass and coupling of the axial-vector tetraquark T_{bc}^0

In this section we extract the spectroscopic parameters of the axial-vector tetraquark T_{bc}^0 from the QCD sum rules. To this end, we start from analysis of the correlation function

$$\Pi_{\mu\nu}(p) = i \int d^4x e^{ipx} \langle 0 | \mathcal{T} \{ J_\mu(x) J_\nu^\dagger(0) \} | 0 \rangle, \quad (1)$$

where $J_\mu(x)$ is the interpolating current to the axial-vector tetraquark T_{bc}^0 . We suggest that T_{bc}^0 is built of the scalar diquark and axial-vector antiquark, and hence its current has the form

$$J_\mu(x) = b_a^T(x) C \gamma_5 c_b(x) \left[\bar{u}_a(x) \gamma_\mu C \bar{d}_b^T(x) - \bar{u}_b(x) \gamma_\mu C \bar{d}_a^T(x) \right]. \quad (2)$$

Here a and b are the color indices and C is the charge conjugation operator. The current (2) has the antisymmetric color structure $[\bar{3}_c]_{bc} \otimes [3_c]_{\bar{u}\bar{d}}$ and describes a four-quark state with the quantum numbers 1^+ , where $b^T C \gamma_5 c$ and $\bar{u} \gamma_\mu C \bar{d}^T$ are the scalar diquark and axial-vector antiquark, respectively.

To derive required sum rules, in accordance with prescriptions of the method we express the correlation function $\Pi_{\mu\nu}(p)$ in terms of the tetraquark's mass m and coupling f . We consider T_{bc}^0 as a ground-state particle, and isolate the first term in $\Pi_{\mu\nu}^{\text{Phys}}(p)$

$$\Pi_{\mu\nu}^{\text{Phys}}(p) = \frac{\langle 0 | J_\mu | T_{bc}^0(p) \rangle \langle T_{bc}^0(p) | J_\nu^\dagger | 0 \rangle}{m^2 - p^2} + \dots \quad (3)$$

Equation (3) is obtained by saturating the correlation function with a complete set of $J^P = 1^+$ states and carrying out the integration over x . Contributions of higher resonances and continuum states to $\Pi_{\mu\nu}^{\text{Phys}}(p)$ are denoted by the dots.

To simplify further the correlator $\Pi_{\mu\nu}^{\text{Phys}}(p)$ it is useful to define the matrix element

$$\langle 0 | J_\mu | T_{bc}^0(p, \epsilon) \rangle = f m \epsilon_\mu, \quad (4)$$

with ϵ_μ being the polarization vector of the T_{bc}^0 state. Then in terms of m and f the correlation function $\Pi_{\mu\nu}^{\text{Phys}}(p)$ takes the form

$$\Pi_{\mu\nu}^{\text{Phys}}(p) = \frac{m^2 f^2}{m^2 - p^2} \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{m^2} \right) + \dots \quad (5)$$

The QCD side of the sum rule is determined by the correlation function $\Pi_{\mu\nu}(p)$, but calculated now by employing the quark propagators

$$\begin{aligned} \Pi_{\mu\nu}^{\text{OPE}}(p) = & i \int d^4x e^{ipx} \text{Tr} \left[\gamma_5 \tilde{S}_b^{aa'}(x) \gamma_5 S_c^{bb'}(x) \right] \\ & \times \left\{ \text{Tr} \left[\gamma_\mu \tilde{S}_d^{a'b}(-x) \gamma_\nu S_u^{b'a}(-x) \right] - \text{Tr} \left[\gamma_\mu \tilde{S}_d^{b'b}(-x) \gamma_\nu S_u^{a'a}(-x) \right] \right. \\ & \left. - \text{Tr} \left[\gamma_\mu \tilde{S}_d^{a'a}(-x) \gamma_\nu S_u^{b'b}(-x) \right] + \text{Tr} \left[\gamma_\mu \tilde{S}_d^{b'a}(-x) \gamma_\nu S_u^{a'b}(-x) \right] \right\}, \quad (6) \end{aligned}$$

where $S_q^{ab}(x)$ is the heavy (b, c)- or light (u, d)-quark propagators. Their explicit expressions can be found in Ref. [21]. In Eq. (6) we use the shorthand notation

$$\tilde{S}_q(x) = C S_q^T(x) C. \quad (7)$$

The correlation function $\Pi_{\mu\nu}(p)$ contains the different Lorentz structures one of which should be chosen to get the sum rules. The invariant amplitudes $\Pi^{\text{Phys}}(p^2)$ and $\Pi^{\text{OPE}}(p^2)$ corresponding to the terms $\sim g_{\mu\nu}$ are convenient for our aim, because they do not receive contributions from the scalar particles.

After picking up and equating corresponding invariant amplitudes, we apply the Borel transformation to both sides of the obtained expression. This is necessary to suppress contributions of the higher resonances and continuum states. Afterwards, one has to subtract continuum contributions, which is achieved by invoking suggestion on the quark-hadron duality. The obtained equality acquires a dependence on auxiliary parameters of the sum rules M^2 and s_0 : first of them is the Borel parameter appeared due to corresponding transformation, the second one s_0 is the continuum threshold parameter that separates the ground-state and higher resonances from each another.

The final sum rule for the mass of the state T_{bc}^0 reads:

$$m^2 = \frac{\int_{\mathcal{M}^2}^{s_0} ds s \rho^{\text{OPE}}(s) e^{-s/M^2}}{\int_{\mathcal{M}^2}^{s_0} ds \rho^{\text{OPE}}(s) e^{-s/M^2}}, \quad (8)$$

where $\mathcal{M} = m_b + m_c$. For the coupling f one obtains the expression

$$f^2 = \frac{1}{m^2} \int_{\mathcal{M}^2}^{s_0} ds \rho^{\text{OPE}}(s) e^{(m^2-s)/M^2}. \quad (9)$$

Here $\rho^{\text{OPE}}(s)$ is the two-point spectral density, which is determined as an imaginary part of the term in $\Pi_{\mu\nu}^{\text{OPE}}(p)$ proportional to $g_{\mu\nu}$, and calculated by taking into account the quark, gluon and mixed vacuum condensates up to dimension ten. Explicit expression of $\rho^{\text{OPE}}(s)$ is rather cumbersome, hence we refrain from providing it here.

In addition to M^2 and s_0 , numerical values of which depend on the considering problem, the sum rules (8) and (9) contain also the vacuum condensates, as well as the masses of b and c -quarks




Fig. 1. The pole contribution as a function of the Borel and continuum threshold parameters M^2 and s_0 .

$$\begin{aligned}
 \langle \bar{q}q \rangle &= -(0.24 \pm 0.01)^3 \text{ GeV}^3, \quad \langle \bar{q}g_s\sigma Gq \rangle = m_0^2 \langle \bar{q}q \rangle, \quad m_0^2 = (0.8 \pm 0.1) \text{ GeV}^2, \\
 \langle \frac{\alpha_s G^2}{\pi} \rangle &= (0.012 \pm 0.004) \text{ GeV}^4, \quad \langle g_s^3 G^3 \rangle = (0.57 \pm 0.29) \text{ GeV}^6, \\
 m_b &= 4.18_{-0.03}^{+0.04} \text{ GeV}, \quad m_c = 1.275_{-0.035}^{+0.025} \text{ GeV}.
 \end{aligned} \tag{10}$$

The parameters M^2 and s_0 should satisfy constraints that are standard for the sum rule computations. Thus, at maximum of the Borel parameter the pole contribution (PC) should be larger than some fixed value, whereas the main criterium to fix the minimum of a Borel window is convergence of the operator product expansion (OPE). Additionally, at minimum M^2 the perturbative contribution has to exceed the nonperturbative terms considerably. Because quantities extracted from the sum rules demonstrate dependence on the auxiliary parameters, the regions for M^2 and s_0 should minimize these side effects, as well.

Our analysis proves that the working regions

$$M^2 \in [5.5, 7] \text{ GeV}^2, \quad s_0 \in [61, 63] \text{ GeV}^2, \tag{11}$$

satisfy all aforementioned restrictions. Thus, within the region $M^2 \in [5.5, 7] \text{ GeV}^2$ the pole contribution decreases approximately from 0.58 till 0.34. A detailed picture for PC is presented in Fig. 1, where we plot the pole contribution as a function of M^2 and s_0 . The minimum M_{\min}^2 is found from analysis of the ratio

$$R(M^2) = \frac{\Pi^{\text{DimN}}(M^2, s_0)}{\Pi(M^2, s_0)}, \tag{12}$$

where $\Pi(M^2, s_0)$ is the Borel transformed and subtracted function $\Pi^{\text{OPE}}(p^2)$. In the present work as a measure of the convergence we use the sum of last three terms in OPE $\text{DimN} = \text{Dim}(8 + 9 + 10)$ and impose the constraint $R(M_{\min}^2) \leq 0.01$ which is fulfilled at 5.5 GeV^2 . The perturbative contribution at $M^2 = 5.5 \text{ GeV}^2$ amounts to 0.68 part of the full result and overshoots contribution of the nonperturbative terms. In Fig. 2 we demonstrate the dependence of the mass m on M^2 and s_0 , where weak residual effects of these parameters are seen.

Our results for m and f read:

$$m = (7105 \pm 155) \text{ MeV}, \quad f = (1.0 \pm 0.2) \times 10^{-2} \text{ GeV}^4. \tag{13}$$

Theoretical errors of the mass is milder than ones of the coupling, nevertheless all these ambiguities do not exceed standard limits of sum rule computations reaching $\pm 2.2\%$ and $\pm 20\%$ of




Fig. 2. The same as in Fig. 1, but for the mass of the tetraquark T_{bc}^0 .

the corresponding central values, respectively. The spectroscopic parameters of the axial-vector tetraquark T_{bc}^0 evaluated in this section from a basis for our further investigations.

3. Semileptonic decays $T_{bc}^0 \rightarrow T_{cc}^+ l \bar{\nu}_l$

As it has been emphasized above for $m = 7105$ MeV the tetraquark T_{bc}^0 is stable against the strong and electromagnetic interactions, because then m resides 85/190 MeV and 45 MeV below the strong and electromagnetic thresholds, respectively. In other words, T_{bc}^0 can dissociate to conventional mesons only due to weak transformations. One of such transitions is weak decay $b \rightarrow W^- c \rightarrow c l \bar{\nu}$ of the heavy b -quark, that triggers the semileptonic decays $T_{bc}^0 \rightarrow T_{cc}^+ l \bar{\nu}_l$ of the tetraquark T_{bc}^0 . It is not difficult to see, that due to large mass difference between the tetraquarks T_{bc}^0 and T_{cc}^+ , all of the decays $T_{bc}^0 \rightarrow T_{cc}^+ l \bar{\nu}_l$ with $l = e, \mu$ and τ are kinematically allowed processes. We restrict ourselves by considering only the dominant process $b \rightarrow W^- c$, because due to smallness of the Cabibbo-Kobayashi-Maskawa (CKM) matrix element $|V_{bu}|^2/|V_{bc}|^2 \simeq 0.01$ the decay $b \rightarrow W^- u$ is suppressed relative to the first one.

At the tree-level, the transition $b \rightarrow W^- c$ is described by means of the effective Hamiltonian

$$\mathcal{H}^{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{bc} \bar{c} \gamma_\mu (1 - \gamma_5) b \bar{l} \gamma^\mu (1 - \gamma_5) \nu_l. \quad (14)$$

Here G_F is the Fermi coupling constant, and V_{bc} is the element of the CKM matrix. After substituting \mathcal{H}^{eff} between the initial and final tetraquark fields and factoring out the leptonic piece we get the matrix element of the current

$$J_\mu^{\text{tr}} = \bar{c} \gamma_\mu (1 - \gamma_5) b, \quad (15)$$

which has to be calculated in terms of the weak form factors $G_i(q^2)$: they parameterize the long-distance dynamics of the transition

$$\begin{aligned} \langle T_{cc}^+(p') | J_\mu^{\text{tr}} | T_{bc}^0(p, \epsilon) \rangle = & \bar{m} G_0(q^2) \epsilon_\mu + \frac{G_1(q^2)}{\bar{m}} (\epsilon p') P_\mu + \frac{G_2(q^2)}{\bar{m}} (\epsilon p') q_\mu \\ & + i \frac{G_3(q^2)}{\bar{m}} \epsilon_{\mu\nu\alpha\beta} \epsilon^\nu p^\alpha p'^\beta. \end{aligned} \quad (16)$$

In Eq. (16) p and ϵ are the momentum and polarization vector of the T_{bc}^0 , p' is the momentum of the scalar tetraquark T_{cc}^+ . Here, we also use the shorthand notations $\bar{m} = m + m_T$ and $P_\mu = p'_\mu +$

p_μ with m_T being the mass of the final-state tetraquark. The $q_\mu = p_\mu - p'_\mu$ is the momentum transferred to the leptons changing within the limits $m_l^2 \leq q^2 \leq (m - m_T)^2$, where m_l is the mass of the lepton l .

The form factors $G_i(q^2)$ are key quantities to be extracted from the sum rules. To this end, we consider the following three-point correlation function:

$$\Pi_{\mu\nu}(p, p') = i^2 \int d^4x d^4y e^{i(p'y - px)} \langle 0 | \mathcal{T} \{ J^T(y) J_\mu^{\text{tr}}(0) J_\nu^\dagger(x) \} | 0 \rangle, \quad (17)$$

where $J_\nu(x)$ and $J^T(y)$ are the interpolating currents corresponding to the states T_{bc}^0 and T_{cc}^+ , respectively. The current $J_\nu(x)$ has been introduced by Eq. (2). The interpolating current for the state T_{cc}^+ is given by the expression:

$$J^T(y) = \epsilon \tilde{\epsilon} [c_b^T(y) C \gamma_\alpha c_c(y)] [\bar{u}_d(y) \gamma^\alpha C \bar{d}_e^T(y)], \quad (18)$$

where $\epsilon \tilde{\epsilon} = \epsilon^{abc} \epsilon^{ade}$. Here, $\epsilon^{abc} [c_b^T C \gamma_\alpha c_c]$ and $\epsilon^{ade} [\bar{u}_d \gamma^\alpha C \bar{d}_e^T]$ are the axial-vector diquark and antidiquark, respectively. Then the scalar designation of the final tetraquark T_{cc}^+ stems naturally from the internal structure of the initial four-quark state T_{bc}^0 , which is the axial-vector particle composed of the scalar diquark $b^T C \gamma_5 c$ and axial-vector antidiquark $\bar{u} \gamma_\mu C \bar{d}^T$. The semileptonic decay $T_{bc}^0 \rightarrow T_{cc}^+ + W^-$ runs through $b \rightarrow W^- c$, which transforms the scalar diquark bc to the final axial-vector cc , leaving, at the same time, unchanged the initial light antidiquark; the light axial-vector antidiquark $\bar{u}\bar{d}$ appears both in the initial and final states. The designation of T_{cc}^+ as an axial-vector requires $\bar{u}\bar{d}$ to be a scalar, which implies additional spin-rearrangement in the initial axial-vector $\bar{u}\bar{d}$ diquark, which evidently suppresses the corresponding process.

Our strategy to derive sum rules for the form factors $G_i(q^2)$ is the same as in all of this kind studies. In fact, to determine the phenomenological side of the sum rule $\Pi_{\mu\nu}^{\text{Phys}}(p, p')$ we express the correlation function $\Pi_{\mu\nu}(p, p')$ in terms of the spectroscopic parameters of particles involving into the decay process. Afterwards we find the QCD side (or OPE) side of the sum rules $\Pi_{\mu\nu}^{\text{OPE}}(p, p')$ by computing the same correlation function in terms of quark propagators. By matching the obtained results and utilizing the quark-hadron duality assumption we extract sum rules and evaluate the physical quantities of interest. Because the quark propagators contain quark, gluon and mixed vacuum condensates, the sum rules express the physical quantities as functions of nonperturbative parameters.

In the context of this approach the function $\Pi_{\mu\nu}^{\text{Phys}}(p, p')$ can be recast into the form

$$\Pi_{\mu\nu}^{\text{Phys}}(p, p') = \frac{\langle 0 | J^T | T_{cc}^+(p') \rangle \langle T_{cc}^+(p') | J_\mu^{\text{tr}} | T_{bc}^0(p, \epsilon) \rangle \langle T_{bc}^0(p, \epsilon) | J_\nu^\dagger | 0 \rangle}{(p^2 - m^2)(p'^2 - m_T^2)} + \dots, \quad (19)$$

where m_T is the mass of T_{cc}^+ . In the expression above we take into account contribution appearing due to only the ground-state particles, denoting contributions of the higher resonances and continuum states by the dots.

Transformation of the ground-state term in $\Pi_{\mu\nu}^{\text{Phys}}(p, p')$ can be completed by detailing the matrix elements in its expression. The matrix element of T_{bc}^0 and $\langle T_{cc}^+(p') | J_\mu^{\text{tr}} | T_{bc}^0(p, \epsilon) \rangle$ are given by Eqs. (4) and (16), respectively. The remaining matrix element $\langle 0 | J^T | T_{cc}^+(p') \rangle$ has a simple form

$$\langle 0 | J^T | T_{cc}^+(p') \rangle = m_T f_T, \quad (20)$$

and depends only on the mass and coupling f_T of the tetraquark T_{cc}^+ . Benefiting from these explicit formulas, for $\Pi_{\mu\nu}^{\text{Phys}}(p, p', q^2)$ we obtain

$$\begin{aligned} \Pi_{\mu\nu}^{\text{Phys}}(p, p', q^2) = & \frac{f m f_T m_T}{(p^2 - m^2)(p'^2 - m_T^2)} \left\{ \bar{m} G_0(q^2) \left(-g_{\mu\nu} + \frac{p_\mu p_\nu}{m^2} \right) + \left[\frac{G_1(q^2)}{\bar{m}} P_\mu \right. \right. \\ & \left. \left. + \frac{G_2(q^2)}{\bar{m}} q_\mu \right] \left(-p'_\nu + \frac{m^2 + m_T^2 - q^2}{2m^2} p_\nu \right) - i \frac{G_3(q^2)}{\bar{m}} \varepsilon_{\mu\nu\alpha\beta} p^\alpha p'^\beta \right\} + \dots \end{aligned} \quad (21)$$

The function $\Pi_{\mu\nu}^{\text{OPE}}(p, p')$ forms the second side of the sum rules:

$$\begin{aligned} \Pi_{\mu\nu}^{\text{OPE}}(p, p') = & i^2 \int d^4 x d^4 y e^{i(p'y - px)} \bar{\epsilon} \tilde{\epsilon} \left(\text{Tr} \left[\gamma^\alpha \tilde{S}_d^{b'e}(x-y) \gamma_\nu S_u^{a'd}(x-y) \right] \right. \\ & \left. - \text{Tr} \left[\gamma^\alpha \tilde{S}_d^{a'e}(x-y) \gamma_\nu S_u^{b'd}(x-y) \right] \right) \left(\text{Tr} \left[\gamma_\mu (1 - \gamma_5) S_b^{ia'}(-x) \gamma_5 \tilde{S}_c^{bb'}(y-x) \gamma_\alpha S_c^{ci}(y) \right] \right. \\ & \left. - \text{Tr} \left[\gamma_\mu (1 - \gamma_5) S_b^{ia'}(-x) \gamma_5 \tilde{S}_c^{cb'}(y-x) \gamma_\alpha S_c^{bi}(y) \right] \right). \end{aligned} \quad (22)$$

The sum rules for the form factors $G_i(q^2)$ can be obtained by equating invariant amplitudes corresponding to the same Lorentz structures in $\Pi_{\mu\nu}^{\text{Phys}}(p, p', q^2)$ and $\Pi_{\mu\nu}^{\text{OPE}}(p, p')$. Because in the three-point sum rules the invariant amplitudes are functions of p'^2 and p^2 , to suppress contributions of higher resonances and continuum states we have to apply the double Borel transformation over these variables. As a result, the final expressions depend on a set of Borel parameters $\mathbf{M}^2 = (M_1^2, M_2^2)$. The continuum subtraction is performed in two channels using two continuum threshold parameters $\mathbf{s}_0 = (s_0, s'_0)$. The form factor $G_0(q^2)$ is obtained by using the structure $g_{\mu\nu}$ and reads:

$$G_0(\mathbf{M}^2, \mathbf{s}_0, q^2) = \frac{1}{\bar{m} f m f_T m_T} \int_{\mathcal{M}^2}^{s_0} ds \int_{4m^2}^{s'_0} ds' \rho_0(s, s', q^2) e^{(m^2-s)/M_1^2} e^{(m_T^2-s')/M_2^2}. \quad (23)$$

The form factors $G_i(q^2)$ ($i = 1, 2, 3$) are derived employing other Lorentz structures in the correlation functions:

$$G_i(\mathbf{M}^2, \mathbf{s}_0, q^2) = \frac{\bar{m}}{f m f_T m_T} \int_{\mathcal{M}^2}^{s_0} ds \int_{4m^2}^{s'_0} ds' \rho_i(s, s', q^2) e^{(m^2-s)/M_1^2} e^{(m_T^2-s')/M_2^2}. \quad (24)$$

The sum rules (23) and (24) are written down in terms of the spectral densities $\rho_i(s, s', q^2)$ which are proportional to the imaginary parts of the corresponding terms in $\Pi_{\mu\nu}^{\text{OPE}}(p, p')$. They contain the perturbative and nonperturbative contributions, and are calculated with dimension-5 accuracy.

To compute the weak form factors $G_i(\mathbf{M}^2, \mathbf{s}_0, q^2)$ we need numerical values of parameters which enter to the sum rules. The vacuum condensates are given in Eq. (10), whereas the spectroscopic parameters of the tetraquark T_{cc}^+ is borrowed from our work [22]. The mass and coupling of the initial particle T_{bc}^0 have been calculated in the previous section; these and other parameters are collected in Table 1. In computations, we impose on the auxiliary parameters \mathbf{M}^2 and \mathbf{s}_0 the same constraints as in the mass calculations: the set (M_1^2, s_0) for the initial particle channel is determined by Eq. (11), whereas the set (M_2^2, s'_0) for T_{cc}^+ is chosen in the form [22]

Table 1

The mass and coupling of the final-state tetraquark T_{cc}^+ and other parameters used in numerical computations.

Quantity	Value
m_T	$(3845 \pm 175) \text{ MeV}$
f_T	$(1.16 \pm 0.26) \times 10^{-2} \text{ GeV}^4$
m_e	0.511 MeV
m_μ	105.658 MeV
m_τ	$(1776.82 \pm 0.16) \text{ MeV}$
G_F	$1.16637 \times 10^{-5} \text{ GeV}^{-2}$
$ V_{bc} $	$(42.2 \pm 0.08) \times 10^{-3}$




Fig. 3. The form factor $|G_0| = |G_0(5 \text{ GeV}^2)|$ as a function of the Borel parameters M_1^2 and M_2^2 at $s_0 = 62 \text{ GeV}^2$ and $s'_0 = 20 \text{ GeV}^2$.

$$M_2^2 \in [3, 4] \text{ GeV}^2, \quad s'_0 \in [19, 21] \text{ GeV}^2. \quad (25)$$

Results of sum rule calculations in the case of $G_0(q^2)$, as an example, are shown in Fig. 3. The similar predictions have been obtained for the remaining form factors as well. The sum rule results for the functions $G_i(q^2)$ are necessary, but not enough to calculate the partial width of the process $T_{bc}^0 \rightarrow T_{cc}^+ l \bar{\nu}_l$. The reason is that these form factors determine its differential decay rate $d\Gamma/dq^2$ (see, Appendix in Ref. [16]). The partial width Γ should be found by integrating $d\Gamma/dq^2$ over q^2 within limits allowed by the kinematical constraints $m_l^2 \leq q^2 \leq (m - m_T)^2$. But sum rules do not cover all this region $m_l^2 \leq q^2 \leq 10.63 \text{ GeV}^2$, and give reliable results within the limits $m_l^2 \leq q^2 \leq 8 \text{ GeV}^2$. Therefore, one has to introduce the model functions $G_i(q^2)$, which at q^2 accessible for the sum rule computations coincide with $G_i(q^2)$, but can be extrapolated to the whole integration region.

The fit functions

$$G_i(q^2) = G_0^i \exp \left[c_1^i \frac{q^2}{m_{\text{fit}}^2} + c_2^i \left(\frac{q^2}{m_{\text{fit}}^2} \right)^2 \right], \quad (26)$$

are convenient for these purposes. Here G_0^i , c_1^i , c_2^i and m_{fit}^2 are the fitted parameters; m_{fit}^2 is equal to 50.48 GeV^2 , numerical values of others are collected in Table 2. Our predictions for the partial width of the semileptonic decay channels are:

Table 2
The parameters of the functions $G_i(q^2)$.

Functions	G_0^i	c_1^i	c_2^i
$G_0(q^2)$	-0.92	0.43	-9.36
$G_1(q^2)$	10.87	2.83	3.69
$G_2(q^2)$	-2.61	0.32	4.44
$G_3(q^2)$	-13.79	2.06	3.31

$$\begin{aligned}
 \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ e^- \bar{\nu}_e) &= (1.44 \pm 0.35) \times 10^{-10} \text{ MeV}, \\
 \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ \mu^- \bar{\nu}_\mu) &= (1.43 \pm 0.34) \times 10^{-10} \text{ MeV}, \\
 \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ \tau^- \bar{\nu}_\tau) &= (4.3 \pm 1.1) \times 10^{-11} \text{ MeV}.
 \end{aligned} \tag{27}$$

Results (27) obtained in this section constitute an important part of the full width of T_{bc}^0 , and will be used below for its evaluation.

4. Two-body weak decays $T_{bc}^0 \rightarrow T_{cc}^+ \pi^- (K^-, D^-, D_s^-)$

The two-body weak decays $T_{bc}^0 \rightarrow T_{cc}^+ \pi^- (K^-, D^-, D_s^-)$ of the tetraquark T_{bc}^0 can be considered in the context of the QCD factorization approach, which allows one to write amplitudes and calculate widths of these processes. This method was successfully applied to study two-body weak decays of the conventional mesons [23,24], and is used here to investigate two-body decays of the tetraquark T_{bc}^0 , when one of the final particles is an exotic meson.

We consider in a detailed form only the decay $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-$, and write down final predictions for remaining channels. At the quark level, the effective Hamiltonian for this decay is given by the expression

$$\tilde{\mathcal{H}}^{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{bc} V_{ud}^* [c_1(\mu) Q_1 + c_2(\mu) Q_2], \tag{28}$$

where

$$Q_1 = (\bar{d}_i u_i)_{V-A} (\bar{c}_j b_j)_{V-A}, \quad Q_2 = (\bar{d}_i u_j)_{V-A} (\bar{c}_j b_i)_{V-A}, \tag{29}$$

and i, j are the color indices. Here $c_1(\mu)$ and $c_2(\mu)$ are the short-distance Wilson coefficients evaluated at the scale μ at which the factorization is assumed to be correct. The shorthand notation $(\bar{q}_1 q_2)_{V-A}$ in Eq. (29) means

$$(\bar{q}_1 q_2)_{V-A} = \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2. \tag{30}$$

The amplitude of this decay can be written down in the following factorized form

$$\mathcal{A} = \frac{G_F}{\sqrt{2}} V_{bc} V_{ud}^* a_1(\mu) \langle \pi^-(q) | (\bar{d}_i u_i)_{V-A} | 0 \rangle \langle T_{cc}^+(p') | (\bar{c}_j b_j)_{V-A} | T_{bc}^0(p, \epsilon) \rangle \tag{31}$$

where

$$a_1(\mu) = c_1(\mu) + \frac{1}{N_c} c_2(\mu), \tag{32}$$

and N_c is the number of quark colors. The amplitude \mathcal{A} describes the process in which the pion π^- is generated directly from the color-singlet current $(\bar{d}_i u_i)_{V-A}$. The matrix element

$\langle T_{cc}^+(p') | (\bar{c}_j b_j)_{V-A} | T_{bc}^0(p, \epsilon) \rangle$ has been introduced by Eq. (16), whereas the matrix element of the pion is given by the expression

$$\langle \pi^-(q) | (\bar{d}_i u_i)_{V-A} | 0 \rangle = i f_\pi q_\mu, \quad (33)$$

and is determined by its decay constant f_π .

Then, it is not difficult to see that \mathcal{A} takes the form

$$\mathcal{A} = i \frac{G_F}{\sqrt{2}} f_\pi V_{bc} V_{ud}^* a_1(\mu) (\epsilon p') \left[-\bar{m} G_0(q^2) + \frac{G_1(q^2)}{\bar{m}} P q + \frac{G_2(q^2)}{\bar{m}} m_\pi^2 \right] \quad (34)$$

The width of the decay $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-$ is

$$\begin{aligned} \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ \pi^-) &= \frac{G_F^2 f_\pi^2}{48\pi m^2} |V_{bc}|^2 |V_{ud}|^2 a_1^2(\mu) \lambda^3(m^2, m_T^2, m_\pi^2) \\ &\times \left[\bar{m}^2 |G_0|^2 + \frac{|G_1|^2}{\bar{m}^2} (m^2 - m_T^2)^2 + \frac{|G_2|^2}{\bar{m}^2} m_\pi^4 - 2 \operatorname{Re}[G_0 G_1^*] (m^2 - m_T^2) \right. \\ &\left. - 2 \operatorname{Re}[G_0 G_2^*] m_\pi^2 + 2 \operatorname{Re}[G_1 G_2^*] \frac{(m^2 - m_T^2) m_\pi^2}{\bar{m}^2} \right], \quad (35) \end{aligned}$$

where the weak form factors $G_i(q^2)$ ($i = 0, 1, 2$) are taken at $q^2 = m_\pi^2$. In Eq. (35) the function $\lambda(m^2, m_T^2, m_\pi^2)$ is given by the formula

$$\lambda(m^2, m_T^2, m_\pi^2) = \frac{1}{2m} \left[m^4 + m_T^4 + m_\pi^4 - 2(m^2 m_T^2 + m^2 m_\pi^2 + m_T^2 m_\pi^2) \right]^{1/2}. \quad (36)$$

The similar analysis can be performed for other decays $T_{bc}^0 \rightarrow T_{cc}^+ K^- (D^-, D_s^-)$ as well: relevant expressions can be obtained from (35) using the spectroscopic parameters of the mesons K^- , D^- , and D_s^- , and by replacements $V_{ud} \rightarrow V_{us}$, V_{cd} , and V_{cs} , respectively.

Numerical computations can be carried out after fixing the spectroscopic parameters of the final-state pseudoscalar mesons, weak form factors, and CKM matrix elements. The masses and decay constants of the final-state pseudoscalar mesons are presented in Table 3. The weak form factors $G_i(q^2)$ ($i = 0, 1, 2$), which are crucial parts of calculations, have been obtained in the previous section. For CKM matrix elements we use $|V_{ud}| = 0.97420 \pm 0.00021$, $|V_{us}| = 0.2243 \pm 0.0005$, $|V_{cd}| = 0.218 \pm 0.004$ and $|V_{cs}| = 0.997 \pm 0.017$. The values of the Wilson coefficients $c_1(m_b)$, and $c_2(m_b)$ with next-to-leading order QCD corrections were presented in Refs. [25–27]

$$c_1(m_b) = 1.117, \quad c_2(m_b) = -0.257. \quad (37)$$

For the decay $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-$, calculations lead to the following result

$$\Gamma(T_{bc}^0 \rightarrow T_{cc}^+ \pi^-) = (1.73 \pm 0.38) \times 10^{-11} \text{ MeV}. \quad (38)$$

Width of this decay is smaller than widths of the semileptonic decays, but is comparable with them. For the remaining weak nonleptonic decays of the tetraquark T_{bc}^0 we get

$$\begin{aligned} \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ K^-) &= (1.27 \pm 0.26) \times 10^{-12} \text{ MeV}, \\ \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ D^-) &= (1.65 \pm 0.35) \times 10^{-12} \text{ MeV}, \\ \Gamma(T_{bc}^0 \rightarrow T_{cc}^+ D_s^-) &= (4.74 \pm 0.99) \times 10^{-11} \text{ MeV}. \quad (39) \end{aligned}$$

Table 3
Masses and decay constants of the pseudoscalar mesons.

Quantity	Value
m_π	139.570 MeV
m_K	(493.677 ± 0.016) MeV
m_D	(1869.61 ± 0.10) MeV
m_{D_s}	(1968.30 ± 0.11) MeV
f_π	131 MeV
f_K	(155.72 ± 0.51) MeV
f_D	(203.7 ± 4.7) MeV
f_{D_s}	(257.8 ± 4.1) MeV

It is seen that partial widths only of the nonleptonic weak decays $T_{bc}^0 \rightarrow T_{cc}^+ D_s^-$ and $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-$ are comparable with widths of the semileptonic modes (27); contribution to the full width of T_{bc}^0 coming from other two weak decays is negligible.

Using Eqs. (27), (38) and (39), it is not difficult to find the full width and mean lifetime of T_{bc}^0

$$\Gamma_{\text{full}} = (3.98 \pm 0.51) \times 10^{-10} \text{ MeV}, \quad \tau = 1.65_{-0.18}^{+0.25} \times 10^{-12} \text{ s}. \quad (40)$$

Predictions for Γ_{full} and τ are among main results of the present work.

5. Strong decays $T_{bc}^0 \rightarrow B^{*-} D^+$ and $T_{bc}^0 \rightarrow \bar{B}^{*0} D^0$

Calculations of the mass of the tetraquark T_{bc}^0 , performed in Section 2, due to uncertainties of the sum rule method do not exclude also prediction $m = 7260$ MeV. In this scenario T_{bc}^0 is strong-interaction unstable particle and decays to conventional mesons $B^{*-} D^+$ and $\bar{B}^{*0} D^0$. It is worth noting that $m = 7260$ MeV is below the thresholds for strong decays $T_{bc}^0 \rightarrow B^- D^{*+}$ and $T_{bc}^0 \rightarrow \bar{B}^0 D^{*0}$, which forbids kinematically these processes. Below we present in a detailed form our analysis of the decay $T_{bc}^0 \rightarrow B^{*-} D^+$ and provide final predictions for $T_{bc}^0 \rightarrow \bar{B}^{*0} D^0$.

In the context of the QCD three-point sum rule method the strong decay $T_{bc}^0 \rightarrow B^{*-} D^+$ can be studied using the correlation function

$$\tilde{\Pi}_{\mu\nu}(p, p') = i^2 \int d^4x d^4y e^{i(p'y - px)} \langle 0 | \mathcal{T} \{ J_\mu^{B^*}(y) J^D(0) J_\nu^\dagger(x) \} | 0 \rangle. \quad (41)$$

Here $J_\nu(x)$, $J^D(x)$ and $J_\mu^{B^*}(x)$ are the interpolating currents for the tetraquark T_{bc}^0 and mesons D^+ and B^{*-} , respectively. The $J_\nu(x)$ is given by Eq. (2), whereas for the remaining two currents we use

$$J_\mu^{B^*}(x) = \bar{u}^i(x) \gamma_\mu b^j(x), \quad J^D(x) = \bar{d}^j(x) i \gamma_5 c^j(x). \quad (42)$$

The 4-momenta of the tetraquark T_{bc}^0 and meson B^{*-} are p and p' , therefore, the momentum of the meson D^+ is $q = p - p'$.

We follow the standard recipes and calculate the correlation function $\tilde{\Pi}_{\mu\nu}(p, p')$ using both the physical parameters of the particles involved into the process, and quark propagators. Separating the ground-state contribution from ones due to higher resonances and continuum states, for the physical side of the sum rule, we get

$$\begin{aligned} \tilde{\Pi}_{\mu\nu}^{\text{Phys}}(p, p') &= \frac{\langle 0 | J_{\mu}^{B^*} | B^{*-}(p', \epsilon') \rangle \langle 0 | J^D | D^+(q) \rangle}{(p'^2 - m_{B^*}^2)(q^2 - m_D^2)} \\ &\times \frac{\langle D^+(q) B^{*-}(p', \epsilon'^*) | T_{bc}^0(p, \epsilon) \rangle \langle T_{bc}^0(p, \epsilon^*) | J_{\nu}^{\dagger} | 0 \rangle}{(p^2 - m^2)} + \dots \end{aligned} \quad (43)$$

The function $\tilde{\Pi}_{\mu\nu}^{\text{Phys}}(p, p')$ can be simplified by expressing the matrix elements in terms of the tetraquark and mesons' physical parameters. The matrix element $\langle T_{bc}^0(p, \epsilon^*) | J_{\nu}^{\dagger} | 0 \rangle$ can be found using Eq. (4). We introduce also the matrix elements of the final-state mesons

$$\langle 0 | J^D | D^+ \rangle = \frac{m_D^2 f_D}{m_c}, \quad \langle 0 | J_{\mu}^{B^*} | B^{*-}(p', \epsilon') \rangle = m_{B^*} f_{B^*} \epsilon'_{\mu}. \quad (44)$$

Here m_D , m_{B^*} and f_D , f_{B^*} are the masses and decay constants of the mesons D^+ and B^{*-} , respectively. In Eq. (44) ϵ'_{μ} is the polarization vector of the meson B^{*-} . We model $\langle D^+(q) B^{*-}(p', \epsilon'^*) | T_{bc}^0(p, \epsilon) \rangle$ in the form

$$\langle D^+(q) B^{*-}(p', \epsilon'^*) | T_{bc}^0(p, \epsilon) \rangle = g_1(q^2) [(p \cdot p')(\epsilon \cdot \epsilon'^*) - (p \cdot \epsilon'^*)(p' \cdot \epsilon)] \quad (45)$$

and denote by $g_1(q^2)$ the strong form factor corresponding to the vertex $T_{bc}^0 B^{*-} D^+$. Then, it is not difficult to see that

$$\begin{aligned} \tilde{\Pi}_{\mu\nu}^{\text{Phys}}(p, p') &= g_1 \frac{m_D^2 m_{B^*} m f f_D f_{B^*}}{m_c (p^2 - m^2)(p'^2 - m_{B^*}^2)(q^2 - m_D^2)} \frac{1}{(q^2 - m_D^2)} \\ &\times \left[\frac{1}{2} (m^2 + m_{B^*}^2 - q^2) g_{\mu\nu} - p_{\mu} p'_{\nu} \right] + \dots \end{aligned} \quad (46)$$

The correlation function $\tilde{\Pi}_{\mu\nu}^{\text{Phys}}(p, p')$ has Lorentz structures proportional to $g_{\mu\nu}$ and $p_{\mu} p'_{\nu}$. We work with the invariant amplitude $\tilde{\Pi}^{\text{Phys}}(p^2, p'^2, q^2)$ that corresponds to the structure $g_{\mu\nu}$. The double Borel transformation of this amplitude over variables p^2 and p'^2 forms the phenomenological side of the sum rule.

To find the QCD side of the three-point sum rule, we calculate $\tilde{\Pi}_{\mu\nu}(p, p')$ in terms of the quark propagators and get

$$\begin{aligned} \tilde{\Pi}_{\mu\nu}^{\text{OPE}}(p, p') &= \int d^4x d^4y e^{i(p'y - px)} \left\{ \text{Tr} \left[\gamma_5 S_c^{jb}(-x) \gamma_5 \tilde{S}_b^{ia}(y-x) \gamma_{\mu} \tilde{S}_u^{ai}(x-y) \gamma_{\nu} S_d^{bj}(x) \right] \right. \\ &\quad \left. - \text{Tr} \left[\gamma_5 S_c^{jb}(-x) \gamma_5 \tilde{S}_b^{ia}(y-x) \gamma_{\mu} \tilde{S}_u^{bi}(x-y) \gamma_{\nu} S_d^{aj}(x) \right] \right\}. \end{aligned} \quad (47)$$

As in the case of the correlation function $\tilde{\Pi}_{\mu\nu}^{\text{Phys}}(p, p')$ here, we also isolate the structure $\sim g_{\mu\nu}$ and find the amplitude $\tilde{\Pi}^{\text{OPE}}(p^2, p'^2, q^2)$. The standard manipulations with invariant amplitudes yield the following sum rule

$$g_1(q^2) = \frac{2m_c}{m_{B^*} m f f_D f_{B^*}} \frac{q^2 - m_D^2}{m^2 + m_{B^*}^2 - q^2} e^{m^2/M_1^2} e^{m_{B^*}^2/M_2^2} \tilde{\Pi}^{\text{OPE}}(\mathbf{M}^2, \mathbf{s}_0, q^2), \quad (48)$$

where $\mathbf{M}^2 = (M_1^2, M_2^2)$, and $\mathbf{s}_0 = (s_0, s'_0)$ are the Borel and continuum threshold parameters. Apart from q^2 , the form factor $g_1(q^2)$ is also a function of the Borel and continuum threshold parameters which, for simplicity, are not shown explicitly in Eq. (48). The set (M_1^2, s_0) corresponds to initial tetraquark channel, whereas (M_2^2, s'_0) describes the channel of the heavy final

meson B^{*-} . Here, $\tilde{\Pi}^{\text{OPE}}(\mathbf{M}^2, \mathbf{s}_0, q^2)$ is the invariant amplitude $\tilde{\Pi}^{\text{OPE}}(p^2, p'^2, q^2)$ after the double Borel transformation and continuum subtraction procedures:

$$\tilde{\Pi}^{\text{OPE}}(\mathbf{M}^2, \mathbf{s}_0, q^2) = \int_{(m_b+m_c)^2}^{s_0} e^{-s/M_1^2} ds \int_{m_b^2}^{s'_0} ds' e^{-s'/M_2^2} \rho(s, s', q^2). \quad (49)$$

The spectral density $\rho(s, s', q^2)$ is calculated as an imaginary part of the relevant amplitude and contains the vacuum condensates up to dimension 5.

The parameters, i.e., the vacuum condensates and masses of the b and c quarks, which are necessary for numerical computations are given by Eq. (10). The mass and coupling of the tetraquark T_{bc}^0 have been calculated in the present work. In computations we also use $m_{D^0} = (1864.84 \pm 0.07)$ MeV and $f_{D^0} = (203.7 \pm 4.7)$ MeV, $m_{B^*} = (5325.2 \pm 0.4)$ MeV and $f_{B^*} = (210 \pm 6)$ MeV, respectively. Parameters of the D meson can be read out from Table 3. The auxiliary parameters for the T_{bc}^0 channel are chosen in accordance with Eq. (11). For the set (M_2^2, s'_0) we use the regions

$$M_2^2 \in [4.5, 5.5] \text{ GeV}^2, \quad s'_0 \in [32, 34] \text{ GeV}^2. \quad (50)$$

The sum rule method for $g_1(q^2)$ gives reliable predictions only for $q^2 < 0$. Therefore, we introduce a variable $Q^2 = -q^2$ and denote the new function as $g_1(Q^2)$. The width of the decay $T_{bc}^0 \rightarrow B^{*-} D^+$ has to be computed using the strong form factor at the mass shell of the D^+ meson $q^2 = m_D^2$. This point is not accessible to sum rule computations, but the problem can be solved by employing a fit function $\mathcal{G}_1(Q^2)$, which at the momenta $Q^2 > 0$ coincides with QCD sum rule predictions, but can be extrapolated to the region of $Q^2 < 0$. Then, using the interpolating function $\mathcal{G}_1(Q^2)$, one can find $g_1(-m_D^2)$. The function $\mathcal{G}_1(Q^2)$ does not differ from ones that we have used in Eq. (26), a difference being only in replacement of the fitting mass with the mass of the tetraquark $m_{\text{fit}}^2 \rightarrow m^2$

$$\mathcal{G}_1(Q^2) = \mathcal{G}_0^1 \exp \left[\tilde{c}_1^1 \frac{Q^2}{m^2} + \tilde{c}_2^1 \left(\frac{Q^2}{m^2} \right)^2 \right]. \quad (51)$$

The parameters \mathcal{G}_0^1 , \tilde{c}_1^1 and \tilde{c}_2^1 have been fixed from numerical analyses $\mathcal{G}_0^1 = 1.11$, $\tilde{c}_1^1 = 14.33$, and $\tilde{c}_2^1 = -120.69$. This function at the mass shell $Q^2 = -m_D^2$ gives

$$g_1 \equiv \mathcal{G}_1(-m_D^2) = (0.25 \pm 0.03) \text{ GeV}^{-1}. \quad (52)$$

The width of decay $T_{bc}^0 \rightarrow B^{*-} D^+$ is determined by the formula

$$\Gamma[T_{bc}^0 \rightarrow B^{*-} D^+] = \frac{g_1^2 m_{B^*}^2}{24\pi} \lambda \left(3 + 2 \frac{\lambda^2}{m_{B^*}^2} \right), \quad (53)$$

where $\lambda = \lambda(m^2, m_{B^*}^2, m_D^2)$.

Using Eqs. (52) and (53), one can easily calculate the width of the decay $T_{bc}^0 \rightarrow B^{*-} D^+$

$$\Gamma[T_{bc}^0 \rightarrow B^{*-} D^+] = (31.1 \pm 6.2) \text{ MeV}. \quad (54)$$

The second process $T_{bc}^0 \rightarrow \overline{B}^{*0} D^0$ can be explored by the same manner. Here, we take into account that interpolating currents have the following forms

$$J_{\mu}^{\bar{B}^{*0}}(x) = \bar{d}^i(x)\gamma_{\mu}b^i(x), \quad J^{D^0}(x) = \bar{u}^j(x)i\gamma_5c^j(x). \quad (55)$$

The remaining operations are standard manipulations in the context of the sum rule method. Therefore, we do not see a necessity to provide a detailed information on them. Let us note only that the fit function $\mathcal{G}_2(Q^2)$ has the parameters $\mathcal{G}_0^1 = 1.11$, $\tilde{c}_1^2 = 14.40$, and $\tilde{c}_2^2 = -121.11$. At the mass shell of the meson D^0 for the strong coupling we get

$$g_2(-m_{D^0}^2) = (0.26 \pm 0.03) \text{ GeV}^{-1}, \quad (56)$$

and

$$\Gamma[T_{bc}^0 \rightarrow \bar{B}^{*0}D^0] = (32.4 \pm 6.3) \text{ MeV}. \quad (57)$$

Then, in the second scenario the full width of the axial-vector tetraquark T_{bc}^0 is

$$\Gamma_{\text{full}} = (63.5 \pm 8.9) \text{ MeV}. \quad (58)$$

This prediction for Γ_{full} is the main result obtained utilizing the second option for m .

6. Analysis and concluding notes

In the present work we have studied, in a rather detailed form, the axial-vector tetraquark T_{bc}^0 . As we have emphasized in Section 1, there are different predictions for its mass and stability properties in the literature. We have calculated the mass m and coupling f of this tetraquark by means of the QCD sum rule method. Our result for m does not allow us to solve unambiguously a problem with stability of the tetraquark T_{bc}^0 . Thus, the central value of the mass 7105 MeV obtained in the present work is below both the strong and electromagnetic thresholds, and therefore in this scenario T_{bc}^0 can transform to conventional mesons only through the weak transitions. But taking into account theoretical errors of computations and using the maximal value of $m = 7260$ MeV, we see that T_{bc}^0 becomes unstable against the strong and electromagnetic decays. We have explored both of these scenarios and calculated the width and lifetime of T_{bc}^0 .

In the framework of the first scenario, we have calculated the partial widths of the semileptonic $T_{bc}^0 \rightarrow T_{cc}^+ l \bar{\nu}_l$ ($l = e, \mu$ and τ) and two-body weak decays $T_{bc}^0 \rightarrow T_{cc}^+ \pi^-$ (K^-, D_s^-, D_s^-) of T_{bc}^0 . Using obtained information on these processes we have evaluated its full width $\Gamma_{\text{full}} = (3.98 \pm 0.51) \times 10^{-10}$ MeV and mean lifetime $\tau \approx 1.7$ ps. In our previous work [20] we computed the same parameters of the scalar tetraquark Z_{bc}^0 . It is instructive to compare parameters of the scalar and axial-vector $bc\bar{u}\bar{d}$ states with each other. The scalar compound Z_{bc}^0 with the mass 6660 MeV has a more stable nature and lives $\tau \approx 21$ ps which is considerably longer than $\tau \approx 1.7$ ps of the T_{bc}^0 .

It is known that, the scalar tetraquark T_{cc}^+ decays strongly to a pair of conventional D^+D^0 mesons [22]. Then, we can estimate branching ratios of different weak decay channels of T_{bc}^0 ; corresponding predictions are collected in Table 4.

If mass of the tetraquark T_{bc}^0 is at around of 7260 MeV, it decays strongly to conventional mesons. In present article we have explored this scenario as well, and calculated partial widths of S -wave decay channels $T_{bc}^0 \rightarrow B^{*-}D^+$ and $T_{bc}^0 \rightarrow \bar{B}^{*0}D^0$. The full width $\Gamma_{\text{full}} = (63.5 \pm 8.9)$ MeV of T_{bc}^0 estimated employing these dominant strong decays characterizes T_{bc}^0 as a typical unstable tetraquark. Branching ratios of the strong decay modes are equal to

$$\mathcal{BR}(T_{bc}^0 \rightarrow B^{*-}D^+) \simeq 0.49, \quad \mathcal{BR}(T_{bc}^0 \rightarrow \bar{B}^{*0}D^0) \simeq 0.51. \quad (59)$$

Table 4

The nonleptonic decay channels of the tetraquark T_{bc}^0 and corresponding branching ratios.

Channels	\mathcal{BR}
$D^+ D^0 e^- \bar{\nu}_e$	0.36
$D^+ D^0 \mu^- \bar{\nu}_\mu$	0.36
$D^+ D^0 \tau^- \bar{\nu}_\tau$	0.11
$D^+ D^0 \pi^-$	0.043
$D^+ D^0 K^-$	0.003
$D^+ D^0 D^-$	0.004
$D^+ D^0 D_s^-$	0.12

Theoretical errors of the sum rule computations and, as a result, different predictions for the mass of the tetraquark T_{bc}^0 do not allow us to interpret it unambiguously as strong- and electromagnetic-interaction stable or unstable particle. The results obtained in the present article can be refined by including into analysis other decay channels of the tetraquark T_{bc}^0 . Thus, the weak transition $c \rightarrow W^+ s$ can give rise to the semileptonic decays $T_{bc}^0 \rightarrow T_{bs}^- \bar{l} \nu_l$, where T_{bs}^- is the scalar tetraquark with content $[bs] [\bar{u}\bar{d}]$. There are also nonleptonic decays of T_{bc}^0 generated by this transition, where final states apart from T_{bs}^- contain the conventional mesons π^+ or K^+ . All these questions deserve further detailed investigations, which will provide useful information on features of the axial-vector tetraquark T_{bc}^0 and may be useful for its experimental and theoretical investigations.

Declaration of competing interest

We declare that there is no conflict of interests regarding this manuscript.

References

- [1] J.P. Ader, J.M. Richard, P. Taxil, Phys. Rev. D 25 (1982) 2370.
- [2] H.J. Lipkin, Phys. Lett. B 172 (1986) 242.
- [3] S. Zouzou, B. Silvestre-Brac, C. Gignoux, J.M. Richard, Z. Phys. C 30 (1986) 457.
- [4] J. Carlson, L. Heller, J.A. Tjon, Phys. Rev. D 37 (1988) 744.
- [5] D. Janc, M. Rosina, Few-Body Syst. 35 (2004) 175.
- [6] Y. Cui, X.L. Chen, W.Z. Deng, S.L. Zhu, HEPNP 31 (2007) 7.
- [7] J. Vijande, A. Valcarce, K. Tsushima, Phys. Rev. D 74 (2006) 054018.
- [8] D. Ebert, R.N. Faustov, V.O. Galkin, W. Lucha, Phys. Rev. D 76 (2007) 114015.
- [9] F.S. Navarra, M. Nielsen, S.H. Lee, Phys. Lett. B 649 (2007) 166.
- [10] M.L. Du, W. Chen, X.L. Chen, S.L. Zhu, Phys. Rev. D 87 (2013) 014003.
- [11] T. Hyodo, Y.R. Liu, M. Oka, K. Sudoh, S. Yasui, Phys. Lett. B 721 (2013) 56.
- [12] A. Esposito, M. Papinutto, A. Pilloni, A.D. Polosa, N. Tantalo, Phys. Rev. D 88 (2013) 054029.
- [13] R. Aaij, et al., LHCb Collaboration, Phys. Rev. Lett. 119 (2017) 112001.
- [14] M. Karliner, J.L. Rosner, Phys. Rev. Lett. 119 (2017) 202001.
- [15] E.J. Eichten, C. Quigg, Phys. Rev. Lett. 119 (2017) 202002.
- [16] S.S. Agaev, K. Azizi, B. Barsbay, H. Sundu, Phys. Rev. D 99 (2019) 033002.
- [17] G.-Q. Feng, X.-H. Guo, B.-S. Zou, arXiv:1309.7813 [hep-ph].
- [18] A. Francis, R.J. Hudspith, R. Lewis, K. Maltman, Phys. Rev. D 99 (2019) 054505.
- [19] T.F. Carames, J. Vijande, A. Valcarce, arXiv:1812.08991 [hep-ph].
- [20] H. Sundu, S.S. Agaev, K. Azizi, Eur. Phys. J. C 79 (2019) 753.

- [21] H. Sundu, B. Barsbay, S.S. Agaev, K. Azizi, *Eur. Phys. J. A* 54 (2018) 124.
- [22] S.S. Agaev, K. Azizi, H. Sundu, *Phys. Rev. D* 99 (2019) 114016.
- [23] M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, *Phys. Rev. Lett.* 83 (1999) 1914.
- [24] M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, *Nucl. Phys. B* 591 (2000) 313.
- [25] A.J. Buras, M. Jamin, M.E. Lautenbacher, *Nucl. Phys. B* 400 (1993) 75.
- [26] M. Ciuchini, E. Franco, G. Martinelli, L. Reina, *Nucl. Phys. B* 415 (1994) 403.
- [27] G. Buchalla, A.J. Buras, M.E. Lautenbacher, *Rev. Mod. Phys.* 68 (1996) 1125.