



Single neutron transfer on ^{23}Ne and its relevance for the pathway of nucleosynthesis in astrophysical X-ray bursts

G. Lotay^{a,*}, J. Henderson^{a,b}, W.N. Catford^a, F.A. Ali^{c,d}, J. Berean^b, N. Bernier^{b,e,1}, S.S. Bhattacharjee^{b,2}, M. Bowry^{b,3}, R. Caballero-Folch^b, B. Davids^{b,f}, T.E. Drake^g, A.B. Garnsworthy^b, F. Ghazi Moradi^c, S.A. Gillespie^{b,4}, B. Greaves^c, G. Hackman^b, S. Hallam^a, D. Hymers^c, E. Kasanda^c, D. Levy^b, B.K. Luna^h, A. Mathews^b, Z. Meiselⁱ, M. Moukaddam^{a,5}, D. Muecher^{b,c}, B. Olaizola^{b,6}, N.A. Orr^j, H.P. Patel^b, M.M. Rajabali^h, Y. Saito^{b,e}, J. Smallcombe^{b,7}, M. Spencer^{a,b}, C.E. Svensson^c, K. Whitmore^f, M. Williams^{b,k}

^a Department of Physics, University of Surrey, Guildford, GU2 7XH, United Kingdom

^b TRIUMF, Vancouver, British Columbia, V6T 2A3, Canada

^c Department of Physics, University of Guelph, Guelph, Ontario N1G 2W1, Canada

^d Department of Physics, College of Education, University of Sulaimani, P.O. Box 334, Sulaimani, Kurdistan Region, Iraq

^e Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

^f Department of Physics, Simon Fraser University, Burnaby, British Columbia, V5A 1S6, Canada

^g Department of Physics, University of Toronto, Toronto, Ontario M5S 1A7, Canada

^h Department of Physics, Tennessee Technological University, Cookeville, TN 38505, USA

ⁱ Department of Physics & Astronomy, Ohio University, Athens, OH 45701, USA

^j LPC-Caen, IN2P3/CNRS, UniCaen, ENSICAEN, Normandie Université, 14000 Caen, France

^k Department of Physics, University of York, Heslington, York, YO10 5DD, United Kingdom

ARTICLE INFO

Article history:

Received 1 April 2022

Received in revised form 24 June 2022

Accepted 28 July 2022

Available online 2 August 2022

Editor: B. Blank

ABSTRACT

We present new experimental measurements of resonance strengths in the astrophysical $^{23}\text{Al}(p, \gamma)^{24}\text{Si}$ reaction, constraining the pathway of nucleosynthesis beyond ^{22}Mg in X-ray burster scenarios. Specifically, we have performed the first measurement of the (d, p) reaction using a radioactive beam of ^{23}Ne to explore levels in ^{24}Ne , the mirror analog of ^{24}Si . Four strong single-particle states were observed and corresponding neutron spectroscopic factors were extracted with a precision of $\sim 20\%$. Using these spectroscopic factors, together with mirror state identifications, we have reduced uncertainties in the strength of the key $\ell = 0$ resonance at $E_r = 157$ keV, in the astrophysical $^{23}\text{Al}(p, \gamma)$ reaction, by a factor of 4. Our results show that the $^{22}\text{Mg}(p, \gamma)^{23}\text{Al}(p, \gamma)$ pathway dominates over the competing $^{22}\text{Mg}(\alpha, p)$ reaction in all but the most energetic X-ray burster events ($T > 0.85$ GK), significantly affecting energy production and the preservation of hydrogen fuel.

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author.

E-mail address: g.lotay@surrey.ac.uk (G. Lotay).

¹ Present address: Department of Physics and Astronomy, University of the Western Cape, P/B X17, Bellville, ZA-7535, South Africa.

² Present address: Institute of Experimental and Applied Physics, Czech Technical University in Prague, Husova 240/5, 110 00 Prague 1, Czech Republic.

³ Present address: School of Computing, Engineering and Physical Sciences, University of the West of Scotland, Paisley PA1 2BE, United Kingdom.

⁴ Present address: National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, Michigan 48824, USA.

⁵ Present Address: Universit'e de Strasbourg, IPHC, 23 rue du Loess, 67037 Strasbourg, France.

⁶ Present Address: ISOLDE, CERN, CH-1211 Geneva 23, Switzerland.

⁷ Present Address: University of Liverpool, Liverpool, L69 3BX, UK.

<https://doi.org/10.1016/j.physletb.2022.137361>

0370-2693/© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

Type-I X-ray bursts represent thermonuclear explosions on the surfaces of accreting neutron stars in close binary systems [1–3]. They exhibit dramatic, recurrent increases in luminosity and constitute the most frequent stellar eruptions to occur in our Galaxy. In between bursts, energy is generated at a constant rate by the β -limited hot CNO cycles [4,5]. However, as the temperature of the accreted material increases, the triple- α reaction becomes favourable, igniting the burst, and nucleosynthesis proceeds along the proton-rich side of stability via the αp process [6] [a series of (p, γ) and (α, p) reactions], and the rp process [3] [a series of (p, γ) reactions and β^+ decays], ending in the Sn-Te mass region.

Recently, advances in computing power have allowed for detailed models of X-ray burst nucleosynthesis to be constructed

[6–9], incorporating complex reaction networks and hundreds of nuclear species ranging from stable isotopes up to the proton drip line. Strikingly, despite the vast number of reactions included, only a handful of nuclear processes have been highlighted as having a noticeable effect on the observational properties of X-ray bursts [10–12]. In particular, the $^{23}\text{Al}(p, \gamma)^{24}\text{Si}$ reaction, which permits flow beyond masses of $A = 22$ in the early phases of the rp process, is postulated to have a strong influence on the inferred surface gravitational redshift $(1+z)$ [12]. The redshift is directly related to the neutron star compactness [13] and thus, any experimental constraints placed on the $^{23}\text{Al}(p, \gamma)$ reaction rate will help to reveal new facets of the underlying compact objects involved. Furthermore, at the ^{22}Mg , rp -process waiting point, the $^{22}\text{Mg}(p, \gamma)^{23}\text{Al}(p, \gamma)$ reaction sequence is expected to compete significantly with the $^{22}\text{Mg}(\alpha, p)$ reaction [14], affecting the overall energy generation in X-ray bursters. Specifically, a prevailing $^{22}\text{Mg}(p, \gamma)^{23}\text{Al}(p, \gamma)$ pathway results in less energetic burning during the burst rise, preserving hydrogen for later burning and extending the burst tail. The exceptional measurements now available for the structure of burst light curves [15,16] are amenable to confront simulations of the burst explosions.

Previous studies of the $^{23}\text{Al}(p, \gamma)$ reaction [17–20] indicate that the rate is dominated by resonant capture on the $5/2^+$ ground state of ^{23}Al to excited states above the proton-emission threshold energy of 3292(19) keV in ^{24}Si [21]. However, the strengths of these resonances remain uncertain, due to the scarcity of experimental data. Most recently, Wolf et al. utilised the $^{23}\text{Al}(d, n)$ reaction to investigate the properties of excited states in ^{24}Si [20]. In that study [20], γ decays were observed from three excited states, including the key $\ell = 0$, proton-unbound resonant level at 3449(5) keV, which is expected to have the most significant influence on the $^{23}\text{Al}(p, \gamma)$ reaction over the temperature range of X-ray bursts. Moreover, by measuring angle-integrated cross sections of excited levels in ^{24}Si , Wolf et al. were able to place the first constraints on proton spectroscopic factors, reducing uncertainties in both the direct and resonant capture components of the $^{23}\text{Al}(p, \gamma)$ reaction [20]. However, the absolute values of spectroscopic factors reported in Ref. [20] carry large uncertainties of order 60% because their extraction relied on the use of shell-model calculations to determine the relative contributions of multiple ℓ -transfers. For example, where states are populated by a mixture of $\ell = 0$ and $\ell = 2$ transfer, the work of Ref. [20] was forced to use the ratio of strengths predicted by the shell model, but it has been pointed out [22] that the shell model consistently fails to predict this ratio correctly. Consequently, the rate of the $^{23}\text{Al}(p, \gamma)$ reaction is still weakly constrained over the temperature range of Type-I X-ray bursts and a more robust experimental measurement is demanded.

A direct measurement of the $^{23}\text{Al}(p, \gamma)$ reaction is not presently feasible. As such, any further experimental constraints must rely on indirect techniques. In this regard, several studies have shown that precise evaluations of proton capture reactions may be achieved via the concept of isospin [23–25]. Specifically, neutron spectroscopic factors of excited states in mirror nuclei, that correspond to analogs of (p, γ) resonances, can be used to accurately determine the strengths of resonances governing the rate of stellar reactions in explosive astrophysical environments [23–25]. In this Letter, we present a first experimental measurement of the $^{23}\text{Ne}(d, p)$ transfer reaction to study excited states in ^{24}Ne . These levels correspond to $T = 2$, mirror analogs of key resonant states in the $^{23}\text{Al}(p, \gamma)^{24}\text{Si}$ reaction. By coupling the TIGRESS γ -array [26] to the SHARC charged-particle detection system [27], neutron spectroscopic factors were extracted to a precision of $\sim 20\%$. This reduces uncertainties in $^{23}\text{Al} + p$ resonance strengths by a factor ~ 4 and, hence, defines the relative importance of the




Fig. 1. Excitation energy spectrum obtained following $^{23}\text{Ne}(d, p)$ transfer. (Inset) Gamma decays observed with a gate placed on the 3871-keV excitation energy peak.

$^{22}\text{Mg}(p, \gamma)^{23}\text{Al}(p, \gamma)$ and $^{22}\text{Mg}(\alpha, p)$ reaction sequences over the temperature range of X-ray bursts.

A beam of radioactive $^{23}\text{Ne}^{2+}$ ions was accelerated to 8.0 MeV/nucleon and an intensity of $\sim 2 \times 10^4$ pps, by the ISAC-II facility at TRIUMF and bombarded a 1 mg/cm² (CD_2)_n foil for 93 hrs. Prompt γ rays were recorded using the TIGRESS array of 12 Compton-suppressed HPGe detectors [26], while charged particles including protons from the $^{23}\text{Ne}(d, p)$ reaction were measured in the SHARC silicon array [27]. Beyond the target, 40 cm downstream, the TRIFOIL detector [28,29] was placed (a 20 μm foil of BC400 plastic scintillator viewed by three photomultiplier tubes and mounted behind a passive stopper foil of 110 μm Al). The TRIFOIL setup (a) stopped the ^{23}Ne beam and ^{24}Ne reaction products in the scintillator and counted them, (b) stopped the ^{23}Ne beam contaminant ($\sim 40\%$ of the beam) and ^{24}Na reaction products in the Al foil so that they had no TRIFOIL tag and (c) also in the Al, stopped fusion-evaporation products from reactions on carbon in the CD_2 target. The TRIFOIL also gave a direct measurement of the average counting rate of the beam over the entire 93 hours of data acquisition to a precision of $< 1\%$. The rejection of ^{23}Ne -induced events was verified by the complete removal of ^{24}Na γ -ray peaks when imposing the TRIFOIL requirement. The beam composition was also measured at regular intervals using a Bragg ionization detector [30] and background from other contaminant isobars was found to be negligible. Energy and efficiency calibrations were performed using standard γ -ray (^{152}Eu and ^{60}Co) and charged-particle (triple alpha) sources. The absolute normalisation was determined using the measured number of incident ^{23}Ne ions, the target thickness and the H:D ratio, as determined from elastic scattering around $\theta_{\text{cm}} = 50^\circ$ measured simultaneously throughout the acquisition.

Fig. 1 illustrates the excitation energy of states in ^{24}Ne populated via the (d, p) reaction. As can be seen, four strongly populated states are observed at 0, 1981, 3871 and 4886 keV, in good agreement with previously reported 0_1^+ , 2_1^+ , 2_2^+ and 3_1^+ levels in ^{24}Ne [32] and the theoretical calculations of Ref. [33]. That being said, additional 4_1^+ and 0_2^+ excited states are also expected in this energy region in ^{24}Ne at 3962 and 4765 keV [32], respectively, which would not be resolvable using proton detection alone, due to the ~ 300 keV (FWHM) excitation energy resolution of SHARC. In this regard, the simultaneous detection of γ rays is of crucial importance. In particular, by placing gates across the observed energy peaks at 3871 and 4886 keV, and viewing coincident γ rays within the TIGRESS array, it was possible to rule out any significant population of the 4_1^+ , 3962-keV and 0_2^+ , 4765-keV excited states via the $^{23}\text{Ne}(d, p)$ reaction. For example, when a gate was placed across the 3871-keV proton peak, we observe 1890- and 1981-keV

Table 1

Properties of excited states in the $T = 2$, $A = 24$ system, as determined in the present work and reported in earlier literature. Excitation energies are given in keV and shell-model spectroscopic factors were determined using the USDA interaction [31]. In Ref. [32], no uncertainties for E_x (^{24}Ne) are given, but we expect ≤ 2 keV based on HPGe calibration. The present $C^2S_{(d,p)}$ is the summed $\ell = 2$ strength, extracted assuming transfer to $0d_{5/2}$ (errors and limits, see text). For comparison, C^2S_{SM} is the sum of USDA shell-model values for $0d_{5/2}$ and $0d_{3/2}$.

E_x (^{24}Ne) [32] ^a	J^π	ℓ_p	$C^2S_{(d,p)}$	C^2S_{SM}	$C^2S_{(d,n)}$ [20]	Analog State in ^{24}Si [20]
0	0^+	2	3.42(68)	3.50	≤ 2.8	0
1981	2^+	0	0.28(6)	0.28	0.6(2)	1874(3)
		2	0.37(7)	0.19	0.4(1)	
3871	2^+	0	0.44(9)	0.42	0.7(4)	3449(5)
		2	0.23(5)	0.17	0.3(2)	
3962	4^+	2	≤ 0.012	0.014	0.07(4)	3471(6)
4765	0^+	2	≤ 0.19	0.21	0.8(4)	4170 ^b
4886	3^+	0	0.56(11)	0.58		4470 ^b
		2	≤ 0.19	0.17		

^a Excitation energy uncertainties are not provided in Ref. [32] but are assumed to be ~ 1 keV, based on the observation of γ -ray transitions.

^b Taken from theoretical calculations of Ref. [33].




Fig. 2. Angular distributions of protons in the $^{23}\text{Ne}(d,p)$ reaction compared with best-fit TWOFNR calculations (statistical error bars only). Data for the 4886-keV state may include the unresolved 4765 keV, 0^+ state (for the $\ell = 2$ component only). For details, see text. Uncertainties shown are statistical only.

γ -ray peaks of equal intensity from the cascade decay from 3871 keV (inset of Fig. 1). The numbers of counts are 76 ± 10 and 84 ± 10 , respectively. The surplus for the 1981-keV peak is 8 ± 14 which is consistent with zero and gives a 2σ upper limit (allowing for the double counting) of 12% of the combined population of the two states. This is the basis of the limit on the spectroscopic factor for the 3962-keV state in Table 1. Consequently, we conclude that the 3962-keV excited state in ^{24}Ne was not appreciably populated and, based on an upper limit analysis of 1981-keV transitions originating from the 4_1^+ state, we set a stringent upper limit on its spectroscopic factor, $C^2S_{(\ell=2)} \leq 0.012$. This is in agreement with our shell model calculations using NuShellX [34] with the USD-A interaction [31], which predict $C^2S_{(\ell=2)} = 0.01$ for the 4_1^+ level in ^{24}Ne . In contrast, a similar procedure to the above was not possible for the expected 4765/4886-keV doublet due to a considerable level of background in the γ -ray energy region of interest. Whilst the observed $\ell = 0$ angular distribution for this doublet, shown in Fig. 2, may be ascribed entirely to the known 3^+ , 4886-keV level in ^{24}Ne [32], we adopt an upper limit of 0.19 for the $\ell = 2$ component of the spectroscopic factor for both the 4765- and 4886-keV excited states in ^{24}Ne .

An angular distribution analysis of the 0-, 1981-, 3871- and 4886-keV excited states in ^{24}Ne , shown in Fig. 2, confirms the spin-parity assignments of Ref. [32]. However, with the exception of the ground state, which necessarily exhibits a pure $\ell = 2$ character, the measured distributions indicate strong mixing between

$\ell = 0$ and $\ell = 2$ transfer for all levels. These observed distributions were then compared with reaction calculations in the Adiabatic Distorted Wave Approximation (ADWA) performed, using the code TWOFNR [35]. Here, the Johnson-Soper adiabatic model [36] was employed with standard parameters [37] using zero range and the Koning-Delaroche [38] global nucleon-nucleus optical potential. We estimate an uncertainty in the overall cross section normalization of $\sim 20\%$, with the dominant contribution coming overwhelmingly from the modelling of the (d,p) reaction itself [37]. The solid angle was calculated accurately from the known geometry, the fitted position of the beam spot and omitting the detector strips excluded from the analysis. The systematic uncertainty in the normalisation of the data arises principally from the uncertainty in the target thickness (taken as 10%) since the total number of incident particles was precisely given by a direct measurement included continuously in the data stream. A summary of the properties of excited states in ^{24}Ne determined in this work is given in Table 1, together with a comparison with our shell-model calculations using the USDA interaction [31]. A proposed matching of analog levels in ^{24}Si is also shown [20]. We have adopted a number of mirror assignments from earlier work [20] and, although the spin-parity assignments of the 3449- and 3471-keV excited states in ^{24}Si are not uniquely defined, we propose analog matchings to the 2_1^+ , 3871-keV and 4_1^+ , 3962-keV levels in ^{24}Ne , respectively, based on mirror energy differences. Specifically, a pairing to the 0_2^+ state would require a very large mirror energy shift of ~ 1.3 MeV (although we note that a recent study [39] suggested that such an assignment may be possible).

The present results show excellent agreement with shell model calculations, especially for $\ell = 0$ transfers. Notably, we find that $\ell = 2$ strengths for strongly mixed states can deviate considerably from theory, as was previously highlighted in Ref. [22]. This is particularly relevant to the extraction of astrophysical data. Specifically, the authors of Ref. [20] were forced to rely on shell model ratios of $\ell = 0$ and $\ell = 2$ strengths in order to analyse their angle-integrated cross sections, but their extracted $\ell = 0$ values are then susceptible to inaccuracies in the shell model theory (the $\ell = 0$ strength determines the important resonance parameters for astrophysics). The present work measures the $\ell = 2$ and $\ell = 0$ strengths independently of any prior constraints and indeed we find clear differences with the results from Ref. [20]. In particular, the values of $C^2S_{\ell=0}$ of the 2_1^+ and 2_2^+ excited levels in the $T = 2$ system are found, respectively, to be 0.28(6) [compared to 0.6(2) for Ref. [20]] and 0.44(9) [compared to 0.7(4)]. The differences in both magnitude and uncertainty have important consequences for the role of the $^{23}\text{Al}(p,\gamma)$ reaction in determining the development of X-ray bursters.

For an evaluation of the astrophysical $^{23}\text{Al}(p,\gamma)$ reaction rate, we consider the contribution of excited states in ^{24}Si at $E_x =$

Table 2

Properties of resonant states in the $^{23}\text{Al}(p, \gamma)$ reaction used in the present analysis, together with a comparison to resonance strengths based on spectroscopic factors reported in Ref. [20]. Excitation energies in ^{24}Si and present C^2S values are as in Table 1. These were used to calculate Γ_p and (together with the Γ_γ from USDA shell-model calculations, see text) $\omega\gamma$. For states shown in Table 1 with $\ell = 0$ and $\ell = 2$ contributions, only the $\ell = 0$ is included here since it overwhelmingly dominates the resonance strength. Upper limits have been determined to a 68% confidence level.

$E_x, ^{24}\text{Si}$ (keV)	E_r (keV)	J^π	ℓ_p	C^2S	Γ_p (eV)	Γ_γ (eV)	Present $\omega\gamma$ (eV)	Previous $\omega\gamma^a$ (eV)
3449(5)	157(20)	2 ⁺	0	0.44(9)	$8.2(17) \times 10^{-5}$	1.8×10^{-2}	$3.4(7) \times 10^{-5}$	$5.4(31) \times 10^{-5}$
3471(6)	179(20)	(4 ⁺) ^b	2	≤ 0.012	$\leq 1.0 \times 10^{-7}$	2.5×10^{-4}	$\leq 7.6 \times 10^{-8}$	$4.4(26) \times 10^{-7}$
		(0 ⁺) ^b	2	≤ 0.19	$\leq 1.6 \times 10^{-6}$	3.0×10^{-5}	$\leq 1.3 \times 10^{-7}$	$5.6(28) \times 10^{-7}$
4170 ^c	878 ^c	0 ⁺	2	≤ 0.19	≤ 32	1.6×10^{-4}	1.3×10^{-5} ^d	-
4470 ^c	1178 ^c	3 ⁺	0	0.56(11)	3.0×10^4	8.9×10^{-3}	5.2×10^{-3} ^d	-

^a Previous resonance strengths have been estimated based on spectroscopic factors reported in Ref. [20].

^b For the 179-keV resonance, we currently favour a 4⁺ assignment based on mirror energy difference arguments. However, for completeness, we provide resonance strength determinations for both 4₁⁺ and 0₂⁺ assignments.

^c Adopted from Ref. [33].

^d Resonance strength determination dominated by theoretically calculated γ -ray partial width.




Fig. 3. (Top) Contribution of the 157-keV resonance to the $^{23}\text{Al}(p, \gamma)$ stellar reaction rate based only on uncertainties in its resonance strength from the present measurement in comparison with those of Ref. [20]. (Bottom) Percentage flow of material from the ^{22}Mg waiting point, through the (α, p) process, based on 1.5σ uncertainties in the present $^{23}\text{Al}(p, \gamma)$ reaction rate in comparison with equivalent 1.5σ uncertainties in the previously reported rate of Wolf et al. [20]. In this case, the contribution of all resonances have been included, as well as uncertainties associated with the reaction Q-value.

3449, 3471, 4170 and 4470 keV, corresponding to resonances in the $^{23}\text{Al} + p$ system at $E_r = 157, 179, 878$ and 1178 keV, respectively (see Table 2; the direct capture component is expected to be negligible for temperatures, $T \geq 0.1$ GK, and we do not foresee any significant departure from the value previously reported in Ref. [19] based on the present results). Here, we adopt spectroscopic factors obtained in the present work for the determination of proton partial widths. The spectroscopic factors of mirror analog states are expected to be nearly identical [40,41]. In assessing the validity of this statement, we performed a comparison of proton

and neutron spectroscopic factors in the mirror systems: $^{17}\text{F}-^{17}\text{O}$ [42,43], $^{21}\text{Na}-^{21}\text{Ne}$ [44–47], $^{25}\text{Al}-^{25}\text{Mg}$ [48,49], $^{29}\text{P}-^{29}\text{Si}$ [50–52], and $^{33}\text{Cl}-^{33}\text{S}$ [52–54], up to excitation energies of $\sim 4 - 5$ MeV. We found that spectroscopic factors agree to within $\sim 12\%$, with a standard deviation of $\sim 10\%$. This is well within the known $\sim 20\%$ uncertainty associated with the extraction of spectroscopic factors from experimentally measured cross sections. As such, we conclude that spectroscopic factors obtained for excited states in ^{24}Ne may be adopted for analog levels in ^{24}Si to a precision consistent with experimental uncertainties. In contrast, γ -ray partial widths were calculated using transition densities from our USDA shell-model calculations, adapted to the actual transition energies between the ^{24}Si states shown in Table 1. It should be noted that the values of Γ_γ are negligible for the determination of $\omega\gamma$ for the 157- and 179-keV resonances, as they are significantly larger than the corresponding proton partial widths, Γ_p . However, in the case of the 878- and 1178-keV states, the opposite is true. For the 1178-keV state, the present value of Γ_γ is in good agreement with the USD results of Ref. [33], whereas our current estimate for the 878-keV resonance is a factor 2 smaller. In the case of the latter, we note that while there is a discrepancy between the USDA and USD calculations, the contribution of the resonance at 878 keV to the overall $^{23}\text{Al}(p, \gamma)$ stellar reaction rate is negligible for temperatures, $T = 0.1 - 2$ GK.

In agreement with previous studies [17,20], we find that the $\ell = 0$ resonance at 157 keV makes the most significant contribution to the $^{23}\text{Al}(p, \gamma)$ stellar reaction rate for $T = 0.1 - 2$ GK. However, in contrast to previous work [20], uncertainties in the strength of the 157-keV resonance have been reduced by a factor of ~ 4 . Consequently, in order to fully assess the astrophysical implications of the current study, we have estimated the uncertainty in the total reaction rate based on the present resonance energies (which have an uncertainty dominated by the reaction Q-value [21]) and the resonance strengths (with uncertainties dominated by the spectroscopic factors, but now much improved). A 1.5σ confidence interval was calculated to properly account for experimental uncertainties in the $^{23}\text{Al}(p, \gamma)$ reaction parameters, which we note also accounts for uncertainties in the $^{22}\text{Mg}(\alpha, p)$ reaction cross section [14]. We note that the authors of Ref. [14] utilised the TALYS code to extend their data into the Gamow energy window for Type-I X-ray bursts and, as such, we presently estimate a $\sim 60\%$ uncertainty in the $^{22}\text{Mg}(\alpha, p)$ rate – this does not include uncertainties associated with centre-of-mass energies in Ref. [14]. By using the Saha equation to determine the $^{22}\text{Mg}(p, \gamma)/^{23}\text{Al}(p, \gamma)$ equilibrium [55], and comparing the present results with those of Ref. [20], we have been able to investigate the relative competition between the $^{22}\text{Mg}(p, \gamma)^{23}\text{Al}(p, \gamma)^{24}\text{Si}$ reaction sequence and the $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$ process path [14] (assuming ignition conditions of Ref. [6] and total accreted mass fractions consistent

with the “zM” model of Ref. [7]). In particular, in defining the temperature at which the $^{22}\text{Mg}(\alpha, p)$ reaction governs 50% of the nucleosynthetic flow in Type-I X-ray bursts as the “tipping” point between the rp - and (α, p) processes, we find that the latter will only become significant at temperatures $\gtrsim 0.85$ GK, as shown in Fig. 3. Such temperatures are only briefly reached for standard X-ray burst model calculations [6,56] and, in ruling out the previously possible lower-temperature onset of the (α, p) process [14,20], we may now conclude that the pathway through the $^{22}\text{Mg}(\alpha, p)$ reaction is not relevant for anything but the most energetic bursters.

In summary, we have performed the first measurement of the $^{23}\text{Ne}(d, p)^{24}\text{Ne}$ transfer reaction. Several strong single-particle states in ^{24}Ne have been identified and their associated neutron spectroscopic factors extracted to a precision of $\sim 20\%$. Using these spectroscopic factors to deduce the properties of resonant states in the astrophysical $^{23}\text{Al}(p, \gamma)$ reaction, we have reduced uncertainties in the strength of the key $E_r = 157$ keV, $\ell = 0$ level, in comparison with the most recent study of Ref. [20], by a factor of ~ 4 , considerably constraining the rate over the temperature range of X-ray bursts. In particular, we find that the $^{23}\text{Al}(p, \gamma)^{24}\text{Si}$ reaction is effective in bypassing the ^{22}Mg waiting point in the rp process (according to standard modelling conditions) for temperatures up to at least 0.85 GK, while the $^{22}\text{Mg}(\alpha, p)$ pathway might play a more prevalent role above 1 GK, the very peak temperature region only rarely reached in X-ray bursts. Further constraints on the $^{23}\text{Al}(p, \gamma)$ reaction would now require a precise determination of the reaction Q-value [21] and, in this regard, we understand that a new measurement of the ^{24}Si mass was recently performed at the National Superconducting Cyclotron Laboratory, USA [57]. The results for resonance strengths, combined with a precise Q-value determination, are now likely to constrain the uncertainties in the nuclear physics data sufficiently tightly to allow the accurate extraction of neutron star mass-radius ratios from current experimental observations of Type-I X-ray bursts [12].

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

The authors would like to thank the TRIUMF beam delivery group for their efforts in providing high-quality beams. This work has been supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), The Canada Foundation for Innovation and the British Columbia Knowledge Development Fund. TRIUMF receives federal funding via a contribution agreement through the National Research Council Canada. UK personnel were supported by the Science and Technology Facilities Council (STFC), JH acknowledges support at the University of Surrey under UKRI Future Leaders Fellowship MR/T022264/1, MR acknowledges support from the U.S. Department of Energy Office of Science through grant No. DE-SC0016988 and ZM acknowledges support from the

U.S. Department of Energy under grants DE-FG02-88ER40387 and DESC001904. We also benefited from support by the National Science Foundation under grant PHY-1430152 (Joint Institute for Nuclear Astrophysics-Center for the Evolution of the Elements).

References

- [1] R.K. Wallace, S.E. Woosley, *Astrophys. J.* 45 (1981) 389.
- [2] H. Schatz, K.E. Rehm, *Nucl. Phys. A* 777 (2006) 601.
- [3] H. Schatz, et al., *Phys. Rev. Lett.* 86 (2001) 3471.
- [4] G.R. Caughlan, W.A. Fowler, *Nat. Phys. Sci.* 238 (1972) 23.
- [5] J. Audouze, J.W. Truran, B.A. Zimmerman, *Astrophys. J.* 184 (1974) 493.
- [6] J.L. Fisker, H. Schatz, F-K. Thielemann, *Astrophys. J. Suppl. Ser.* 174 (2008) 261.
- [7] S.E. Woosley, et al., *Astrophys. J. Suppl. Ser.* 151 (2004) 75.
- [8] B. Paxton, et al., *Astrophys. J. Suppl. Ser.* 220 (2015) 15.
- [9] J. José, F. Moreno, A. Parikh, C. Iliadis, *Astrophys. J. Suppl. Ser.* 189 (2010) 204.
- [10] A. Parikh, J. José, F. Moreno, C. Iliadis, *Astrophys. J. Suppl. Ser.* 178 (2008) 110.
- [11] R.H. Cyburt, et al., *Astrophys. J.* 830 (2016) 2.
- [12] Z. Meisel, G. Merz, S. Medvid, *Astrophys. J.* 872 (2019) 84.
- [13] Z. Meisel, *Astrophys. J.* 860 (2018) 147.
- [14] J.S. Randhawa, et al., *Phys. Rev. Lett.* 125 (2020) 202701.
- [15] D.K. Galloway, M.P. Muno, J.M. Hartman, D. Psaltis, D. Chakrabarty, *Astrophys. J.* 179 (2008) 360.
- [16] D.K. Galloway, et al., *Astrophys. J. Suppl. Ser.* 249 (2020) 32.
- [17] H. Schatz, et al., *Phys. Rev. Lett.* 79 (1997) 3845.
- [18] K. Yoneda, et al., *Phys. Rev. C* 74 (2006) 021303(R).
- [19] A. Banu, et al., *Phys. Rev. C* 86 (2012) 015806.
- [20] C. Wolf, et al., *Phys. Rev. Lett.* 122 (2019) 232701.
- [21] M. Wang, et al., *Chin. Phys. C* 41 (2017) 3.
- [22] G. Lotay, et al., *Eur. Phys. J. A* 56 (2020) 3.
- [23] C. Iliadis, P.M. Endt, N. Prantzos, W.J. Thompson, *Astrophys. J.* 524 (1999) 434.
- [24] S.D. Pain, et al., *Phys. Rev. Lett.* 114 (2015) 212501.
- [25] V. Margerin, et al., *Phys. Rev. Lett.* 115 (2015) 062701.
- [26] G. Hackman, C.E. Svensson, *Hyperfine Interact.* 225 (2014) 241.
- [27] C.Aa. Diget, et al., *J. Instrum.* 6 (P02005) (2011).
- [28] G.L. Wilson, et al., *Phys. Lett. B* 759 (2016) 417.
- [29] A. Matta, et al., *Phys. Rev. C* 99 (2019) 044320.
- [30] S. Cruz, et al., *Phys. Lett. B* 786 (2018) 94.
- [31] B.A. Brown, W.A. Richter, *Phys. Rev. C* 74 (2006) 034315.
- [32] C.R. Hoffman, et al., *Phys. Rev. C* 68 (2003) 034304.
- [33] H. Herndl, J. Görres, M. Wiescher, B.A. Brown, L. Van Wormer, *Phys. Rev. C* 52 (1995) 2.
- [34] B.A. Brown, W.D.M. Rae, *Nucl. Data Sheets* 120 (2014) 115.
- [35] J.A. Tostevin, University of Surrey version of the code TWOFNR (of M. Toyama, M. Igarashi and N. Kishida) and code FRONT (private communication), <http://www.nucleartheory.net/NPG/code.htm>.
- [36] R.C. Johnson, P.J.R. Soper, *Phys. Rev. C* 1 (1970) 976.
- [37] J. Lee, et al., *Phys. Rev. C* 75 (2007) 064320.
- [38] A.J. Koning, J.P. Delaroche, *Nucl. Phys. A* 713 (2003) 231.
- [39] B. Longfellow, et al., *Phys. Rev. C* 101 (2020) 031303(R).
- [40] N.K. Timofeyuk, R.C. Johnson, A.M. Mukhamedzhanov, *Phys. Rev. Lett.* 91 (2003) 232501.
- [41] N.K. Timofeyuk, P. Descouvemont, R.C. Johnson, *Eur. Phys. J. A* 27 (2006) 269.
- [42] F. Ajzenberg-Selove, *Nucl. Phys. A* 375 (1982) 1.
- [43] M.D. Cooper, W.F. Hornyak, P.G. Roos, *Nucl. Phys. A* 218 (1974) 249.
- [44] A. Terakawa, et al., *Phys. Rev. C* 48 (1993) 2775.
- [45] A.M. Mukhamedzhanov, et al., *Phys. Rev. C* 73 (2006) 035806.
- [46] A.J. Howard, J.O. Pronko, C.A. Whitten Jr., *Nucl. Phys. A* 152 (1970) 317.
- [47] A.R. Stanford, P.A. Quin, *Nucl. Phys. A* 342 (1980) 283.
- [48] F. Entezami, A.K. Basak, O. Karban, P.M. Lewis, S. Roman, *Nucl. Phys. A* 366 (1981) 1.
- [49] G.P.A. Berg, R. Das, S.K. Datta, P.A. Quin, *Nucl. Phys. A* 289 (1977) 15.
- [50] W.W. Dykoski, D. Dehnhard, *Phys. Rev. C* 13 (1976) 80.
- [51] R.J. Peterson, C.A. Fields, R.S. Raymond, J.R. Thieke, J.L. Ullman, *Nucl. Phys. A* 408 (1983) 221.
- [52] M.C. Mermaz, et al., *Phys. Rev. C* 4 (1971) 1778.
- [53] R.A. Morrison, *Nucl. Phys. A* 140 (1970) 97.
- [54] R.L. Kozub, *Phys. Rev. C* 5 (1972) 413.
- [55] H. Schatz, et al., *Phys. Rep.* 294 (1998) 167.
- [56] G. Merz, Z. Meisel, *Mon. Not. R. Astron. Soc.* 500 (2021) 2958.
- [57] D. Puentes, et al., private communication.