



B_k spin vertex models and quantum algebras

Doron Gepner

Department of Particle Physics and Astrophysics, Weizmann Institute, Rehovot 76100, Israel

Received 6 May 2020; received in revised form 3 July 2020; accepted 14 July 2020

Available online 21 July 2020

Editor: Hubert Saleur

Abstract

We construct new solvable vertex models based on the spin representation of the Lie algebra B_k . We use these models to study the algebraic structure underlying such vertex theories. We show that all the B_k spin vertex models obey a version of the BMW algebra along with extra relations that are called n -CB (conformal braiding) algebras. These algebras were discussed before for various IRF (interaction round the face) models. Here we establish that the same algebras hold for vertex models.

© 2020 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Solvable lattice models in two dimensions are an excellent playing ground to study phase transitions, integrable models and knot theory. For reviews see [1,2].

Of particular significance to us is the algebraic structure underlying solvable lattice models. Examples of such algebras are the Temperley–Lieb algebra [3] and the Birman–Murakami–Wenzl algebra (BMW) [4,5]. These algebras had a major role in the solution of the models as well as applications such as knot theory. In particular, in knot theory we mention the celebrated work of Jones [6] and the works of Wadati et al. reviewed in ref. [2]. See also the book [7].

In recent works [8–11], the algebraic structure of IRF solvable lattice models was established. These works were based on the Yang–Baxter equation and the ansatz for Baxterization put forward in ref. [12], generalizing the two blocks Baxterization of Jones [6], to more than two blocks. An algebra was described for any number of blocks and called the n -CB algebra (conformal

E-mail address: doron.gepner@weizmann.ac.il.

braiding), where n is the number of blocks (defined as the order of the polynomial equation satisfied by the Boltzmann weights). The n -CB algebra includes the Temperley–Lieb algebra and a version of BMW algebra along with additional relations.

Our aim here is to study the n -CB algebra for vertex models. For recent works on vertex models, see [13–16]. We establish that the same algebra is obeyed by vertex models. For this study, we construct new vertex models based on the spin representation of the simple Lie algebra $SO(2k + 1)$ which is denoted as B_k , for arbitrary positive integer k . These models are described for any number of blocks which is $n = k + 1$. Previously, a Yang–Baxter solution was described for the vector representation of B_k (for a review see [2] and refs. therein.)

2. Vertex models and their Baxterization

Vertex lattice models are described by an element of $\text{End}(V \otimes V)$ where V is some vector space. It will be convenient to describe these elements using a matrix notation. Namely, if $R \in \text{End}(V \otimes V)$ then we may write,

$$R(v_\mu \otimes v_\nu) = R_{\mu,\nu}^{\bar{\mu},\bar{\nu}}(v_{\bar{\mu}} \otimes v_{\bar{\nu}}), \quad (2.1)$$

where ν and μ are basis vectors of the vector space V . Here, we include the indices of R .

The matrix R depends on the spectral parameter $R(u)$ where u is some complex number. The solvability of the model is encapsulated in the Yang–Baxter equation (YBE) which can be written as an equation in $\text{End}(V \otimes V \otimes V)$,

$$(R(u) \otimes 1)(1 \otimes R(u + v))(R(v) \otimes 1) = (1 \otimes R(v))(R(u + v) \otimes 1)(1 \otimes R(u)). \quad (2.2)$$

This equation can be expanded in terms of matrix elements, eq. (2.1), to give,

$$\sum_{\alpha,\beta,\gamma} R_{j,k}^{\beta,\alpha}(u) R_{i,\beta}^{l,\gamma}(u + v) R_{\gamma,\alpha}^{m,n}(v) = \sum_{\alpha,\beta,\gamma} R_{i,j}^{\alpha,\beta}(v) R_{\beta,k}^{\gamma,n}(u + v) R_{\alpha,\gamma}^{l,m}(u). \quad (2.3)$$

We assume that $R(u)$ is a trigonometric solution of the Yang–Baxter equation.

The vertex models may obey a number of properties in addition to the YBE. The initial condition,

$$R_{i,j}^{k,l}(0) = \delta_{ik} \delta_{jl}. \quad (2.4)$$

The inversion relation,

$$\sum_{m,n} R_{i,j}^{m,n}(u) R_{m,n}^{l,k}(-u) = \rho(u) \rho(-u) \delta_{il} \delta_{jk}, \quad (2.5)$$

where $\rho(u)$ is a function, to be specified later. Also, crossing symmetry,

$$R_{ji}^{k,l}(u) = R_{k,j}^{l,\bar{i}}(\lambda - u) \left[\frac{r(i)r(l)}{r(j)r(k)} \right]^{1/2}, \quad (2.6)$$

where λ is the crossing parameter. Here \bar{i} is the charge conjugation of i and the crossing multiplier is $r(i)$, where $r(\bar{i}) = 1/r(i)$. Usually, in a vector model, we will have that $\bar{v} = -v$. We have the reflection symmetry,

$$R_{i,j}^{m,n}(u) = R_{m,n}^{i,j}(u). \quad (2.7)$$

Finally, we have the charge conservation,

$$R_{m,n}^{i,j} = 0, \quad \text{unless } m + n = i + j. \tag{2.8}$$

We find it convenient to define an operator form for the R matrix. We define the matrix,

$$X_i(u) = \sum_{m,n,a,b} R_{m,n}^{a,b}(u) I^{(1)} \otimes \dots \otimes I^{(i-1)} \otimes e_{a,m}^{(i)} \otimes e_{b,n}^{(i+1)} \otimes I^{(i+2)} \otimes \dots \otimes I^{(n)}, \tag{2.9}$$

where \otimes means tensor product, $I^{(i)}$ is the identity matrix at position i , and e_{ab} is a matrix whose elements are given by $(e_{rs})_{lm} = \delta_{rl}\delta_{sm}$. We define in a similar fashion other operators. It is then clear that the YBE, eq. (2.2), can be written as

$$\begin{aligned} X_i(u)X_j(v) &= X_j(v)X_i(u), \quad \text{if } |i - j| \geq 2, \\ X_i(u)X_{i+1}(u+v)X_i(v) &= X_{i+1}(v)X_i(u+v)X_{i+1}(u). \end{aligned} \tag{2.10}$$

We will build the vertex model from the data of a fixed conformal field theory. Given the conformal field theory \mathcal{O} , let V be the representation of some primary field in \mathcal{O} . The vertex model is given in terms of the representations that appear in the tensor product of V . We assume that the theory is real and that $[V]$ is a real representation. Thus, we have the fusion product,

$$[V] \times [V] = \sum_{i=0}^{n-1} [\psi_i], \tag{2.11}$$

where $[x]$ denotes the primary field x . Here n is an integer which is called the number of blocks and $[\psi_0] = 1$, is the unit representation. The order of the fields in eq. (2.11) is important as will be discussed later. For real models a rule of thumb appears for this order. Namely, the field ψ_{i+1} appears in the tensor product of ψ_i and the adjoint representation (for quantum groups). Thus, in particular, $\psi_0 = 1$ and ψ_1 is the adjoint representation. The complete implementation of this rule is presently not clear. For each representation that appears in this fusion product we define a projection operator P_i onto this representation. For this purpose, we define the limit of the trigonometric solution of the Yang–Baxter equation, $X_i(u)$, as,

$$X_i = \lim_{u \rightarrow i\infty} e^{i(n-1)u} X_i(u), \quad X_i^t = \lim_{u \rightarrow -i\infty} e^{-i(n-1)u} X_i(u). \tag{2.12}$$

The eigenvalues of X_i can be seen from conformal field theory to be,

$$\lambda_i = \epsilon_i e^{i\pi(2\Delta_v - \Delta_i)}, \tag{2.13}$$

where Δ_v is the conformal dimension of the primary field $[V]$, Δ_i is the conformal dimension of the representation $[\psi_i]$ and $\epsilon_i = \pm 1$ indicating whether the product is symmetric or antisymmetric.

From X_i we can define the a th projection operator as,

$$P_i^a = \prod_{p \neq a} \left[\frac{X_i - \lambda_p I}{\lambda_a - \lambda_p} \right], \tag{2.14}$$

where $a = 0, 1, \dots, n - 1$ and I is the unit operator. We have the following relations for the projection operators,

$$X_i = \sum_{a=0}^{n-1} \lambda_a P_i^a, \tag{2.15}$$

$$\sum_{a=0}^{n-1} P_i^a = I, \quad P_i^a P_i^b = \delta_{ab} P_i^a. \quad (2.16)$$

From the projection operator one may build the solution to the YBE, $X_i(u)$. It is basically the same conjecture as for the IRF models described in ref. [12]. We define the parameters by,

$$\zeta_i = \pi(\Delta_{i+1} - \Delta_i)/2, \quad (2.17)$$

and $\lambda = \zeta_0$ is the crossing parameter and $i = 0, 1, \dots, n-2$. The trigonometric solution to the Yang–Baxter equation ansatz is then,

$$X_i(u) = \sum_{a=0}^{n-1} f_a(u) P_i^a, \quad (2.18)$$

where the functions $f_a(u)$ are given by,

$$f_a(u) = \left[\prod_{r=1}^a \sin(\zeta_{r-1} - u) \right] \left[\prod_{r=a+1}^{n-1} \sin(\zeta_{r-1} + u) \right] / \left[\prod_{r=1}^{n-1} \sin(\zeta_{r-1}) \right]. \quad (2.19)$$

From the ansatz it is easy to see that the inversion relation, eq. (2.5), holds with

$$\rho(u) = \prod_{r=1}^{n-1} \frac{\sin(\zeta_{r-1} - u)}{\sin(\zeta_{r-1})}. \quad (2.20)$$

The crossing equation, eq. (2.6), holds with the crossing parameter $\lambda = \zeta_0$.

The order of the fields $[\psi_i]$ is important and the YBE holds only for one particular order. We will specify below the order which is suitable for specific models.

An interesting question is the relation between the CFT used to define the model and the conformal field theories which arise at the criticality of the models. A partial answer, based on (D. Gepner, unpublished work), is that the critical field theories are cosets of the original theory, where in one limit, the original theory is in the denominator, whereas in the other it is in the numerator of the coset theory. The full coset theory is presently unknown, except in some examples. For a review see ref. [2].

We are interested in the algebra obeyed by these models. For this purpose, we define the operators,

$$G_i = 2^{n-1} e^{-i(n-1)\zeta_0/2} \left[\prod_{r=1}^{n-1} \sin(\zeta_{r-1}) \right] X_i, \quad (2.21)$$

and

$$G_i^{-1} = 2^{n-1} e^{i(n-1)\zeta_0/2} \left[\prod_{r=1}^{n-1} \sin(\zeta_{r-1}) \right] X_i^t, \quad (2.22)$$

where X_i and X_i^t are given by eq. (2.12). We also define the operator,

$$E_i = X_i(\lambda). \quad (2.23)$$

The inversion relation eqs. (2.5), (2.20) implies that as defined $G_i G_i^{-1} = I$.

From the crossing relation, eq. (2.6), it follows that E_i can be expressed as follows,

$$E_{a,b}^{m,n} = \delta_{\bar{a},b} \delta_{\bar{m},n} r(n)r(b), \tag{2.24}$$

where $r(a)$ is the crossing multiplier. Here we reverted back to the explicit notation for E_i . From the above equation, it follows that E_i obeys the relation,

$$E_i E_{i\pm 1} E_i = E_i, \tag{2.25}$$

where we used the equation $r(\bar{m}) = 1/r(m)$. From the ansatz eqs. (2.18), (2.19) it follows that

$$E_i^2 = b E_i, \tag{2.26}$$

where

$$b = \prod_{r=1}^{n-1} \frac{\sin(\zeta_0 + \zeta_{r-1})}{\sin(\zeta_{r-1})}. \tag{2.27}$$

These two equations together are the celebrated Temperley–Lieb algebra [3]. Thus, we proved that any real vertex model, with any number of blocks, obeys the Temperley–Lieb algebra, assuming that the ansatz eq. (2.18)-(2.19) holds.

Since E_i is proportional to P_i^0 we find the following relations,

$$G_i E_i = E_i G_i = l^{-1} E_i, \tag{2.28}$$

where l is given by,

$$l = i^{n-1} \exp \left[i(n-1)\zeta_0/2 + i \sum_{r=0}^{n-2} \zeta_r \right]. \tag{2.29}$$

From the YBE, eq. (2.2), we find that G_i obeys the braid group relation,

$$G_i G_j = G_j G_i \quad \text{if } |i-j| \geq 2, \quad G_i G_{i+1} G_i = G_{i+1} G_i G_{i+1}. \tag{2.30}$$

From the ansatz, eqs. (2.18), (2.19), and from the equation $\sum_a P_i^a = I$ we find the skein relation,

$$G_i^{n-2} = a E_i + \sum_{r=-1}^{n-3} b_r G_i^r, \tag{2.31}$$

where the coefficients a and b_r are expressed as functions of the parameters ζ_i , which can be calculated from the ansatz, eqs. (2.18), (2.19).

3. Vertex models and quantum groups

We utilize now the powerful method for constructing solutions to the YBE vertex models, eq. (2.2), which is quantum groups [17–19].

The definition of the quantum group is as follows [17,18]. Let $A = (a_{ij})$ be a Cartan matrix of a simple Lie algebra G . Let $\{\alpha_j\}$ and $\{h_j\}$ be the simple roots and coroots, for $1 \leq i \leq N$, such that $\langle h_i | \alpha_j \rangle = a_{ij}$. For a parameter q which is nonzero we define $q_i = q^{(\alpha_i, \alpha_i)}$, where (\cdot) is the invariant inner product in \mathfrak{h}^* .

The generators of the quantum group are $\{k_i^{\pm 1}, e_i, f_i\}_{1 \leq i \leq N}$. They obey the relations,

$$k_i k_i^{-1} = k_i^{-1} k_i = 1, \quad [k_i, k_j] = 0, \tag{3.1}$$

$$k_i e_j k_i^{-1} = q_i^{a_{ij}} e_j, \quad k_i f_j k_i^{-1} = q_i^{-a_{ij}} f_j, \tag{3.2}$$

$$[e_i, f_j] = \delta_{ij} (k_i^2 - k_i^{-2}) / (q_i^2 - q_i^{-2}). \tag{3.3}$$

There are additional relations, (3D) and (3E) of ref. [17], but we will not require these.

For $q \rightarrow 1$ the quantum algebra, denoted by $U_q(G)$ reduces to the simple Lie algebra G . (Actually, the quantum group can be defined for any Kac–Moody algebra [17,18].)

We shall need the co-product of the quantum group $U_q(G)$. This is the homomorphism $\Delta^{(m)}U \rightarrow \otimes^m U$ (m fold tensor product), defined by

$$\Delta^{(m)}(k_i) = k_i \otimes k_i \otimes \dots \otimes k_i, \tag{3.4}$$

$$\Delta^{(m)}(X_i) = \sum_{\nu=1}^m k_i \otimes \dots \otimes k_i \otimes^{\nu} X_i \otimes k_i^{-1} \otimes \dots \otimes k_i^{-1}, \tag{3.5}$$

for $X_i = e_i$ or $X_i = f_i$. The co-product obey the same quantum group $U_q(G)$.

In the following we will assume that q is not a root of unity, unless otherwise specified. In this case, the irreducible representations of $U_q(G)$ are labeled by the irreducible representations of G and have the same dimensions.

The solution to the vertex YBE, eq. (2.2), commutes with the co-product,

$$[R, \Delta^{(2)}(X_i)] = 0, \tag{3.6}$$

for any $X_i = e_i$ or f_i or k_i . This equation is not enough to determine the R matrix. However, given a solution to this equation, it is guaranteed to have the same eigenvectors but not the same eigenvalues as the R matrix. Thus, they share the same projection operators, eq. (2.14). Assuming that the number of distinct eigenvalues of R is n , where n is the number of blocks, than the projection operators are given as in eq. (2.14),

$$P^a = \prod_{p \neq a} \left[\frac{R - \lambda_p I}{\lambda_a - \lambda_p} \right], \tag{3.7}$$

where λ_p are the eigenvalues of R . We can then use our ansatz eqs. (2.18), (2.19) to get the full trigonometric solution of the YBE.

Each projection operator P^a is associated to some representation in the tensor product $g \in V \times V$, where V is the representation used to define the vertex model. The projection operator can be written as,

$$(P^g)_{a,b}^{c,d} = \sum_{\lambda} \langle g \lambda | V a V b \rangle \langle g \lambda | V c V d \rangle, \tag{3.8}$$

where λ runs over the weights of the representation g and $\langle g \lambda | V a V b \rangle$ is the quantum group Wigner coefficient of this tensor product. P^g is the vertex projection operator with the weights a, b, c, d which are weights of the representation V . From this equation, it is clear that the projection operator vanishes unless $a + b = c + d$, eq. (2.8). For $SU(2)$ this was described in ref. [19]. In this reference, it was shown that for $SU(2)$ the vertex and the IRF models have the same Baxterization.

4. B_k spin vertex models

Our purpose is to describe solvable vertex models based on the algebra B_k , or $SO(2k + 1)$, where the representation V is the spinor representation. We use the basis for B_k where the simple roots are $\alpha_n = \epsilon_n - \epsilon_{n+1}$, for $n = 1, 2, \dots, k - 1$ and $\alpha_k = \epsilon_k$. Here ϵ_i are orthogonal unit vectors. The spinor representation has the highest weight $(\epsilon_1 + \epsilon_2 + \dots + \epsilon_k)/2$ and the weights of this representation are $(\pm\epsilon_1 \pm \epsilon_2 \pm \dots \pm \epsilon_k)/2$. We find it convenient to add $1/2$ to these weights and to represent the weights of the spinor representation by m where $m_i = 0$ or 1 .

We look for a solution C for the spinor representation of the algebra B_k , which commutes with the co-product, eq. (3.6). Such a solution was described recently in a paper by Wenzl [20]. The solution C is an element of $\text{End}(V \otimes V)$ where V denotes the spinor representation. It is given by [20],

$$C_{m,n}^{b,c} = \sum_{j=1}^k \delta_{m_j, 1-n_j} (-q^2)^{\{m-n\}_j} \delta_{b, \bar{m}^j} \delta_{c, \bar{n}^j} + (-1)^k \delta_{m,b} \delta_{n,c} (-q^2)^{\{m-n\}_k} / [2], \tag{4.1}$$

where

$$\{m\}_j = \sum_{r=1}^j m_r, \tag{4.2}$$

and \bar{n}^j is equal to n except at the j th coordinate where it is $1 - n_j$. Here $[2] = q + q^{-1}$. Here $m, n, b, c = 0$ or 1 are weights of the spinor representation shifted by $1/2$. The matrix C , so constructed, commutes with the co-product eq. (3.6).

The eigenvalues of the matrix C were computed by Wenzl [20]. They are given by

$$\lambda_j = (-1)^j s(k + \frac{1}{2} - j), \quad \text{for } j = 0, 1, \dots, k, \tag{4.3}$$

where

$$s(x) = \frac{q^{2x} - q^{-2x}}{q^2 - q^{-2}}. \tag{4.4}$$

There are $k + 1$ distinct eigenvalues of C . Thus, this is a $k + 1$ blocks theory. Each eigenvalue corresponds to a representation in the tensor product $V \times V$, where V is the spinor representation. The j th eigenvalue λ_j corresponds to the representation $V_j = \wedge^j v$ where v is the vector representation. The highest weight of the representation V_j is $\epsilon_1 + \epsilon_2 + \dots + \epsilon_j$. It is the fully anti-symmetric representation in the tensor of j vector representations.

We assume that q is not a root of unity and is nonzero. To connect with section 3, we identify

$$q^2 = \exp[\pi i / (r + g)], \tag{4.5}$$

where r is the level of the WZW model based on B_k , at level r and g is the dual Coxeter number,

$$g = 2k - 1. \tag{4.6}$$

We assume that the level r is not a real rational number, so that q is not a root of unity. The dimension of the representation with highest weight Λ in a WZW theory is given by

$$\Delta_\Lambda = \frac{\Lambda(\Lambda + 2\rho)}{2(r + g)}. \tag{4.7}$$

Here ρ is half the sum of positive roots and $C_\Lambda = \Lambda(\Lambda + 2\rho)$ is the Casimir of the representation Λ . See, e.g. [21].

As explained in section 3, the eigenvectors of C are the projections of the solution of the YBE to the different representations. We thus define,

$$(P^a)_{m,n}^{b,c} = \prod_{p \neq a} \left[\frac{C - \lambda_p I}{\lambda_a - \lambda_p} \right], \tag{4.8}$$

where the product is in $\text{End}(V \otimes V)$ and I is the identity map.

We know from equation (2.13) that the eigenvalues of the R matrix are given by $\epsilon_j \exp[-i\pi \Delta_j]$ up to an irrelevant constant. Thus, we need to compute the second Casimir of the representations V_j , since the dimensions of the representations are computed in terms of the Casimir, eq. (4.7). The Casimir is given by

$$C(V_j) = C_j = j(2k + 1 - j). \tag{4.9}$$

Thus the eigenvalues of R are

$$\beta_j = \epsilon_j q^{-C(V_j)}, \tag{4.10}$$

where ϵ_j is a sign which is harder to compute. To give this sign we define,

$$(h_0, h_1, \dots, h_k) = (0, 2, 4, \dots, k, k - 1, k - 3, \dots, 1), \tag{4.11}$$

for even k . For odd k ,

$$(h_0, h_1, \dots, h_k) = (0, 2, \dots, k - 1, k, k - 2, k - 4, \dots, 1). \tag{4.12}$$

Then the sign ϵ_j is given by,

$$\epsilon_{h_s} = (-1)^s. \tag{4.13}$$

We are now in position to construct the R matrix as

$$R_{m,n}^{a,b} = \sum_{j=0}^k \beta_j (P^j)_{m,n}^{a,b}. \tag{4.14}$$

This is since we know the projection operators from eq. (4.8) and the eigenvalues of R from eq. (4.10).

We can now check that the R matrix, so constructed, obeys the braiding relation,

$$\sum_{\alpha, \beta, \gamma} R_{j,k}^{\beta, \alpha} R_{i,\beta}^{l, \gamma} R_{\gamma, \alpha}^{m, n} = \sum_{\alpha, \beta, \gamma} R_{i,j}^{\alpha, \beta} R_{\beta, k}^{\gamma, n} R_{\alpha, \gamma}^{l, m}. \tag{4.15}$$

We checked this R matrix numerically for $k = 2, 3, 4, 5, 6$ and it is, indeed, obeyed for various weights and for general q .

We can now build the full solution to the YBE, eq. (2.3). We need to compute the parameters ζ_i . To do this, we need to know the order of the operators ψ_i in eq. (2.11). In fact, the order of the representations is given by h_r . Thus, we have

$$\zeta_j = (C_{h_{j+1}} - C_{h_j})/2, \tag{4.16}$$

for $j = 0, 1, \dots, k - 1$. We also replace the $\sin(x)$ in eq. (2.19) by

$$\sin(x) \rightarrow p(x) = q^x - q^{-x}. \tag{4.17}$$

Then the solution to the YBE, eq. (2.3), assumes the form,

$$X_{m,n}^{a,b}(u) = \sum_{j=0}^k f_j(u) (P^{h_j})_{m,n}^{a,b}, \tag{4.18}$$

where

$$f_a(u) = \left[\prod_{j=1}^a p(\zeta_{j-1} - u) \right] \left[\prod_{j=a+1}^k p(\zeta_{j-1} + u) \right] / \left[\prod_{j=1}^k p(\zeta_{j-1}) \right], \tag{4.19}$$

where $a = 0, 1, \dots, k$.

For example for $k = 6$ we have, $(\zeta_0, \zeta_1, \dots, \zeta_5) = (11, 7, 3, -1, -5, -9)$. The crossing parameter is always $\lambda = \zeta_0$.

We can now check numerically that the matrix $X_i(u)$ so defined obeys the Yang–Baxter equation, eq. (2.3). We checked this numerically for $k = 2, 3, 4, 5, 6$ for various values of the weights and the spectral parameters and indeed the YBE is obeyed for general q .

Actually, our results hold also for q which is a root of unity. We take $q^2 = \exp[i\pi s/(r + g)]$, as in eq. (4.5), where r and s are now integers such that, $\gcd(s, r + g) = 1$. Then, if the level r is greater or equal two, then the fusion rule in eq. (2.11) is the same as the tensor product, since the representations ψ_i appear at level two. Namely, all the representations appear if the level is greater or equal two. Thus, the ansatz eqs. (4.18), (4.19), holds as it is. We checked this for various algebras of the type B_k and various integer levels, r , and indeed the YBE is obeyed for q which is a root of unity, as well. Thus, for levels greater than one, exactly the same solution holds. We call these models for rational level, the restricted models.

5. n -CB algebra and B_k vertex models

The B_k vertex models are $k + 1$ blocks models. For $k = 2$ this is a three blocks model. Thus, it is natural that the model would obey the BMW algebra [4,5], as we will show. We use the operator notation eq. (2.9) and define the operators G_i and E_i as before, eqs. (2.21)-(2.23). The relations of the BMW algebra are,

$$G_i - G_i^{-1} = m(1 - E_i), \tag{5.1}$$

$$G_i G_j = G_j G_i \text{ if } |i - j| \geq 2, \quad G_i G_{i+1} G_i = G_{i+1} G_i G_{i+1}, \tag{5.2}$$

$$E_i E_{i\pm 1} E_i = E_i, \quad E_i^2 = b E_i, \tag{5.3}$$

$$G_{i\pm 1} G_i E_{i\pm 1} = E_i G_{i\pm 1} G_i = E_i E_{i\pm 1}, \quad G_{i\pm 1} E_i G_{i\pm 1} = G_i^{-1} E_{i\pm 1} G_i^{-1}, \tag{5.4}$$

$$G_{i\pm 1} E_i E_{i\pm 1} = G_i^{-1} E_{i\pm 1}, \quad E_{i\pm 1} E_i G_{i\pm 1} = E_{i\pm 1} G_i^{-1}, \tag{5.5}$$

$$G_i E_i = E_i G_i = l^{-1} E_i, \tag{5.6}$$

$$E_i G_{i\pm 1} E_i = l E_i, \quad E_i G_{i\pm 1}^{-1} E_i = l^{-1} E_i, \tag{5.7}$$

where

$$b = m^{-1}(l - l^{-1}) + 1, \tag{5.8}$$

and l and b are given by eqs. (2.27), (2.29) for three blocks, $n = 3$.

We checked the BMW relations eqs. (5.1)–(5.8) for the B_2 vertex model and indeed they are all obeyed for various weights and general q . We find,

$$b = -(q^4 + q^2 + q^{-2} + q^{-4}), \quad (5.9)$$

and

$$l = -q^5, \quad m = q + q^{-1} \quad (5.10)$$

In fact, as we show, the BMW algebra is also obeyed for $k > 2$, except for the skein relation, eq. (2.31). The relations eqs. (5.2), (5.3), (5.6) were already proved in section 2 for all the vertex models, eqs. (2.25), (2.26), (2.28), (2.30), along with the new skein relation eq. (2.31). It remains to check the other relations. We checked them for $k = 3, 4, 5, 6$ with various weights, and general q , and indeed they are all obeyed. The parameters l and b are given by eqs. (2.27), (2.29). We call this algebra BMW' .

We checked the BMW' algebra also for the restricted models and it also holds. Our discussion below applies equally well to the restricted models as they also obey the same ansatz and the same YBE.

In ref. [8–11], we analyzed the Yang–Baxter equation assuming only BMW' algebra and the ansatz eqs. (2.18), (2.19). We established this only for IRF models and not for vertex models. However, all the assumptions are exactly the same, even though the definition of the operators is different, eq. (2.9). Thus, the same conclusions we found by expanding the YBE still hold. We found that for three blocks, $k = 2$, we get a weak version of the BMW algebra [10]. For four blocks, $n = 4$, we get an algebra which we called 4–CB, which is BMW' , along with one additional relation. The additional relations are enlisted in the appendix. For five blocks $n = 5$ we get additional 19 relations which are quite bulky. This method can be used to compute the algebra for any number of blocks, n , which we call n -CB algebra.

Since all of the assumptions are the same for IRF models as for the vertex models, we conclude that the n -CB algebra holds for the B_k vertex models, with $n = k + 1$. We conjecture that the n -CB algebra is obeyed for all the solvable vertex models with n blocks, for which the ansatz eqs. (2.18), (2.19) holds.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

It is my pleasure to thank Hans Wenzl for many discussions, encouragement and for sending me his paper [20]. I am also grateful to Ida Deichaite for remarks on the manuscript and valuable impetus. I also thank Jian-Rong Li for comments. I thank Vladimir Belavin for help with the calculations.

Appendix A. Four CB relations

For completeness, we summarize here the four blocks relations [10]. The skein relation is given by

$$G_i^2 = ie^{-\frac{1}{2}i\zeta_0 - i\zeta_1 - i\zeta_2} \left(1 - e^{2i\zeta_1} + e^{2i\zeta_1 + 2i\zeta_2} \right) G_i + ie^{-\frac{3}{2}i\zeta_0 + i\zeta_1 - i\zeta_2} G_i^{-1} + \frac{e^{-2i\zeta_0 - 2i\zeta_1 - 2i\zeta_2} (e^{2i\zeta_1} - 1) (1 + e^{2i\zeta_0 + 2i\zeta_1 + 2i\zeta_2}) (e^{2i\zeta_2} - 1)}{(e^{2i\zeta_0 + 2i\zeta_2} - 1)} E_i - e^{-i\zeta_0 - 2i\zeta_2} \left(1 - e^{2i\zeta_2} + e^{2i\zeta_1 + 2i\zeta_2} \right). \quad (A.1)$$

The last relation follows from the Yang Baxter equation and the ansatz eq. (2.18)-(2.19). It is

$$g(i, i + 1, i) = g(i + 1, i, i + 1), \quad (A.2)$$

where

$$g = a_{1,2,4} + a_{1,3,1} + a_{4,2,1} + iq^{-\zeta_0/2 + \zeta_1 - \zeta_2} (a_{1,3,4} + a_{4,2,4} + a_{4,3,1}) + iq^{\zeta_0/2 - \zeta_1 + \zeta_2} (a_{2,3,4} + a_{4,1,4} + a_{4,3,2}) + i \frac{q^{\zeta_1 + \zeta_2}}{(q^{2\zeta_1} - 1)(q^{2\zeta_2} - 1)} \left(q^{\zeta_0/2} a_{1,2,1} + q^{-\zeta_0/2} a_{2,1,2} \right) + za_{4,3,4}, \quad (A.3)$$

where

$$z = \frac{q^{-\zeta_0 - 2\zeta_1 - 2\zeta_2} (q^{2\zeta_1} - 1)(q^{2\zeta_2} - 1)}{q^{2\zeta_0 + 2\zeta_2 - 1}} \times \left(2q^{2\zeta_0 + 2\zeta_2} + 2q^{2\zeta_0 + 2\zeta_1 + 2\zeta_2} + q^{4\zeta_0 + 2\zeta_1 + 4\zeta_2} + 1 \right). \quad (A.4)$$

We denoted by $a_{i,j,k}(r, s, t)$ the element of the algebra $a_i[r]a_j[s]a_k[t]$ where $a_i[r]$ is G_r , G_r^{-1} , E_r or 1_r , if $i = 1, 2, 3, 4$, respectively. Here, $q = e^i$.

We checked these relations for the B_3 vertex model, which is a four blocks model, numerically, and indeed they hold for various values of the heights and for general values of q .

References

- [1] R.J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press, London, England, 1982.
- [2] M. Wadati, T. Deguchi, Y. Akutsu, *Phys. Rep.* 180 (4) (1989) 247.
- [3] N. Temperley, E. Lieb, *Proc. R. Soc. A* 322 (1971) 251.
- [4] J.S. Birman, H. Wenzl, *Trans. Am. Math. Soc.* 313 (1) (1989) 313.
- [5] J. Murakami, *Osaka J. Math.* 24 (4) (1987) 745.
- [6] V.F.R. Jones, *Int. J. Mod. Phys. A* 5 (1990) 441.
- [7] L.H. Kauffman, *Knots and Physics*, World Scientific, Singapore, 1991.
- [8] V. Belavin, D. Gepner, *Nucl. Phys. B* 938 (2019) 223.
- [9] V. Belavin, D. Gepner, *J. High Energy Phys.* 02 (2019) 033.
- [10] V. Belavin, D. Gepner, J.R. Li, R. Tessler, *J. High Energy Phys.* 11 (2019) 155.
- [11] V. Belavin, D. Gepner, arXiv:2001.09280, 2020.
- [12] D. Gepner, arXiv:hep-th/9211100v2, 1992.
- [13] J.H. Arbeitman, S. Mantilla, I. Sodeman, *Phys. Rev. B* 99 (2019) 245108.
- [14] A. Bossard, W. Galleas, *J. Math. Phys.* 60 (2019) 103509.
- [15] B. Brubaker, V. Buciumas, D. Bump, H. Gustafsson, arXiv:1902.01795v3, 2019.
- [16] K.S. Nirov, A.V. Razumov, *SIGMA* 15 (2019) 068.
- [17] M. Jimbo, *Lett. Math. Phys.* 10 (1985) 63.
- [18] V.G. Drinfeld, *Dokl. Akad. Nauk SSSR* 283 (5) (1985) 1060.
- [19] V. Pasquier, *Commun. Math. Phys.* 118 (1988) 355.
- [20] H. Wenzl, *Dualities for spin representation*, arXiv:2005.11299, 2020.
- [21] P. Francesco, P. Mathieu, D. Senechal, *Conformal Field Theory*, Springer, 1997.