



Excluding static and spherically symmetric black holes in Einsteinian cubic gravity with unsuppressed higher-order curvature terms

Antonio De Felice^a, Shinji Tsujikawa^{b,*}

^a Center for Gravitational Physics and Quantum Information, Yukawa Institute for Theoretical Physics, Kyoto University, 606-8502, Kyoto, Japan

^b Department of Physics, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo 169-8555, Japan

ARTICLE INFO

Article history:

Received 12 May 2023

Received in revised form 20 June 2023

Accepted 26 June 2023

Available online 29 June 2023

Editor: M. Trodden

ABSTRACT

Einsteinian cubic gravity is a higher-order gravitational theory in which the linearized field equations of motion match Einstein's equations on a maximally symmetric background. This theory allows the existence of a static and spherically symmetric black hole solution where the temporal and radial metric components are equivalent to each other ($f = h$), with a modified Schwarzschild geometry induced by cubic curvature terms. We study the linear stability of the static and spherically symmetric vacuum solutions against odd-parity perturbations without dealing with Einsteinian cubic gravity as an effective field theory where the cubic curvature terms are always suppressed relative to the Ricci scalar. Unlike General Relativity containing one dynamical perturbation, Einsteinian cubic gravity has three propagating degrees of freedom in the odd-parity sector. We show that at least one of those dynamical perturbations always behaves as a ghost mode. We also find that one dynamical degree of freedom has a negative sound speed squared $-1/2$ for the propagation of high angular momentum modes. Thus, the static and spherically symmetric hairy black hole solutions realized by unsuppressed cubic curvature terms relative to the Ricci scalar are excluded by ghost and Laplacian instabilities.

© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

General Relativity (GR) has been successful in describing the gravitational interaction between submillimeter and solar-system scales [1–5]. At extremely small distances close to the Planck length, however, it is expected that GR is replaced by a quantum theory of gravity with an ultraviolet completion. The attempt for the construction of a power-counting renormalizable gravitational theory was advocated by Stelle [6] by taking into account quadratic-order curvatures to the Einstein-Hilbert action. In string theory, the low-energy effective action also contains quadratic curvature corrections known as a Gauss-Bonnet (GB) term [7,8]. Moreover, higher-order curvature terms have played a prominent role in conformal field theory with holography [9–12].

In gravitational theories where the field equations of motion contain derivatives higher than second order in the metric tensor $g_{\mu\nu}$, the system can be unstable due to the emergence of Ostrogradski instability [13,14] associated with a Hamiltonian unbounded from below. To avoid such a problem, Lanczos [15] and Lovelock [16] constructed gravitational theories with second-order field equations of motion for general, curved backgrounds. Exploiting polynomial functions of the Riemann curvature tensors in four-dimensional spacetime endowed with four-dimensional diffeomorphism invariance, Lovelock showed that the symmetric and divergence-free tensors $A_{\mu\nu}$, which depend on $g_{\mu\nu}$ and its derivatives up to second order, are expressed by a linear combination of the Einstein tensor $G_{\mu\nu}$ and the metric tensor $g_{\mu\nu}$. In spacetime dimensions higher than four, there is the quadratic-order GB curvature scalar affecting the spacetime dynamics. In four dimensions, the GB term corresponds to an Euler density which does not contribute to the field equations of motion. To extract its effect in four dimensions, the GB term needs to be coupled to other degrees of freedom (DOFs) such as scalar or vector fields [17–25].

If we consider cubic-order curvature combinations constructed from the Riemann tensors, there is also an additional Euler density in the action corresponding to the surface integral in four dimensions [16]. According to the Lovelock theorem, there are no other nontrivial cubic-order terms keeping the field equations of motion up to second order in general, curved backgrounds. If we consider some specific

* Corresponding author.

E-mail address: tsujikawa@waseda.jp (S. Tsujikawa).

spacetime, however, it is possible to construct nontrivial cubic-order theory whose graviton spectrum shares a similar property to that in GR. On a maximally symmetric background, which includes the Minkowski and de Sitter spacetimes, there is a unique cubic combination \mathcal{P} which keeps the structure of linearized perturbation equations of motion in Einstein gravity [26]. In this theory, which is dubbed *Einsteinian cubic gravity* (ECG), the cubic curvature term can give rise to derivatives higher than second order in general, curved backgrounds, so the spacetime dynamics is generally different from that in GR [27].

On a static and spherically symmetric (SSS) background in a vacuum configuration, ECG admits the existence of a black hole (BH) solution whose temporal and radial metric components (denoted as f and h , respectively) coincide with each other [28–31] (see also Refs. [32–35]). Since f and h are affected by the cubic curvature term, the background geometry is still different from the Schwarzschild BH solution. One can also construct a cubic gravity theory by imposing the condition $f = h$ (up to a time reparametrization freedom in f) on the SSS background [36–38]. This construction allows the existence of the Lagrangian \mathcal{P} mentioned above as well as two additional cubic combinations \mathcal{C} and \mathcal{C}' defined in Ref. [36]. As we will see later in Sec. 2, both \mathcal{C} and \mathcal{C}' vanish for $f = h$ and hence only the Lagrangian \mathcal{P} contributes to the single differential equation for f ($= h$).

If we apply ECG to the cosmological dynamics on the Friedmann-Lemaître-Robertson-Walker (FLRW) spacetime, the field equations of motion arising from \mathcal{P} contain derivatives higher than second order. On the FLRW background, there is a unique combination $\mathcal{P} - 8\mathcal{C}$ that leads to the second-order Friedmann equation [39,40]. This theory—dubbed *cosmological Einsteinian cubic gravity* (CECG)—was applied to the dynamics of inflation and late-time cosmological epochs [39–45]. On an exact de Sitter background, tensor perturbations with two polarized modes propagate as in GR without pathological behavior. If one considers a spatially homogeneous Bianchi type I manifold close to the isotropic de Sitter spacetime, however, there are three dynamical propagating DOFs associated with linear perturbations in the odd-parity sector. In Ref. [46], it was shown that, in the regime of small anisotropies, such theory possesses at least one ghost mode as well as short-time-scale tachyonic instability. Hence CECG cannot be used to describe a viable geometric inflationary scenario.

Given that there is an instability problem of CECG on the anisotropic cosmological background, we are now interested in the linear stability of SSS vacuum solutions in ECG. For this purpose, we do not deal with ECG as a trivial effective field theory (EFT) where the cubic Lagrangian is always strongly suppressed relative to the Einstein-Hilbert term. We consider odd-parity perturbations according to the Regge-Wheeler formulation [47] without necessarily imposing the condition $f = h$. We show that the Lagrangian \mathcal{P} in ECG gives rise to three propagating DOFs in the odd-parity sector. We find that there is at least one ghost mode and that the squared propagation speed of one of the dynamical perturbations is negative for large multipoles in the regime where the effective mass of perturbations is below the Planck mass, or the cutoff of the theory M . The presence of these ghost and Laplacian instabilities excludes the SSS vacuum solutions in ECG with unsuppressed higher-order curvature terms, including the BH solution with $f = h$.

This paper is organized as follows. In Sec. 2, we revisit the SSS BH solution present in ECG. In Sec. 3, we derive the second-order action of odd-parity perturbations and show how the ghost and Laplacian instabilities arise for the large frequency and momentum modes. Sec. 4 is devoted to conclusions.

2. Cubic gravity

General cubic gravity theories consist of a combination of cubic products of the Riemann tensor $R_{\alpha\beta\gamma\delta}$, Ricci tensor $R_{\mu\nu}$, and Ricci scalar R . We consider the following eight cubic Lagrangians:

$$\begin{aligned} \mathcal{L}_1 &= R_{\alpha}{}^{\beta}{}_{\gamma}{}^{\delta} R_{\beta}{}^{\mu}{}_{\delta}{}^{\nu} R_{\mu}{}^{\alpha}{}_{\nu}{}^{\gamma}, & \mathcal{L}_2 &= R_{\alpha\beta}{}^{\gamma\delta} R_{\gamma\delta}{}^{\mu\nu} R_{\mu\nu}{}^{\alpha\beta}, & \mathcal{L}_3 &= R_{\alpha\beta\gamma\delta} R^{\alpha\beta\gamma}{}_{\mu} R^{\delta\mu}, & \mathcal{L}_4 &= R R_{\alpha\beta\gamma\delta} R^{\alpha\beta\gamma\delta}, \\ \mathcal{L}_5 &= R_{\alpha\beta\gamma\delta} R^{\alpha\gamma} R^{\beta\delta}, & \mathcal{L}_6 &= R_{\alpha}{}^{\beta} R_{\beta}{}^{\gamma} R_{\gamma}{}^{\alpha}, & \mathcal{L}_7 &= R R_{\alpha\beta} R^{\alpha\beta}, & \mathcal{L}_8 &= R^3. \end{aligned} \quad (2.1)$$

Taking into account the Einstein-Hilbert term $M_{\text{pl}}^2 R/2$, where M_{pl} is the reduced Planck mass, we express the total action as

$$S = \frac{M_{\text{pl}}^2}{2} \int d^4x \sqrt{-g} \left(R + \sum_{i=1}^8 c_i \mathcal{L}_i \right), \quad (2.2)$$

where g is a determinant of the metric tensor $g_{\mu\nu}$, and c_i 's are constants.

ECG is constructed to possess a transverse and massless graviton spectrum as in GR on a maximally symmetric background [26]. Requiring also that the relative coefficients of different curvature terms are the same in all dimensions, there is a unique combination \mathcal{P} which is neither trivial nor topological in four dimensions. This nontrivial cubic interaction corresponds to the choice of coefficients $c_1 = 12$, $c_2 = 1$, $c_5 = -12$, and $c_6 = 8$, i.e.,

$$\mathcal{P} = 12\mathcal{L}_1 + \mathcal{L}_2 - 12\mathcal{L}_5 + 8\mathcal{L}_6. \quad (2.3)$$

In general curved spacetime including the SSS background as well as the FLRW background, the equations of motion following from the Lagrangian \mathcal{P} are higher than the second order. In such cases, higher-order derivatives can induce extra DOFs in comparison to those in GR.

In Ref. [36], the authors took a different approach to the construction of cubic gravity theories (dubbed generalized quasi-topological gravity) by demanding that the vacuum SSS solution is fully characterized by a single field equation. In general, the line element on the SSS background is given by

$$ds^2 = -f(r)dt^2 + h^{-1}(r)dr^2 + r^2 \left(d\theta^2 + \sin^2\theta d\varphi^2 \right), \quad (2.4)$$

where f and h are functions of the radial coordinate r . If $f(r)$ is proportional to $h(r)$ such that $f(r) = Nh(r)$, where N is a constant, the time and radial components of the field equations coincide with each other. This gives the following constraints among the coefficients in Eq. (2.2):

$$c_4 = \frac{3c_1 - 36c_2 - 14c_3}{56}, \quad c_5 = -\frac{3c_1 + 48c_2 + 14c_3}{7}, \quad c_7 = \frac{6c_1 + 96c_2 + 14c_3 - 21c_6}{28}, \quad c_8 = -\frac{3c_1 + 20c_2 - 7c_6}{56}. \quad (2.5)$$

The six-dimensional Euler density \mathcal{X}_6 corresponds to the coefficients $c_1 = 8$, $c_2 = -4$, $c_3 = 24$, and $c_6 = -16$, but this is a topological term that does not affect the field equations of motion. One of the remaining three cubic interactions is the Lagrangian \mathcal{P} given by Eq. (2.3). There are also two additional terms

$$C = \mathcal{L}_3 - \frac{1}{4}\mathcal{L}_4 - 2\mathcal{L}_5 + \frac{1}{2}\mathcal{L}_7, \quad (2.6)$$

$$C' = \mathcal{L}_6 - \frac{3}{4}\mathcal{L}_7 + \frac{1}{8}\mathcal{L}_8, \quad (2.7)$$

which correspond to the coefficients $c_1 = 0$, $c_2 = 0$, $c_3 = 1$, $c_6 = 0$ and $c_1 = 0$, $c_2 = 0$, $c_3 = 0$, $c_6 = 1$, respectively. As we will see below, for $f = h$, the contributions of the Lagrangians C and C' to the field equations of motion vanish. On the FLRW spacetime, the Lagrangian \mathcal{P} gives rise to the Friedmann equation higher than second-order. The specific combination $\mathcal{P} - 8C$ leads to the second-order field equations on the FLRW background [39,40]. If we apply this latter cubic theory to inflation, the presence of small anisotropies close to the de Sitter background generates the instability of cosmological solutions [46].

In both ECG and CECG, the dynamical DOFs are more than those in GR (two tensor polarizations) around general, curved backgrounds. The property that the propagating DOFs around the maximally symmetric background degenerate to those in GR implies that the disappearing DOFs are, in general, strongly coupled [44,48]. In such theories, the maximally symmetric background corresponds to a singular surface at which the coefficients of higher-order kinetic terms appearing in the perturbation equations of motion are degenerate. This degeneracy leads to the divergence of couplings of the canonically normalized fields, which is a signal of the strong coupling. On the background different from the maximally symmetric spacetime, some pathological behavior like the instability of perturbations usually arises as it happens for the cosmological background [44,46]. In this paper, we would like to study whether this is also the case for BHs on the SSS background.

On the SSS background given by the line element (2.4), let us consider cubic gravity theories given by the action

$$S = \frac{M_{\text{pl}}^2}{2} \int d^4x \sqrt{-g} (R + \alpha \mathcal{P} + \kappa C + \mu C'), \quad (2.8)$$

where α , κ , μ are constants. Then, the quantities (2.3), (2.6), and (2.7) reduce, respectively, to

$$\begin{aligned} \mathcal{P} = & \frac{1}{f^4 r^4} [3f'^4 h^3 r^2 - 3f^3 h'(2f''hr + f'h'r - 2f'h)(h'r - 2h + 2) - 6f'^2 f h^2 r \{f''hr + f'(h'r + h - 1)\} \\ & + 6rf' f^2 h \{f'h'(h'r + h) + f''h(h'r + 2h - 2)\}], \end{aligned} \quad (2.9)$$

$$C = -\frac{3}{8f^4 r^4} (f'h - h'f)^2 \left[(2f''fh - hf'^2 + f'h'f)r^2 - 4f^2(h - 1) \right], \quad (2.10)$$

$$C' = \frac{1}{2}C, \quad (2.11)$$

where a prime represents the derivative with respect to r . If f is equal to h , both C and C' vanish. Varying the action (2.8) with respect to f and h , we obtain the third-order differential equations for h and f , respectively.¹ Setting $f(r) = h(r)$, the two differential equations coincide with each other, giving

$$rf' + f - 1 + \frac{6\alpha}{r^3} [rf'^2(4f - 1) + rf\{r^2 f''^2 + 4f''(f - 1) - 2rf'''(f - 1)\} + f'f(r^3 f''' - 4r^2 f'' - 4f + 4)] = 0. \quad (2.12)$$

For $\alpha = 0$, the solution to Eq. (2.12) is $f = h = 1 - r_h/r$, where r_h is an integration constant corresponding to the horizon radius. For $\alpha \neq 0$, there are corrections to the Schwarzschild metric. At large distances away from the horizon ($r \gg r_h$), we derive the solution to Eq. (2.12) under the following expansion

$$f(r) = h(r) = 1 - \frac{r_h}{r} + \sum_{i=2} c_i \left(\frac{r_h}{r}\right)^i, \quad (2.13)$$

where c_i 's are constants. Substituting Eq. (2.13) into Eq. (2.12) and computing the coefficients at each order, we obtain $c_2 = c_3 = c_4 = c_5 = 0$, $c_6 = 54\alpha/r_h^4$, and $c_7 = -46\alpha/r_h^4$. Then, the leading-order correction to f arises at sixth order in the expansion (2.12), so that²

$$f(r) = h(r) = 1 - \frac{r_h}{r} + \frac{54\alpha r_h^2}{r^6} \left[1 + \mathcal{O}\left(\frac{r_h}{r}\right) \right]. \quad (2.14)$$

The solution (2.14) has been derived by requiring that $f(r) = h(r)$ without explicitly imposing any requirement on the value of α . The same solution also follows without assuming the condition $f(r) = h(r)$ in ECG with

¹ If we relax the assumption of staticity, it can be shown that for this theory the Birkhoff theorem does not hold in general. Therefore, there could be other solutions that in principle have some relevance but will be in general time-dependent. We will not investigate their existence, however, as we will already set strong constraints on the theory just by looking at their static limit.

² Here we do not study whether the series converges, as in any case, the solution loses its validity for large enough values of r , but we assume that it can be considered as a good approximation in the physical range of r of interest to the real numerical solution having suitable asymptotically flat boundary conditions.

$$\mathcal{P} \neq 0, \quad \mathcal{C} = 0, \quad \mathcal{C}' = 0. \tag{2.15}$$

In this case, we write the large-distance solutions as

$$f(r) = 1 - \frac{r_h}{r} + \sum_{i=2} c_i \left(\frac{r_h}{r}\right)^i, \quad h(r) = 1 - \frac{r_h}{r} + \sum_{i=2} d_i \left(\frac{r_h}{r}\right)^i. \tag{2.16}$$

We substitute Eq. (2.16) into the third-order differential equations of f and h and obtain the coefficients c_i and d_i . This gives rise to the same coefficients c_i derived above, with $c_i = d_i$, so that the large-distance solution is again given by Eq. (2.14). Hence the BH solution with $f(r) = h(r)$ generically arises in ECG.

3. Odd-parity perturbations on the SSS background

We study the stability of SSS vacuum solutions in ECG given by the action

$$\mathcal{S} = \frac{M_{\text{pl}}^2}{2} \int d^4x \sqrt{-g} (R + \alpha \mathcal{P}). \tag{3.1}$$

In our analysis, we do not restrict ourselves to the EFT regime where $\alpha \mathcal{P}$ is always suppressed relative to R . The modification to the Schwarzschild BH in GR can be significant by allowing for the possibility that $\alpha \mathcal{P}$ can be as large as R . This is analogous to Starobinsky's model given by the Lagrangian $L = R + \beta R^2$ [49], where the cosmic acceleration occurs in the regime $\beta R^2 \gtrsim R$. If we stick to the EFT regime with $\beta R^2 \ll R$, one cannot accommodate the physics of inflation driven by the quadratic curvature term.

On the background (2.4), we decompose the metric tensor into $g_{\mu\nu} = g_{\mu\nu}^{(0)} + h_{\mu\nu}$, where $g_{\mu\nu}^{(0)}$ and $h_{\mu\nu}$ correspond to the background and perturbed parts, respectively. Although we are primarily interested in the stability of the BH solution (2.14), we do not impose the condition $f = h$ for the background metric from the beginning. Under the rotation in the (θ, φ) plane, the metric perturbations $h_{\mu\nu}$ can be separated into odd- and even-parity modes [47,50]. Expanding $h_{\mu\nu}$ in terms of the spherical harmonics $Y_{lm}(\theta, \varphi)$, the odd- and even-modes have parities $(-1)^{l+1}$ and $(-1)^l$, respectively. In the odd-parity sector, the components of $h_{\mu\nu}$ are given by [51–55]

$$\begin{aligned} h_{tt} = h_{tr} = h_{rr} &= 0, & h_{ab} &= 0, \\ h_{ta} &= \sum_{l,m} Q(t, r) E_{ab} \nabla^b Y_{lm}(\theta, \varphi), & h_{ra} &= \sum_{l,m} W(t, r) E_{ab} \nabla^b Y_{lm}(\theta, \varphi), \\ h_{ab} &= \frac{1}{2} \sum_{l,m} U(t, r) [E_a^c \nabla_c \nabla_b Y_{lm}(\theta, \varphi) + E_b^c \nabla_c \nabla_a Y_{lm}(\theta, \varphi)], \end{aligned} \tag{3.2}$$

where Q , W , and U depend on t and r , and the subscripts a and b denote either θ or φ . E_{ab} is an antisymmetric tensor with nonvanishing components $E_{\theta\varphi} = -E_{\varphi\theta} = \sin\theta$. In a strict sense, we should write subscripts l and m for the variables Q , W , and U , but we omit them for brevity. Under the gauge transformation $x_\mu \rightarrow x_\mu + \xi_\mu$, where $\xi_t = 0$, $\xi_r = 0$, and $\xi_a = \sum_{l,m} \Lambda(t, r) E_{ab} \nabla^b Y_{lm}(\theta, \varphi)$, metric perturbations transform as $Q \rightarrow Q + \dot{\Lambda}$, $W \rightarrow W + \Lambda' - 2\Lambda/r$, and $U \rightarrow U + 2\Lambda$. There are several gauge-invariant combinations like

$$\hat{W} \equiv W - \frac{1}{2}U' + \frac{1}{r}U, \quad \hat{Q} \equiv Q - \frac{1}{2}\dot{U}. \tag{3.3}$$

We fix the residual gauge DOF by choosing

$$U = 0, \tag{3.4}$$

under which $\hat{W} = W$ and $\hat{Q} = Q$.

For the purpose of expanding the action (3.1) up to second order in odd-parity perturbations, we will focus on the axisymmetric modes ($m = 0$) without loss of generality because nonaxisymmetric modes ($m \neq 0$) can be restored under a suitable rotation. For the integrals with respect to θ and φ , we exploit the following properties

$$\begin{aligned} \int_0^{2\pi} d\varphi \int_0^\pi d\theta (Y_{l0,\theta})^2 \sin\theta &= l(l+1), & \int_0^{2\pi} d\varphi \int_0^\pi d\theta \left[\frac{(Y_{l0,\theta})^2}{\sin\theta} + (Y_{l0,\theta\theta})^2 \sin\theta \right] &= l^2(l+1)^2, \\ \int_0^{2\pi} d\varphi \int_0^\pi d\theta \left[\left(\frac{2}{\sin\theta} - \frac{3}{\sin^3\theta} \right) (Y_{l0,\theta})^2 + \frac{3}{\sin\theta} (Y_{l0,\theta\theta})^2 + \sin\theta (Y_{l0,\theta\theta\theta})^2 \right] &= \frac{1}{4} l^2(l+1)^2 (4l^2 - 2l - 3). \end{aligned} \tag{3.5}$$

After the integration with respect to θ and φ , the second-order action of perturbations, which is expressed in the form $S^{(2)} = \int dt dr L$, contains time derivatives such as \dot{W}^2 , \dot{Q}'^2 , and \dot{Q}^2 . They can be factored out as

$$L_K \equiv \frac{3\alpha M_{\text{pl}}^2 \sqrt{h} [2f'hr + f(2-2h-rh')l(l+1)]}{2f^{5/2}r^2} \left(\ddot{W} - \dot{Q}' + \frac{2\dot{Q}}{r} \right)^2, \tag{3.6}$$

where $L_K \in L$. Then, the rest of the Lagrangian $L - L_K$ does not contain products like $\ddot{W}\dot{Q}'$ and $\ddot{W}\dot{Q}$. We introduce a Lagrange multiplier χ that helps us to understand the presence and behavior of propagating DOFs. Then, the Lagrangian equivalent to L can be written as $L - L_K + \dot{L}_K$ where

$$\tilde{L}_K = \frac{3\alpha M_{\text{pl}}^2 \sqrt{h} [2f'hr + f(2 - 2h - rh')]l(l+1)}{2f^{5/2}r^2} \left[2\chi \left(\ddot{W} - \dot{Q}' + \frac{2\dot{Q}}{r} \right) - \chi^2 \right]. \tag{3.7}$$

Indeed, the variation of (3.7) with respect to χ leads to $\chi = \ddot{W} - \dot{Q}' + 2\dot{Q}/r$, so that (3.7) reduces to (3.6). We note that the gauge-invariant field $\chi = \ddot{W} - \dot{Q}' + 2\dot{Q}/r$ corresponds to a time derivative of the dynamical perturbation $\tilde{\chi} = \dot{W} - Q' + 2Q/r$ in GR [47,53,55] (up to a free function of r). Integrating the Lagrangian $\chi \ddot{W}$ by parts in Eq. (3.7), we obtain the product $-\dot{W} \dot{\chi}$ between the two first derivatives. As we will see below, there are three propagating DOFs $\vec{\chi} = (W, \chi, Q)$ in this system. In other words, we generally need to give six independent initial conditions to determine the time evolution of $\vec{\chi}$.

After the integration by parts, the total second-order can be expressed in the form $\mathcal{S}^{(2)} = \int dt dr L$, where

$$L = a_1 \dot{W}^2 + a_2 \dot{Q}^2 + 2a_3 \dot{W} \dot{\chi} + a_4 \left(\dot{W}' - Q'' + \frac{2Q'}{r} \right)^2 + a_5 W'^2 + a_6 Q'^2 + a_7 W^2 + a_8 \chi^2 + a_9 Q^2 + a_{10} W' \dot{Q} + a_{11} \dot{W} Q' + a_{12} \dot{\chi} Q' + a_{13} \dot{W} Q + a_{14} \chi \dot{Q}, \tag{3.8}$$

with a_1, \dots, a_{14} being functions of r alone. Varying the Lagrangian (3.8) with respect to W, χ , and Q , we obtain the field equations of motion for these perturbations. They contain the derivatives up to the fourth order, e.g., \ddot{W}'' , \dot{Q}''' .

In the following, we study the propagation of fast oscillating modes by assuming the solutions in the form

$$\vec{\chi} = \vec{\chi}_0 e^{i(\omega t - kr)}, \quad \text{with} \quad \vec{\chi}_0 = (W_0, \chi_0, Q_0). \tag{3.9}$$

Because of staticity, here, the coefficients $\vec{\chi}_0$ are supposed to be functions slowly varying in the r -direction, so that in the WKB domain they satisfy for instance the condition $|\dot{\vec{\chi}}_0| \ll |k\vec{\chi}_0|$. For the same reason, we will also suppose that $|\omega'| \ll |k\omega| \simeq |\omega^2|$. As a consequence, we will also consider the wavenumber k and the multipole l in the ranges $kr_h \gg 1$ and $l \gg 1$.

Then, each coefficient in Eq. (3.8) has the following l dependence:

$$a_1 = b_1 l^4, \quad a_2 = b_2 l^4, \quad a_3 = b_3 l^2, \quad a_4 = b_4 l^2, \quad a_5 = b_5 l^4, \quad a_6 = b_6 l^4, \quad a_7 = b_7 l^6, \quad a_8 = b_8 l^2, \\ a_9 = b_9 l^6, \quad a_{10} = b_{10} l^4, \quad a_{11} = b_{11} l^4, \quad a_{12} = b_{12} l^2, \quad a_{13} = b_{13} l^4, \quad a_{14} = b_{14} l^2, \tag{3.10}$$

where

$$b_1 = \frac{3\alpha M_{\text{pl}}^2 h^{1/2} [fr(2f''hr + f'h'r - 6f'h) + 2f^2(h'r + 2h - 2) - f'^2 hr^2]}{4f^{5/2}r^4}, \\ b_2 = \frac{3\alpha M_{\text{pl}}^2 [(2f''f - f'^2)hr + h'f(f'r - 2f)]}{2f^{7/2}h^{1/2}r^3}, \\ b_3 = \frac{r^2}{f}b_1 - \frac{hr^2}{2}b_2 + \frac{b_0}{2fh}, \quad b_4 = fhb_3 + \frac{3}{2}b_0, \quad b_5 = f^2h^2b_2 - \frac{f}{r^2}b_0, \quad b_6 = b_1 + \frac{2}{hr^2}b_0, \\ b_7 = -\frac{b_5}{2hr^2}, \quad b_8 = b_3, \quad b_9 = \frac{f}{2r^2}b_2, \quad b_{10} = -\frac{3\alpha M_{\text{pl}}^2 \sqrt{h}(3f'h + h'f)}{f^{3/2}r^3}, \\ b_{11} = -2b_1 - 2fhb_2 - b_{10} - \frac{b_0}{hr^2}, \quad b_{12} = -2b_3, \tag{3.11}$$

with

$$b_0 = \frac{3\alpha M_{\text{pl}}^2 h^{3/2}(f'h - h'f)}{f^{3/2}r}. \tag{3.12}$$

We do not show the explicit forms of b_{13} and b_{14} , as they are not needed in the following discussion.

Exploiting the approximation of large values of ω, k , and l in the field equations of motion, we ignore terms proportional to $i\omega, ik$ and l relative to those proportional to ω^2, k^2 and l^2 . Then, the perturbation equations of motion following from the Lagrangian (3.8) can be expressed in the form

$$\mathbf{A} \vec{\chi}_0^T = 0, \tag{3.13}$$

where \mathbf{A} is the 3×3 matrix whose components are given by

$$A_{11} = 2l^2 \left[(b_4k^2 + b_1l^2)\omega^2 + b_5k^2l^2 + b_7l^4 \right], \quad A_{22} = 2l^2 b_8, \quad A_{33} = 2l^2 \left(b_2l^2\omega^2 + b_4k^4 + b_6k^2l^2 + b_9l^4 \right), \\ A_{12} = A_{21} = 2l^2 b_3\omega^2, \quad A_{13} = A_{31} = l^2(2b_4k^3 - b_{10}kl^2 - b_{11}kl^2)\omega, \quad A_{23} = A_{32} = -l^2 b_{12}k\omega. \tag{3.14}$$

3.1. Ghost instability

To derive the no-ghost conditions, we pick up the matrix components in \mathbf{A} containing terms proportional to ω^2 , i.e.,

$$\mathbf{K} = 2\omega^2 \begin{pmatrix} K_{11} & K_{12} & 0 \\ K_{12} & 0 & 0 \\ 0 & 0 & K_{33} \end{pmatrix}, \tag{3.15}$$

where

$$K_{11} = l^2(b_4k^2 + b_1l^2), \quad K_{12} = l^2b_3, \quad K_{33} = l^4b_2. \quad (3.16)$$

The absence of ghosts requires the following three conditions

$$K_{11} > 0, \quad -K_{12}^2 > 0, \quad -K_{12}^2K_{33} > 0. \quad (3.17)$$

The explicit form of K_{12} is given by

$$K_{12} = -\frac{3\alpha M_{\text{pl}}^2 h^{1/2} [r(2f'h - h'f) + 2f(1-h)]}{2f^{5/2}r^2} l^2. \quad (3.18)$$

So long as $\alpha \neq 0$, the second condition of Eq. (3.17) is always violated and hence we have at least one ghost mode. The existence of the dynamical perturbation χ , which is kinetically coupled to W , gives rise to the ghost DOF. This result generally holds without restricting the background BH solution to the form (2.14), as we have not assumed the condition $f = h$.

It is important to understand how many ghost DOFs are generated in ECG, at least for the vacuum case. A simple way to see this is to make the field redefinitions: $W = W_2 - K_{12} \chi_2 / K_{11}$, $\chi = \chi_2$, and $Q = Q_2$. The Jacobian of the transformation always has a determinant equal to unity, so that the transformation is well defined. On using these new fields, the kinetic matrix becomes diagonal with elements K_{11} , $-K_{12}^2 / K_{11}$, and K_{33} . For the BH solution (2.14), which is approximately valid for $r \gg r_h$, we find that

$$K_{33} \simeq -\frac{9\alpha M_{\text{pl}}^2 r_h l^4}{r^5}, \quad \frac{K_{33}}{K_{11}} \simeq \frac{2l^2}{k^2 r^2} > 0. \quad (3.19)$$

This shows that, at least for the approximate BH solution, K_{11} and K_{33} share the same sign. Therefore, in the range of validity of this solution, we would have one ghost only if $\alpha < 0$, whereas we would have two ghost DOFs if instead $\alpha > 0$. This calculation also shows that the kinetic term of, e.g., the mode Q , tends to vanish as $r_h/r \rightarrow 0$. This limit then needs to be taken with care.

3.2. Radial propagation

The propagation speeds $c_r = dr_*/d\tau$ along the radial direction, where $r_* = \int dr/\sqrt{h}$ is the rescaled radial coordinate and $\tau = \int \sqrt{f} dt$ is the proper time, are expressed as $c_r = (fh)^{-1/2}(\partial\omega/\partial k)$. We compute the eigenvalue of the matrix \mathbf{A} and expand it with respect to the large momentum k . This amounts to considering the modes in the range $kr_h \gg l \gg 1$. The leading-order k^8 dependent term vanishes on account of the relation $b_{12} = -2b_3$, so the dominant terms in $\det \mathbf{A}$ are those proportional to k^6 . Then, we obtain the following three solutions to the squared radial propagation speeds:

$$c_{r1}^2 = \frac{4b_4b_8f^4r^4}{9\alpha^2M_{\text{pl}}^4h^2[(h'r + 2h - 2)f - 2f'hr]^2} = 1 + \frac{3r(f'h - h'f)}{(h'r + 2h - 2)f - 2f'hr}, \quad (3.20)$$

$$c_{r2}^2 = \frac{-(b_1 + b_6 + b_{10} + b_{11}) + \sqrt{\mathcal{D}_1}}{2b_2fh} = 1, \quad (3.21)$$

$$c_{r3}^2 = \frac{-(b_1 + b_6 + b_{10} + b_{11}) - \sqrt{\mathcal{D}_1}}{2b_2fh} = 1 - \frac{2f(f'h - h'f)}{2f''fhr - f'r(hf' - fh') - 2hf'^2}, \quad (3.22)$$

where

$$\mathcal{D}_1 = b_1^2 + 2b_1b_6 - 4b_2b_5 + b_6^2 + b_{10}^2 + 2(b_1 + b_6 + b_{11})b_{10} + 2(b_1 + b_6)b_{11} + b_{11}^2 = \frac{9\alpha^2M_{\text{pl}}^4h(f'h - h'f)^2}{f^3r^6}. \quad (3.23)$$

For the BH solution (2.14) we have $f = h$, in which case all the three squared propagation speeds (3.20)-(3.22) reduce to 1.

3.3. Angular instability

The propagation speeds $c_\Omega = r d\theta/d\tau$ along the angular direction are expressed as $c_\Omega = r\omega/(\sqrt{f}l)$. Expanding $\det \mathbf{A}$ with respect to large l in the range $l \gg kr_h \gg 1$, the leading-order terms, which are proportional to l^4 , give the following squared angular propagation speeds:

$$c_{\Omega 1}^2 = \frac{r^2(b_1b_8 + \sqrt{\mathcal{D}_2})}{2b_3^2f} = 1, \quad (3.24)$$

$$c_{\Omega 2}^2 = \frac{r^2(b_1b_8 - \sqrt{\mathcal{D}_2})}{2b_3^2f} = \frac{[2f''fhr - f'(f'hr - h'fr + 2fh)]r}{2f[r(h'f - 2f'h) + 2f(h-1)]}, \quad (3.25)$$

$$c_{\Omega 3}^2 = -\frac{b_9r^2}{b_2f} = -\frac{1}{2}, \quad (3.26)$$

where

$$\begin{aligned} \mathcal{D}_2 &= b_8(b_1^2 b_8 + 4b_3^2 b_7) \\ &= \frac{81\alpha^4 M_{\text{pl}}^8 h^2 [r(2f'h - h'f) - 2f(h-1)]^2 [2f''fhr^2 - f'r(f'hr - h'fr - 2fh) - 2f^2(h'r + 2h - 2)]^2}{64f^{10}r^{12}}. \end{aligned} \quad (3.27)$$

Using the BH solution (2.14) in the regime $r \gg r_h$, the second propagation speed squared (3.25) has the following behavior

$$c_{\Omega 2}^2 = 1 - \frac{720\alpha r_h}{r^5} + \mathcal{O}(r^{-10}), \quad (3.28)$$

which quickly approaches 1 at large distances.

The third propagation speed squared (3.26) is negative, so there is Laplacian instability along the angular direction. This instability arises from the unhealthy propagation of the perturbation Q . In the region slightly away from the horizon, the typical time for the instability to occur is of order $T \simeq \mathcal{O}(1)/|\omega| \approx r_h/(\sqrt{f}l) \ll r_h$. We also note that the condition $f = h$ was not used for the derivation of $c_{\Omega 3}^2$. Due to this angular instability besides the appearance of a ghost, the SSS BH solutions with unsuppressed cubic-order terms are not stable.

3.4. Instability versus EFT mass breaking limit

In what follows, we study the regime of validity of the BH solution and conditions under which our previous analysis can be trusted, especially regarding the presence of instabilities. Indeed, we know that one or two ghosts are present and that the field Q acquires a negative propagation speed squared along the angular direction. The last instability is purely (and already) classical, and as such, probably the most serious one. We would like to see here whether this instability for the field Q occurs anywhere in the spatial slicing with l in the range $l \gg kr_h$. In order to study this, we consider the partial differential equations, focusing on the unstable mode, which corresponds to the mass term of the perturbation Q .

Focusing on the term which makes Q unstable and also looking for the case $r \gg r_h$, we find that

$$\ddot{Q} \simeq \left(\frac{l^2}{2r^2} - \frac{r_h^2}{36\alpha} \frac{r^3}{r_h^3} \right) Q + \dots \quad (3.29)$$

where the dots stand for other terms which are irrelevant to the following discussion. For example, in Eq. (3.29), we also have a term of the form

$$\frac{r^5}{36l^2\alpha r_h} Q'', \quad (3.30)$$

but in the following, we will show that, for the Laplacian instability to occur, we need to be in a region of the spatial geometry (and parameter space) for which $l^2 \gg r^5/(18|\alpha|r_h)$. This sets the coefficient of Q'' very small. Furthermore, since we are considering the case in which the angular instability occurs, we also need to assume that the angular derivatives overcome the r -derivatives, that is, $l^2|Q|/r^2 \gg |Q''|$. For the same reason, the coefficients in the terms $-r^5\dot{W}'/(36l^2\alpha r_h)$ and $-r^4\dot{W}/(18l^2\alpha r_h)$ are suppressed and can be thought to be subdominant, also because the field W does not exhibit instabilities along the angular direction (as it propagates with a speed approximately equal to unity).

In any case, from Eq. (3.29), we can define an effective squared mass, the self-coupling term independent of l (or k), for the variable Q as follows

$$\mu_Q^2 = \frac{r_h^2}{36|\alpha|} \frac{r^3}{r_h^3}, \quad (3.31)$$

where we suppose for the moment that $\alpha > 0$. We see that a negative value of α would make the instability even worse, but our goal here is not trying to solve the partial differential equations, even approximately, for $r \rightarrow \infty$. This is because, in fact, it is clear that the term proportional to l^2 in Eq. (3.29) decreases as r increases, whereas the opposite happens for μ_Q^2 . Evidently, this increase in the mass of the perturbation Q cannot approach infinity. In fact, we would expect that the theory reaches an EFT cutoff so that it needs to be ultraviolet (UV) completed. In other words, this behavior implies that the asymptotically flat limit for the theory needs a UV completion. Let us call this cutoff mass scale $M \lesssim M_{\text{pl}}$. Then, in this case, the instability really occurs only if

$$\frac{r_h^2}{36|\alpha|} \frac{r^3}{r_h^3} \ll M^2. \quad (3.32)$$

This implies that, if the instability occurs at about $r = \beta r_h$, then we require that

$$M_{\text{pl}}^2 \gtrsim M^2 \gg \beta^3 \frac{r_h^2}{36|\alpha|} \quad \text{or} \quad |\alpha| \gg \frac{\beta^3}{36(4\pi)^2} \frac{M_S^2}{M_{\text{pl}}^4 M^2}, \quad (3.33)$$

where M_S is the BH mass related to r_h as $r_h = M_S/(4\pi M_{\text{pl}}^2)$. If the inequality (3.33) does not hold, we should expect that the solution is outside the EFT domain. In this case, it should either be replaced by another solution (maybe time-dependent) or instead, the spherically symmetric description of gravity in this theory already requires an appropriate UV completion.

At the same time, we require that the BH solution (2.14), if it is inside the EFT domain, should not be so different from the standard Schwarzschild solution. This statement holds true if

$$\frac{54|\alpha|r_h^2}{r^6} \ll \frac{r_h}{r}, \quad (3.34)$$

in particular for $r = \beta r_h$. This condition leads to

$$|\alpha| \ll \frac{\beta^5}{54(4\pi)^4} \frac{M_S^4}{M_{\text{pl}}^8}. \quad (3.35)$$

Putting these two requirements on α together, we find

$$\alpha_{\text{min}}^{r=\beta r_h} \equiv \frac{\beta^3}{36(4\pi)^2} \frac{M_S^2}{M_{\text{pl}}^4 M^2} \ll |\alpha| \ll \alpha_{\text{max}}^{r=\beta r_h} \equiv \frac{\beta^5}{54(4\pi)^4} \frac{M_S^4}{M_{\text{pl}}^8}. \quad (3.36)$$

This condition makes sense only if $\alpha_{\text{min}}^{r=\beta r_h} \ll \alpha_{\text{max}}^{r=\beta r_h}$, that is $M \gg (M_{\text{pl}}/M_S)M_{\text{pl}}$, i.e. $M \gg 5.3 \times 10^{-21}$ GeV for a solar mass BH. A larger BH would make this bound weaker. Hence, in the following, we will consider the range $(M_{\text{pl}}/M_S)M_{\text{pl}} \ll M \lesssim M_{\text{pl}}$. It should be noted that this relation depends on the BH mass. Expressing the dimensionful coupling α in terms of a length scale ℓ_α as $\alpha = \ell_\alpha^4$ and saturating the cutoff mass scale to $M \simeq M_{\text{pl}}$ in order to be able to discuss a more concrete example, the inequality (3.36) translates to

$$\frac{\beta^{3/4}}{2\sqrt{6}\pi} \left(\frac{M_S}{M_{\text{pl}}}\right)^{1/2} \ell_{\text{pl}} \ll |\ell_\alpha| \ll \frac{\beta^{5/4}}{4\pi(54)^{1/4}} \frac{M_S}{M_{\text{pl}}} \ell_{\text{pl}}, \quad (3.37)$$

where $\ell_{\text{pl}} = 1/M_{\text{pl}} = 8.1 \times 10^{-35}$ m is the reduced Planck length. If we consider a BH with $M_S = M_\odot = 2.0 \times 10^{30}$ kg and $\beta = 2$, for instance, the inequality (3.37) corresponds to 3.4×10^{-16} m $\ll |\ell_\alpha| \ll 2.6 \times 10^3$ m. For $|\ell_\alpha|$ close to the upper limit of (3.37), the cubic curvature terms can give rise to appreciable deviation from BHs in GR in the vicinity of the horizon ($r_h \simeq 3 \times 10^3$ m). For increasing M_S and β , both the upper and lower limits of $|\ell_\alpha|$ tend to be larger.

Now, if the relation (3.36) or (3.37) holds, then we can trust the solution as a valid EFT, but it would be unstable, because of the negative angular squared speed of propagation. Indeed, this instability would take place if

$$\frac{l^2}{2r^2} \gg \mu_Q^2, \quad \text{or} \quad l^2 \gg \frac{r^5}{18|\alpha|r_h} \gg 1. \quad (3.38)$$

Assuming also that $l^2/(2r^2) \ll M^2 \lesssim M_{\text{pl}}^2$, we find

$$3 \frac{\alpha_{\text{max}}^{r=\beta r_h}}{|\alpha|} \ll l^2 \ll \frac{\beta^2}{8\pi^2} \frac{M_S^2 M^2}{M_{\text{pl}}^4}. \quad (3.39)$$

These last inequalities are compatible with the conditions (3.36).

We do not find it interesting to discuss the behavior of the BH solution in the limit $r \rightarrow \infty$, as in this case, we soon hit the EFT breaking scale, and hence the analysis would lose its validity.

After the initial submission of this paper, we noticed that Bueno *et al.* put a paper on the arXiv claiming that the BH instability can be avoided within the EFT regime of ECG [56]. First of all, we would like to stress that, unlike the discussion given above, the EFT approach taken in Ref. [56] means that the cubic Lagrangian $\ell_\alpha^4 \mathcal{P}$ is always suppressed relative to the Einstein-Hilbert term. In this case, the cutoff mass scale M for the validity of the theory is related to the cubic coupling constant ℓ_α^4 as $M = 1/\ell_\alpha$. As also the authors admit, the difference in our approach consists of relaxing this bound and considering instead general values for the cutoff M . In this sense, our approach is more general, but it is true, on the other hand, that our bound does not apply to all the parameter spaces in ECG theories. For the choice $M = 1/\ell_\alpha$, the condition (3.32) for the occurrence of Laplacian instability translates to $r^3 \ll 36\ell_\alpha^2 r_h$. The EFT approach requires that the corrections induced by the cubic Lagrangian are small outside the BH horizon. Since $\ell_\alpha \ll r_h$ in this case, the inequality (3.32) holds in the regime $r \ll r_h$, i.e., deep inside the horizon.

As we already mentioned at the beginning of this section, unlike Ref. [56], we have not dealt with the ECG as an EFT where the cubic Lagrangian $\ell_\alpha^4 \mathcal{P}$ is always suppressed relative to R . In this case, the cutoff mass scale M in Eq. (3.32) is not necessarily restricted to taking values of the order of $1/\ell_\alpha$. Then, the Laplacian instability is present outside the horizon ($\beta \geq 1$) for the coupling constant ℓ_α in the wide range of (3.37). In other words, whenever the cubic Lagrangian starts to be comparable to the Ricci scalar, the problem of Laplacian instability emerges besides the ghost and strong coupling problems. Since we always need to be in the EFT domain with $|\ell_\alpha^4 \mathcal{P}| \ll |R|$ to avoid this pathological behavior, it is not possible to deal with any nonperturbative phenomenon in the vicinity of the horizon [48]. Whenever nonperturbative effects come into play in BH physics, we hit the aforementioned problems due to the breakdown of EFT.

4. Conclusions

ECG is a cubic-order gravitational theory constructed to share the massless graviton spectrum similar to that in GR on a maximally symmetric background. Since the Lagrangian \mathcal{P} is beyond the domain of Lovelock theories, the field equations of motion contain derivatives higher than second order in general, curved geometries. At the same time, this suggests that there should be additional propagating DOFs to those in GR for the spacetime different from the maximally symmetric background. On the SSS background given by the line element (2.4), we studied the propagation of odd-parity perturbations and the resulting linear stability conditions of propagating DOFs. In this procedure, we did not deal with the ECG as a rather trivial EFT where the cubic Lagrangian is always strongly suppressed relative to the Einstein-Hilbert term.

At the background level, the metric components of SSS vacuum solutions in ECG obey the third-order single differential Eq. (2.12) with $f = h$. At large distances away from the horizon, the SSS BH solution is approximately given by Eq. (2.14). On using the large-distance

expansion (2.16) of metrics without imposing the condition $f = h$, we also obtain the same result as Eq. (2.14). This shows the universality of SSS BH solutions with $f = h$.

In Sec. 3, we expanded the action (3.1) in ECG up to quadratic order in odd-parity perturbations to see the propagation of dynamical DOFs. There are some higher-order derivatives appearing as the form (3.6) in the action. By introducing a gauge-invariant Lagrange multiplier $\chi = \dot{W} - \dot{Q}' + 2\dot{Q}/r$, one can express Eq. (3.6) as the equivalent Lagrangian (3.7). Then, the total Lagrangian is expressed by the form (3.8), which consists of three dynamical perturbations $\vec{\chi} = (W, \chi, Q)$. In comparison to GR, which contains only one dynamical perturbation $\tilde{\chi} = \dot{W} - Q' + 2Q/r$ in the odd-parity sector, there are two more propagating DOFs in ECG.

For high frequencies and large radial and angular momenta, we derived the perturbation equations in the form (3.13) by assuming the solution as the WKB form (3.9). The ghosts are absent under the three conditions (3.17), but the second condition is always violated for $\alpha \neq 0$. This is the outcome of a kinetic mixing of $\dot{\chi}$ and \dot{W} without the $\dot{\chi}^2$ term. We showed that all the radial propagation speeds reduce to 1 for the BH solution (2.14). Along the angular direction, the squared propagation speed of the perturbation Q is negative ($c_{\Omega 3}^2 = -1/2$) for the coupling constant α in the range (3.36). Thus, the SSS BH in ECG with unsuppressed higher-order curvature terms is excluded by both ghost and Laplacian instabilities.

The results found in this paper are analogous to what happens in the anisotropic cosmological background in CECG [46]. In this case, there are also three propagating DOFs, with the appearance of ghosts and tachyonic instabilities on the quasi-de Sitter background with small anisotropies. We also note that a strong coupling problem arises for both ECG and CECG due to the degeneracy of propagating DOFs around the maximally symmetric background [44,48]. These pathologies may be avoided by restricting ourselves to the EFT domain in which the cubic Lagrangians are strongly suppressed relative to the Einstein-Hilbert term, but the problems of ghosts, instabilities, and strong couplings manifest themselves in the regime where the cubic curvature terms become comparable to R . These results suggest that going beyond the Lovelock domain with unsuppressed cubic curvature terms can cause problems in the spacetime geometry different from the maximally symmetric background. It will be of interest to explore further whether the similar property persists or not for higher-order gravitational theories containing quadratic and quartic curvature terms.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgements

We thank Jose Beltrán Jiménez and Alejandro Jiménez-Cano for useful correspondence. The work of ADF was supported by the Japan Society for the Promotion of Science Grants-in-Aid for Scientific Research No. 20K03969 and by grant PID2020-118159GB-C41 funded by MCIN/AEI/10.13039/501100011033. ST is supported by the Grant-in-Aid for Scientific Research Fund of the JSPS No. 22K03642 and Waseda University Special Research Project No. 2023C-473.

References

- [1] C.D. Hoyle, U. Schmidt, B.R. Heckel, E.G. Adelberger, J.H. Gundlach, D.J. Kapner, H.E. Swanson, *Phys. Rev. Lett.* 86 (2001) 1418, arXiv:hep-ph/0011014.
- [2] E.G. Adelberger, B.R. Heckel, A.E. Nelson, *Annu. Rev. Nucl. Part. Sci.* 53 (2003) 77, arXiv:hep-ph/0307284.
- [3] A. De Felice, S. Tsujikawa, *Living Rev. Relativ.* 13 (2010) 3, arXiv:1002.4928 [gr-qc].
- [4] T. Clifton, P.G. Ferreira, A. Padilla, C. Skordis, *Phys. Rep.* 513 (2012) 1, arXiv:1106.2476 [astro-ph.CO].
- [5] C.M. Will, *Living Rev. Relativ.* 17 (2014) 4, arXiv:1403.7377 [gr-qc].
- [6] K.S. Stelle, *Phys. Rev. D* 16 (1977) 953.
- [7] B. Zwiebach, *Phys. Lett. B* 156 (1985) 315.
- [8] D.J. Gross, J.H. Sloan, *Nucl. Phys. B* 291 (1987) 41.
- [9] D.M. Hofman, *Nucl. Phys. B* 823 (2009) 174, arXiv:0907.1625 [hep-th].
- [10] R.C. Myers, B. Robinson, *J. High Energy Phys.* 08 (2010) 067, arXiv:1003.5357 [gr-qc].
- [11] R.C. Myers, M.F. Paulos, A. Sinha, *J. High Energy Phys.* 08 (2010) 035, arXiv:1004.2055 [hep-th].
- [12] J. Oliva, S. Ray, *Class. Quantum Gravity* 27 (2010) 225002, arXiv:1003.4773 [gr-qc].
- [13] M. Ostrogradsky, *Mem. Acad. St. Petersburg* 6 (1850) 385.
- [14] R.P. Woodard, *Scholarpedia* 10 (2015) 32243, arXiv:1506.02210 [hep-th].
- [15] C. Lanczos, *Ann. Math.* 39 (1938) 842.
- [16] D. Lovelock, *J. Math. Phys.* 12 (1971) 498.
- [17] R.R. Metsaev, A.A. Tseytlin, *Nucl. Phys. B* 293 (1987) 385.
- [18] I. Antoniadis, J. Rizos, K. Tamvakis, *Nucl. Phys. B* 415 (1994) 497, arXiv:hep-th/9305025.
- [19] M. Gasperini, M. Maggiore, G. Veneziano, *Nucl. Phys. B* 494 (1997) 315, arXiv:hep-th/9611039.
- [20] P. Kantz, N.E. Mavromatos, J. Rizos, K. Tamvakis, E. Winstanley, *Phys. Rev. D* 54 (1996) 5049, arXiv:hep-th/9511071.
- [21] T. Torii, H. Yajima, K.-i. Maeda, *Phys. Rev. D* 55 (1997) 739, arXiv:gr-qc/9606034.
- [22] S. Kawai, M.-a. Sakagami, J. Soda, *Phys. Lett. B* 437 (1998) 284, arXiv:gr-qc/9802033.
- [23] C. Cartier, E.J. Copeland, R. Madden, *J. High Energy Phys.* 01 (2000) 035, arXiv:hep-th/9910169.
- [24] S. Tsujikawa, R. Brandenberger, F. Finelli, *Phys. Rev. D* 66 (2002) 083513, arXiv:hep-th/0207228.
- [25] K. Aoki, S. Tsujikawa, arXiv:2303.13717 [gr-qc].
- [26] P. Bueno, P.A. Cano, *Phys. Rev. D* 94 (2016) 104005, arXiv:1607.06463 [hep-th].
- [27] P. Bueno, P.A. Cano, V.S. Min, M.R. Visser, *Phys. Rev. D* 95 (2017) 044010, arXiv:1610.08519 [hep-th].
- [28] R.A. Hennigar, R.B. Mann, *Phys. Rev. D* 95 (2017) 064055, arXiv:1610.06675 [hep-th].
- [29] P. Bueno, P.A. Cano, *Phys. Rev. D* 94 (2016) 124051, arXiv:1610.08019 [hep-th].
- [30] P. Bueno, P.A. Cano, *Class. Quantum Gravity* 34 (2017) 175008, arXiv:1703.04625 [hep-th].
- [31] R.A. Hennigar, M.B.J. Poshteh, R.B. Mann, *Phys. Rev. D* 97 (2018) 064041, arXiv:1801.03223 [gr-qc].

- [32] C. Adair, P. Bueno, P.A. Cano, R.A. Hennigar, R.B. Mann, *Phys. Rev. D* 102 (2020) 084001, arXiv:2004.09598 [gr-qc].
- [33] A.M. Frassino, J.V. Rocha, *Phys. Rev. D* 102 (2020) 024035, arXiv:2002.04071 [hep-th].
- [34] D.J. Burger, W.T. Emond, N. Moynihan, *Phys. Rev. D* 101 (2020) 084009, arXiv:1910.11618 [hep-th].
- [35] P.A. Cano, D. Pereñíguez, *Phys. Rev. D* 101 (2020) 044016, arXiv:1910.10721 [hep-th].
- [36] R.A. Hennigar, D. Kubizňák, R.B. Mann, *Phys. Rev. D* 95 (2017) 104042, arXiv:1703.01631 [hep-th].
- [37] P. Bueno, P.A. Cano, J. Moreno, A. Murcia, *J. High Energy Phys.* 11 (2019) 062, arXiv:1906.00987 [hep-th].
- [38] P. Bueno, P.A. Cano, R.A. Hennigar, *Class. Quantum Gravity* 37 (2020) 015002, arXiv:1909.07983 [hep-th].
- [39] G. Arciniega, P. Bueno, P.A. Cano, J.D. Edelstein, R.A. Hennigar, L.G. Jaime, *Phys. Lett. B* 802 (2020) 135242, arXiv:1812.11187 [hep-th].
- [40] G. Arciniega, J.D. Edelstein, L.G. Jaime, *Phys. Lett. B* 802 (2020) 135272, arXiv:1810.08166 [gr-qc].
- [41] A. Cisterna, N. Grandi, J. Oliva, *Phys. Lett. B* 805 (2020) 135435, arXiv:1811.06523 [hep-th].
- [42] C. Erices, E. Papantonopoulos, E.N. Saridakis, *Phys. Rev. D* 99 (2019) 123527, arXiv:1903.11128 [gr-qc].
- [43] I. Quiros, R. García-Salcedo, T. Gonzalez, J.L.M. Martínez, U. Nucamendi, *Phys. Rev. D* 102 (2020) 044018, arXiv:2003.10516 [gr-qc].
- [44] J. Beltrán Jiménez, A. Jiménez-Cano, *J. Cosmol. Astropart. Phys.* 01 (2021) 069, arXiv:2009.08197 [gr-qc].
- [45] P.A. Cano, K. Fransen, T. Hertog, *Phys. Rev. D* 103 (2021) 103531, arXiv:2011.13933 [hep-th].
- [46] M.C. Pookkillath, A. De Felice, A.A. Starobinsky, *J. Cosmol. Astropart. Phys.* 07 (2020) 041, arXiv:2004.03912 [gr-qc].
- [47] T. Regge, J.A. Wheeler, *Phys. Rev.* 108 (1957) 1063.
- [48] J.B. Jiménez, A. Jiménez-Cano, arXiv:2306.07095 [gr-qc].
- [49] A.A. Starobinsky, *Phys. Lett. B* 91 (1980) 99.
- [50] F.J. Zerilli, *Phys. Rev. Lett.* 24 (1970) 737.
- [51] A. De Felice, T. Suyama, T. Tanaka, *Phys. Rev. D* 83 (2011) 104035, arXiv:1102.1521 [gr-qc].
- [52] H. Motohashi, T. Suyama, *Phys. Rev. D* 84 (2011) 084041, arXiv:1107.3705 [gr-qc].
- [53] T. Kobayashi, H. Motohashi, T. Suyama, *Phys. Rev. D* 85 (2012) 084025, Erratum: *Phys. Rev. D* 96 (2017) 109903, arXiv:1202.4893 [gr-qc].
- [54] R. Kase, L.A. Gergely, S. Tsujikawa, *Phys. Rev. D* 90 (2014) 124019, arXiv:1406.2402 [hep-th].
- [55] R. Kase, S. Tsujikawa, *Phys. Rev. D* 105 (2022) 024059, arXiv:2110.12728 [gr-qc].
- [56] P. Bueno, P.A. Cano, R.A. Hennigar, arXiv:2306.02924 [hep-th].