

Spin-2 Kaluza-Klein scattering in a stabilized warped backgroundR. Sekhar Chivukula^{1,*}, Dennis Foren^{1,†}, Kirtimaan A. Mohan^{2,‡}, Dipan Sengupta^{1,3,§} and Elizabeth H. Simmons^{1,||}¹*Department of Physics and Astronomy, University of California,
9500 Gilman Drive, San Diego, California 92093, USA*²*Department of Physics and Astronomy, Michigan State University,
567 Wilson Road, East Lansing, Michigan 48824, USA*³*ARC Centre of Excellence for Dark Matter Particle Physics, Department of Physics,
University of Adelaide, South Australia 5005, Australia* (Received 27 June 2022; accepted 3 January 2023; published 16 February 2023)

Scattering amplitudes involving massive spin-2 particles typically grow rapidly with energy. In this paper we demonstrate that the anomalous high-energy growth of the scattering amplitudes cancel for the massive spin-2 Kaluza-Klein modes arising from compactified five-dimensional gravity in a stabilized warped geometry. Generalizing previous work, we show that the two sum rules which enforce the cancellations between the contributions to the scattering amplitudes coming from the exchange of the (massive) radion and those from the exchange of the tower of Goldberger-Wise scalar states (admixture of the original gravitational and scalar fields of the theory) still persist in the case of the warping which would be required to produce the hierarchy between the weak and Planck scales in a Randall-Sundrum model. We provide an analytic proof of one combination of these generalized scalar sum rules and show how the sum rule depends on the Einstein equations determining the background geometry and the mode-equations and normalization of the tower of physical scalar states. Finally, we provide a consistent and self-contained derivation of the equations governing the physical scalar modes, and we list, in appendixes, the full set of sum rules ensuring proper high-energy growth of all $2 \rightarrow 2$ massive spin-2 scattering amplitudes.

DOI: [10.1103/PhysRevD.107.035015](https://doi.org/10.1103/PhysRevD.107.035015)**I. INTRODUCTION**

Historically, extradimensional theories of gravity were introduced soon after Einstein's discovery of the general theory of relativity. In the original form extra dimensions were introduced by Kaluza and Klein (KK) to unify electromagnetism with gravity, the only two fundamental forces known at the time [1,2]. Extradimensional models have continued to evolve since the late 1970s, thanks in large part to the development of string theory. Over the past three decades, low-energy realizations of extradimensional models gained prevalence as well-motivated scenarios of physics beyond the standard model. One of the most popular and phenomenologically viable models of extra dimensions is the Randall-Sundrum model [3,4], wherein a

compact extra dimension in anti-de Sitter space is used to generate relative exponential factors; this factor allows particles fixed to a brane to interact at electroweak strength while also ensuring bulk-propagating gravity is weak in the observed extended four-dimensional (4D) space, thus providing a geometric solution for the hierarchy problem of the standard model.

Low-energy four-dimensional effective field theories arising from compactified theories of gravity involve towers of interacting spin-0 and spin-2 fields (and potentially spin-1 fields as well, though these are often eliminated by imposing an orbifold symmetry on the compact extra dimension). The massive spin-2 resonances—sometimes called KK gravitons—are particularly interesting. The existence of self-interactions between these KK gravitons is problematic because typically scattering amplitudes between massive spin-2 particles grow far too rapidly with energy to keep unitarity constraints satisfied much beyond the mass of the lightest massive spin-2 state involved. For example, theories of massive gravity that extend 4D general relativity by adding a Fierz-Pauli mass term [5] result in 2-to-2 scattering amplitudes for the helicity-zero channel (the channel whose amplitude has the highest energy growth) that grow like $s^5 / (m_{\text{FP}}^8 M_{\text{Pl}}^2)$ [6], where m_{FP} is the mass of the graviton, s the squared center-of-mass energy, and M_{Pl} the reduced Planck mass. Adding carefully chosen potential

*rschivukula@physics.ucsd.edu

†dennisforen@gmail.com

‡kamohan@msu.edu

§disengupta@physics.ucsd.edu

||ehsimmons@ucsd.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

terms to the model [7,8] can soften matrix element growth down to $\mathcal{O}(s^3)$. However, compactified theories of extradimensional gravity should—due to higher-dimensional diffeomorphism invariance, requiring a smooth $m_{\text{KK}} \rightarrow 0$ limit—grow at most as $\mathcal{O}(s)$.

As we have demonstrated in earlier work [9–12], while the individual contributions to massive helicity-zero spin-2 scattering amplitudes do each grow like $\mathcal{O}(s^5)$ in compactified gravity theories, cancellations occurring between these contributions ultimately ensure the overall 2-to-2 scattering amplitudes grow no faster than $\mathcal{O}(s)$.¹ These cancellations require that the couplings and masses of the Kaluza-Klein spin-2 and spin-0 modes be constrained to satisfy various sum-rule relations, which we have shown to be satisfied in both flat and warped compactifications.

In the original formulation of the RS1 (Randall-Sundrum I) model, extradimensional gravity generates a massless radion in the effective 4D theory.² A massless radion couples to the trace of the stress-energy tensor, yielding a Brans-Dicke-like theory, in odds with predictions of the general theory of relativity. Additionally, this massless radion sources a Casimir force [15], thereby destabilizing the extra dimension and leading to its collapse. A naive computation of scattering amplitudes of massive spin-2 KK particles in a compactified theory of gravity, with a radion mass introduced by hand, reveals that scattering amplitudes grow like $m_{\text{radion}}^2 s^2 / (m_{\text{KK}}^4 M_{\text{Pl}}^2)$, i.e., $\mathcal{O}(s^2)$ instead of $\mathcal{O}(s)$. To stabilize the extra dimension in a way that retains validity of the theory up to the Planck scale, the radion must not only be made massive, but its mass must be generated *dynamically*. These dynamics must also provide additional contributions guaranteeing that the overall scattering amplitude grows no faster than $\mathcal{O}(s)$ —as we have explicitly shown in [16] and explore here in detail.

In the same year that Randall and Sundrum published their model, Goldberger and Wise published a dynamical mechanism for stabilizing the model’s extra dimension [17,18]. Their mechanism shares conceptual similarities with the standard technique of generating massive gauge bosons by spontaneously breaking the associated underlying gauge symmetry, wherein dynamics producing a nonzero vacuum expectation value (VEV) for a gauge-variant operator induces mixing between the longitudinal components of the gauge bosons with Goldstone bosons. In the Goldberger-Wise (GW) mechanism, a new five-dimensional (5D) bulk scalar field $\hat{\Phi}(x, y)$ is appended to the RS1 model and included in new potential terms that also involve (via standard gravitational factors) the usual RS1 metric fields. That bulk scalar field then spontaneously acquires a background profile $\phi_0(y)$ with

nonconstant dependence on the extradimensional coordinate y , causing mixing between background fluctuations \hat{f} of the bulk scalar field and scalar fluctuations $\hat{\sigma}$ of the RS1 metric. While this could in principle yield two physically relevant superpositions of the bulk scalar field and scalar metric fluctuations, one combination is automatically forced to vanish in unitary gauge, leaving just one physical 5D scalar field in the spectrum.

Quantitatively, the new mixed scalar sector decomposes into a tower of spin-0 modes, all described by a single Sturm-Liouville equation with nontrivial Robin boundary conditions involving delta function contributions at the boundaries of the RS1 geometry. In this way, the GW mechanism ultimately generates an infinite tower of physical massive spin-0 states. But what happened to the radion? If the background profile of the bulk scalar field were, instead, made constant in the extradimensional coordinate, no extradimensional symmetries would be spontaneously broken and all but the lowest state in the spin-0 tower would cease to mix with the gravitational sector of the theory. In this limit, the lowest state would become massless, and—indeed—its couplings would exactly match those of the original unstabilized RS1 radion. Hence, in the GW-stabilized RS1 model we are considering, the radion should be associated with the lightest *massive* state among an entire tower of massive spin-0 states.

The assessment of the validity of the GW-stabilized RS1 effective field theory proceeds as for the corresponding unstabilized case, by calculating $2 \rightarrow 2$ massive KK graviton scattering, now with an extended scalar sector, as compared to only the radion in the unstabilized case. An account of this calculation was provided in [16], where we formulated an extended set of sum rules required to ensure that scattering amplitudes were well behaved in a stabilized theory of extra dimension without reference to any explicit GW model. In addition, we proposed a simple model of a stabilized-but-approximately-flat extra dimension, the “flat stabilized” model, and we demonstrated that the revised sum rules were satisfied in this model.

In this work we extend prior results into a new domain by computing the couplings and masses of the scalar and spin-2 states in a Randall-Sundrum model with a Goldberger-Wise stabilization mechanism in the phenomenologically interesting case in which the warping reproduces the hierarchy between the weak and Planck scales. We provide a self-contained derivation of the equations governing the physical scalar modes. We show how one combination of the generalized sum rules in particular relies explicitly on the equations determining the background bulk scalar field and metric, including the scalar mode wave functions and their normalization conditions. We introduce a model in which the Goldberger-Wise dynamics are a small perturbation away from the

¹See also [13,14].

²Along with the usual massless 4D graviton.

unstabilized warped RS1 model. We then demonstrate numerically that all of the sum rules needed to ensure that the anomalous growth of the scattering amplitudes cancel are satisfied to leading nontrivial order in perturbation theory.

The computations performed in the pure gravity sector in this paper and the preceding ones [9–11,16], demonstrating that amplitudes in compactified extradimensional theories grow no faster than $\mathcal{O}(s)$, have important phenomenological consequences. The underlying higher-dimensional diffeomorphism invariance that ensures that scattering amplitudes are well behaved also guarantees that scattering amplitudes involving matter particles should also be compliant with the same principle. For example, in calculations for cosmological observables such as relic abundances of dark matter with KK graviton portals (for both freeze-in and freeze-out), the velocity averaged cross sections must be properly estimated at large \sqrt{s} . An erroneous estimate with anomalously growing cross sections would lead to inaccurate predictions for cosmological observables within the scope of these models. The subject of KK graviton/massive graviton portals to dark matter have been considered extensively in the literature [19–21]. In many of these works, the velocity averaged cross section has been incorrectly estimated due to anomalously growing scattering amplitudes. An application of our works was considered in [22] where some of these issues were accurately addressed within unstabilized models. An application of this work will be to accurately estimate KK graviton portal scenarios in freeze-in/freeze-out mechanisms within the phenomenologically relevant Goldberger-Wise models, with a massive radion. In addition, the Goldberger-Wise scalar sector of such models has been neglected due to its complexity. Here, we work out the details of the scalar sector. These issues will be considered further in future work.

The rest of the paper is organized as follows. In Sec. II we describe the Lagrangian of the Goldberger-Wise Randall-Sundrum model and set notation for the background geometry. In Sec. III we describe the spin-0 and spin-2 mode expansions. Our analysis of the Kaluza-Klein expansions for this system follows the computations of [23–26], generalized to de Donder gauge, and is presented in detail for completeness and clarity in Appendix A. A review of Kaluza-Klein mode scattering and couplings, and description of the version of the sum rules of [16] used here, as well as a description of the

analytic proof of one combination of the sum rules involving the scalar sector is given in Sec. IV. Details of the analytic proof of the sum rule are given in Appendix B 5 a. We also provide, in the totality of Appendix B, a complete list of the sum-rule relations which must be satisfied for all $2 \rightarrow 2$ massive spin-2 scattering amplitudes to grow no faster than $\mathcal{O}(s)$ —completing the analyses begun in [9–12]. A description and the analysis of the perturbative warped-stabilized model is given in Sec. V. In particular, our numerical checks of the sum rules in this model are illustrated in Figs. 3 and 4 of Sec. V C. Our perturbative analysis requires a slight generalization of Rayleigh-Schrödinger perturbation theory to account for perturbations in the weight function of the corresponding Sturm-Liouville problem, and this formalism is described in Appendix C. Our conclusions are given in Sec. VI. *Mathematica* [27] files giving the expressions for all the spin-2 and spin-0 perturbative wave functions can be found on GitHub [28].

II. THE LAGRANGIAN

In this section we outline schematically how a canonical 4D effective Lagrangian is derived from a 5D RS1 model stabilized by the Goldberger-Wise mechanism. We provide a self-contained discussion of the details of this derivation in Appendix A utilizing arguments similar to those found in Refs. [24–26], generalized to de Donder gauge to enable consistent scattering amplitude computations for processes involving the (massless) graviton. In this section we specify our notation and outline the results needed to present our computations.

We begin by writing down the Lagrangian which consists of the following terms:

$$\mathcal{L}_{5D} \equiv \mathcal{L}_{EH} + \mathcal{L}_{\Phi\Phi} + \mathcal{L}_{\text{pot}} + \mathcal{L}_{\text{GHY}} + \Delta\mathcal{L}. \quad (1)$$

Here \mathcal{L}_{EH} comes from the usual Einstein-Hilbert action, $\mathcal{L}_{\Phi\Phi}$ and \mathcal{L}_{pot} are the kinetic and potential terms, respectively, of a bulk scalar field $\hat{\Phi}(x, y)$, \mathcal{L}_{GHY} is the Gibbons-Hawking-York (GHY) boundary term [29,30], and $\Delta\mathcal{L}$ is a useful total derivative we define in Appendix A. The combination of \mathcal{L}_{GHY} and $\Delta\mathcal{L}$ is required to have a well-posed variational principle for the gravitational action [31]]. This Lagrangian is a function of the 5D metric G , which we parametrize in terms of a 4D metric perturbation $g_{\mu\nu}$ and a scalar metric perturbation \hat{r} as [32]

$$[G_{MN}] = \begin{pmatrix} g_{\mu\nu} \exp\left[-2\left(A(y) + \frac{e^{2A(y)}}{2\sqrt{6}}\kappa\hat{r}(x, y)\right)\right] & 0 \\ 0 & -\left(1 + \frac{e^{2A(y)}}{\sqrt{6}}\kappa\hat{r}(x, y)\right)^2 \end{pmatrix} \quad (2)$$

in terms of coordinates $x^M \equiv (x^\mu, y)$, where $y \in (-\pi r_c, +\pi r_c]$ parametrizes an orbifolded extra dimension³; for convenience we define $\varphi \equiv y/r_c$ and use y or φ interchangeably as the coordinate of the fifth dimension. Note that this form of the metric was used in our previous works [9–11] for the unstabilized RS1 metric, following [32] to bring the quadratic Lagrangian to a canonical form from the outset. In the case of the stabilized model, the situation is further complicated by a nontrivial mixing between the bulk scalar field and the scalar metric fluctuations. As we will explain below and demonstrate explicitly in Appendix A, the total derivative $\Delta\mathcal{L}$ helps us bring the Lagrangian into a canonical form.

The warp factor $A(y)$ encodes the warped background geometry. In RS1 [3], in which the extra dimension is unstabilized, $A(y) = k|y|$, where k is related to the space-time curvature. The Goldberger-Wise mechanism [17,18] complicates the background geometry such that the specific form of $A(y)$ becomes dependent on the details of the mechanism's bulk scalar interactions. Crucially, the scalar (bulk and boundary) potential terms in \mathcal{L}_{pot} are chosen such that the scalar field gains a y -dependent background field value, and the trade-off between the contributions to the action from bulk kinetic energy terms $\mathcal{L}_{\Phi\Phi}$ and the scalar potential(s) stabilizes the size of the extra dimension.

The Lagrangian thus far is written in terms of the metric perturbations $\{g_{\mu\nu}, \hat{r}\}$ and the bulk scalar field $\hat{\Phi}$. We next expand $g_{\mu\nu}$ and $\hat{\Phi}$ about their background values:

$$\begin{aligned} g_{\mu\nu}(x, y) &\equiv \eta_{\mu\nu} + \kappa \hat{h}_{\mu\nu}(x, y), \\ \hat{\Phi}(x, y) &\equiv \frac{1}{\kappa} \hat{\phi} \equiv \frac{1}{\kappa} [\phi_0(y) + \hat{f}(x, y)]. \end{aligned} \quad (3)$$

The background metric $\eta_{\mu\nu} = \text{Diag}(+1, -1, -1, -1)$ of $g_{\mu\nu}$ is determined by demanding Lorentz invariance along the extended dimensions, while the background value ϕ_0/κ of $\hat{\Phi}$ must be found by solving the theory's field equations. We normalize the Lagrangian such that the 5D gravitational coupling κ is related to the 5D Planck mass $M_{\text{Pl},5\text{D}}$ according to $\kappa^2 = 4/M_{\text{Pl},5\text{D}}^3$. Note that the factors of κ (units: energy^{-3/2}) included in Eq. (3) are such that ϕ_0 and \hat{f} are unitless in natural units. Following KK decomposition, the 5D tensor field $\hat{h}_{\mu\nu}$, the 5D scalar field \hat{r} , and the 5D scalar fluctuation field \hat{f} give rise to an infinite tower of 4D states.

Perturbatively expanding the Lagrangian Eq. (1) order-by-order in κ yields terms containing various powers of $h_{\mu\nu}$,

³That is, the extra dimension is a circle in which the points with coordinates y and $-y$ are identified. As we will see, this view of the extra dimension (as opposed to treating it as a line segment) allows us to motivate and use the boundary conditions of the Kaluza-Klein mode equations at the orbifold fixed points at $y = 0$ and $y = \pi$ more easily.

\hat{f} , and \hat{r} . In particular, at quadratic order in the fields, we find a complicated expression; c.f. Eqs. (A58)–(A63). Thankfully, there are residual five-dimensional diffeomorphism transformations which leave the form of Eq. (2) invariant—these transformations allow us to reorganize how the physical content is embedded in the fields and thereby attain explicitly canonical quadratic Lagrangians. This process will also mix the 5D fields \hat{r} and \hat{f} (and their constituent 4D states) together in a process that eventually leaves a single scalar tower of physical states. In particular, to ultimately bring the quadratic 5D Lagrangian into a form suitable for generating canonical 4D Lagrangians, we impose the gauge-fixing constraint⁴

$$(\partial_\varphi \phi_0) \hat{f}(x, y) \equiv \sqrt{6} e^{2A(\varphi)} (\partial_\varphi \hat{r}) \quad (4)$$

to eliminate the field \hat{f} in terms of \hat{r} .⁵ In this gauge the 5D theory's independent field degrees of freedom consist only of the 5D scalar field \hat{r} and the 5D tensor field $\hat{h}_{\mu\nu}$. To yield a 4D effective theory, each of these 5D fields is subsequently decomposed into a tower of 4D KK modes. We emphasize here that bringing the Lagrangian to a canonical form is a nontrivial task, and it is of paramount importance to figure out all the interactions of both the gravitational and the scalar sector that will eventually determine the structure of the matrix elements and the couplings.

To calculate the desired matrix elements, we require the cubic and quartic self-interactions of the 5D tensor field $\hat{h}_{\mu\nu}$ as well as the $\hat{h}\hat{h}\hat{r}$ cubic interaction. The \hat{h} self-interactions (and their 4D effective equivalents) are changed from our previous works [9,10,12] only in the specific choice of $A(y)$. Following integration by parts and the elimination of total derivatives, we find that the $\hat{h}\hat{h}\hat{r}$ interaction is similarly identical to the unstabilized case [11]:

$$\mathcal{L}_{hhr} = -\frac{\kappa}{2r_c^2} \sqrt{\frac{3}{2}} e^{-2A(\varphi)} [(\hat{h}')^2 - \hat{h}'_{\mu\nu} (\hat{h}^{\mu\nu})'] \hat{r}. \quad (5)$$

Thus, the primary difference between the stabilized and unstabilized cases as far as $\hat{h}\hat{h}\hat{r}$ is concerned regards the KK decomposition of the 5D field $\hat{r}(x, y)$. In the unstabilized case, \hat{r} generates only a single massless scalar state $\hat{r}(x)$ (see footnote 5)—the usual RS1 radion. In the stabilized case, \hat{r} has nontrivial y -dependence and instead generates an infinite tower of massive scalars $\{\hat{r}^{(i)}(x)\}$, wherein the lightest of these scalars (with KK number $i = 0$) is identified as the massive radion and the heavier

⁴A demonstration that one can always impose this gauge constraint can be found in Refs. [24,26].

⁵Note that in the limit in which there is no nontrivial scalar background, $\phi'_0 \rightarrow 0$, this constraint leaves only the constant (φ -independent) mode of \hat{r} in the theory—a mode corresponding to the massless radion of the unstabilized theory.

states are called GW scalars.⁶ From here on, we will drop the argument φ in the warp factor $A(\varphi)$ for convenience.

III. KALUZA-KLEIN MODE EXPANSIONS

Upon KK decomposition, the scalar and tensor modes come from extradimensional wave functions which satisfy one-dimensional Sturm-Liouville (SL) problems. For completeness and notational consistency, we provide details of the derivation of the SL problems in the tensor and scalar sectors in Appendix A—following the procedures originally described in [25,26] (see also Ref. [16]). We report the results here for the convenience of the reader and to highlight the differences that arise when solving these problems in the stabilized RS1 model.

A. Spin-2 sector

The tensor field $h_{\mu\nu}(x, y)$ is decomposed into a tower of 4D KK states $\hat{h}_{\mu\nu}^{(n)}(x)$ in the usual way as follows: recalling $\varphi \equiv y/r_c$,

$$\hat{h}_{\mu\nu}(x, y) = \frac{1}{\sqrt{\pi r_c}} \sum_{n=0}^{+\infty} \hat{h}_{\mu\nu}^{(n)}(x) \psi_n(\varphi). \quad (6)$$

Here r_c is the radius of the extra dimension and $\psi_n(\varphi)$ is the 5D wave function of the n th mode that satisfies the following SL differential equation:

$$\partial_\varphi [e^{-4A} \partial_\varphi \psi_n] = -\mu_n^2 e^{-2A} \psi_n, \quad (7)$$

where the wave functions satisfy the boundary conditions where $(\partial_\varphi \psi_n) = 0$ at $\varphi \in \{0, \pi\}$. The eigenvalues $\mu_n = m_n r_c$ are the masses m_n of the n th spin-2 KK mode. The wave functions are normalized as follows:

$$\frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi e^{-2A} \psi_m \psi_n = \delta_{m,n}, \quad (8)$$

and satisfy the completeness relation

$$\delta(\varphi_2 - \varphi_1) = \frac{1}{\pi} e^{-2A} \sum_{j=0}^{+\infty} \psi_j(\varphi_1) \psi_j(\varphi_2). \quad (9)$$

The form of this SL problem is identical to the unstabilized case and differs only in how the new background geometry influences the warp factor A . In the unstabilized case, the warp factor is simply $kr_c|\varphi|$. In the stabilized case, the bulk

⁶Note that, having chosen to express the physical degrees of freedom in terms of the scalar field \hat{r} in Eq. (4), the *form* of the couplings between the massive spin-2 states and the tower of GW states is precisely the same as the form of the radion coupling in RS1. However, as we will see, the mode equation and normalization conditions for the physical GW scalars lead to additional complications.

scalar potential modifies the background geometry such that $A(\varphi)$ satisfies the (Einstein) equation

$$A'' = \frac{1}{12} \left[(\phi'_0)^2 + 4 \sum_{i=1,2} V_i r_c \delta_i \right] \quad (10)$$

in terms of the background scalar field ϕ_0 and the brane-localized potentials $V_{1,2}$ at $\varphi = 0$ and $\varphi = \pi$, respectively (refer to Appendix A for additional details).

B. Spin-0 sector

The spin-0 sector of the model arises from two sources. The first is the scalar metric fluctuation (where even the lightest mode will be y -dependent in the stabilized model), and the second is the new bulk scalar field. The two sectors mix via the gauge condition noted in Eq. (4). Consequently, we attain a single physically relevant 5D physical scalar perturbation $\hat{r}(x, y)$. Similar to the tensor perturbation, the KK decomposition of the 5D scalar field $\hat{r}(x, y)$ into a tower of spin-0 KK modes proceeds by introducing extradimensional wave functions $\gamma_i(\varphi)$ and a tower of 4D scalar fields $\hat{r}^{(i)}(x)$ parametrized as follows:

$$\hat{r}(x, y) = \frac{1}{\sqrt{\pi r_c}} \sum_{i=0}^{+\infty} \hat{r}^{(i)}(x) \gamma_i(\varphi). \quad (11)$$

The mode equation that brings the 5D scalar Lagrangian to canonical form, however, is quite different from the tensor case and involves nontrivial boundary terms

$$\begin{aligned} & \partial_\varphi \left[\frac{e^{2A}}{(\phi'_0)^2} (\partial_\varphi \gamma_i) \right] - \frac{e^{2A}}{6} \gamma_i \\ & = -\mu_{(i)}^2 \frac{e^{4A}}{(\phi'_0)^2} \gamma_i \left\{ 1 + \frac{2\delta(\varphi)}{\left[2\ddot{V}_1 r_c - \frac{\phi''_0}{\phi'_0} \right]} + \frac{2\delta(\varphi - \pi)}{\left[2\ddot{V}_2 r_c + \frac{\phi''_0}{\phi'_0} \right]} \right\}, \end{aligned} \quad (12)$$

where $\phi'_0 \equiv (\partial_\varphi \phi_0)$ and the eigenvalues $\mu_{(n)} = m_{(n)} r_c$ are the masses $m_{(n)}$ of the n th scalar KK mode. The Dirac deltas enforce the following (orbifold) boundary conditions:

$$\begin{aligned} (\partial_\varphi \gamma_i)|_{\varphi=0} & = - \left[2\ddot{V}_1 r_c - \frac{\phi''_0}{\phi'_0} \right]^{-1} \mu_{(i)}^2 e^{2A} \gamma_i|_{\varphi=0}, \\ (\partial_\varphi \gamma_i)|_{\varphi=\pi} & = + \left[2\ddot{V}_2 r_c + \frac{\phi''_0}{\phi'_0} \right]^{-1} \mu_{(i)}^2 e^{2A} \gamma_i|_{\varphi=\pi}, \end{aligned} \quad (13)$$

where $\ddot{V}_{1,2}$ are second functional derivatives of the brane potentials evaluated at the background-field configuration. Note that these boundary conditions reduce to Neumann form in the “stiff-wall” limit, $\ddot{V}_{1,2} \rightarrow +\infty$, a limit which will be useful to us during our numerical work in Sec. V.

In the form of Eq. (12), the Sturm-Liouville nature of the problem is manifest, and the orthogonality and completeness of the wave functions follow immediately [24,25,33,34].⁷

Due to the mixing between the gravitational and bulk scalar sectors in Eq. (4), however, an unconventional normalization of the scalar wave functions is required to bring the scalar kinetic energy terms to canonical form,⁸

$$\delta_{mn} = \frac{6\mu_{(n)}^2}{\pi} \int_{-\pi}^{+\pi} d\varphi \gamma_m \gamma_n \frac{e^{4A}}{(\phi'_0)^2} \times \left\{ 1 + \frac{2\delta(\varphi)}{[2\dot{V}_1 r_c - \frac{\phi''_0}{\phi'_0}] + \frac{2\delta(\varphi - \pi)}{[2\dot{V}_2 r_c + \frac{\phi''_0}{\phi'_0}]} \right\} \quad (14)$$

$$= \frac{6}{\pi} \int_{-\pi}^{+\pi} d\varphi \left[\frac{e^{2A}}{(\phi'_0)^2} \gamma'_m \gamma'_n + \frac{e^{2A}}{6} \gamma_m \gamma_n \right], \quad (15)$$

where the second line follows by applying the differential equation (12) and integration-by-parts on the periodic doubled orbifold. The scalar wave function completeness relation follows from Eq. (14):

$$\delta(\varphi_2 - \varphi_1) = \frac{6}{\pi} \frac{e^{4A(\varphi_1)}}{(\phi'_0(\varphi_1))^2} \times \left\{ 1 + \frac{2\delta(\varphi_1)}{[2\dot{V}_1 r_c - \frac{\phi''_0}{\phi'_0}] + \frac{2\delta(\varphi_1 - \pi)}{[2\dot{V}_2 r_c + \frac{\phi''_0}{\phi'_0}]} \right\} \times \sum_{j=0}^{+\infty} \mu_{(j)}^2 \gamma_j(\varphi_1) \gamma_j(\varphi_2). \quad (16)$$

The second form of the normalization condition in Eq. (15) is useful in computational work, since it does not rely on knowledge of the eigenvalues. Details are given in Appendix A.

Because of the Neumann boundary conditions ($\partial_\varphi \psi_n$) = 0 and the simplicity of Eq. (7), there is always a massless

⁷The completeness of the solutions to the scalar Sturm-Liouville problem in Eq. (12) and the positivity of the scalar mass-squared eigenvalues $\mu_{(i)}^2$ are only assured if the coefficients of the δ -function terms are *non-negative* [33,34]. We will assume that the brane potentials and background field are such as to satisfy this condition (as they do in the stiff-wall limit we use later). Physically this constraint is more easily understood in the analogous case of the modes of a string: in that case the δ -functions can be understood as representing point masses which can freely move at the boundary of the string, and the coefficients are proportional to these masses and must therefore be positive for stability.

⁸Because $\mu_{(i)}^2 > 0$ in the GW model, this normalization choice (albeit unusual) is consistent. Taking the unstabilized limit $\phi'_0 \rightarrow 0$ [in which $\mu_{(0)} \rightarrow 0$ and all other scalar states decouple; see Eq (4)], however, requires care. This limit is discussed in the context of the “flat-stabilized” model in [16].

spin-2 mode (with a wave function constant in φ) in the tensor tower. This is not the case for the scalar tower. Due to the nonconstant potential of the background scalar, along with its vacuum expectation value, the lightest spin-0 state (identified as the radion with a wave function γ_0) acquires a mass $\mu_{(0)}$.

IV. MASSIVE SPIN-2 COUPLINGS, SCATTERING AMPLITUDES, AND SUM RULES

As discussed extensively in the literature (see, for example, [7,8], and references therein), phenomenological calculations incorporating massive spin-2 states often generate matrix element diagrams which exhibit anomalous high-energy behavior. Extradimensional models of gravity possess an underlying 5D diffeomorphism invariance that ensures their amplitudes are well behaved. That is, any overall bad high-energy growth necessarily signals the omission of additional important physics. Such omissions can produce erroneous phenomenological results.⁹

In our previous work we analyzed the diagrams which contribute to $2 \rightarrow 2$ massive spin-2 mode scattering for several variants of the Randall-Sundrum I model; within each of these analyses, we found there exist individual diagrams which diverge as fast as $\mathcal{O}(s^5)$ at high energies (s being the usual Mandelstam parameter) and that cancellations between diagrams ensure the total matrix element only grows as fast as $\mathcal{O}(s)$ [9–12]. This genuine $\mathcal{O}(s)$ growth is important because it ensures the 4D effective theory breaks down at an energy scale consistent with the physics of the underlying extradimensional theory. The central purpose of this paper is to verify the various coupling relations and sum rules [16] required to ensure these cancellations in a general Goldberger-Wise-stabilized Randall-Sundrum I model, report how most of these rules may be proved in full generality, and (in the next section) numerically demonstrate leading $\mathcal{O}(s)$ growth of the matrix element to second-order in a solvable warped stabilized model.

In this section we review the definitions of the KK mode couplings relevant to the scattering computations (Sec. IV A), describe the sum rules which apply to these couplings, and show how they are related to the physics of the Goldberger-Wise model (Sec. IV B), and provide a brief summary of the sum rules we numerically verify (Sec. IV C).

A. Scattering amplitudes and couplings

The tree-level diagrams relevant to the aforementioned $2 \rightarrow 2$ matrix element are shown in Fig. 1; details can be

⁹Following our previous computations [9–11] it was shown in [22] that in freeze-out computations with KK-graviton portal dark matter scenarios exhibit amplitudes that grow no faster than $\mathcal{O}(s)$ in an unstabilized model. The stabilized and phenomenologically relevant RS1 model is significantly more difficult to calculate and will be presented in a future work.




FIG. 1. Matrix element diagrams contributing to $n, n \rightarrow n, n$ massive spin-2 KK boson scattering. Here n refers to the KK mode number of the external state. The intermediate states x include a massive radion, the graviton, a tower of massive spin-2 KK bosons, and a tower of GW scalars.

found in [9–12,16]. The general strategy employed in this paper follows that of the unstabilized case. Briefly, the total matrix element \mathcal{M} involves a contact diagram (\mathcal{M}_c) as well as infinite sums over the diagrams ($\mathcal{M}_{j,X}$, $\mathcal{M}_{(i),X}$) describing $X = s, t$, and u -channel exchanges of intermediate spin-2 states and spin-0 states,

$$\mathcal{M} = \mathcal{M}_c + \sum_{X \in \{s,t,u\}} \left[\sum_{j=0}^{+\infty} \mathcal{M}_{j,X} + \sum_{i=0}^{+\infty} \mathcal{M}_{(i),X} \right],$$

where in general KK numbers within parentheses [the (i)] refer to those of the spin-0 states, while those without parentheses (the j) reference the spin-2 states. We expand the matrix element \mathcal{M} in a Taylor series in s as [11]

$$\mathcal{M}(s, \theta) = \sum_{\sigma \in \frac{1}{2}\mathbb{Z}} \overline{\mathcal{M}}^{(\sigma)}(\theta) \cdot s^\sigma \quad (17)$$

and isolate the kinematic factors and the couplings. Generally, 5D diffeomorphism demands that all coefficients of s^σ with $\sigma > 1$ in this expansion must vanish. At present, because we consider the process involving helicity-zero external states, half-integer values of σ automatically vanish. Demanding $\mathcal{O}(s)$ growth at most in this matrix element necessitates relationships between various masses and coupling structures in the theory.

The couplings present in each of these diagrams come from wave function overlap integrals attained following Kaluza-Klein decomposition of the fields in the Lagrangian; this procedure of attaining a 4D effective theory from a Lagrangian such as Eq. (1) via KK decomposition is explained in detail in Refs. [11,12].

- (i) The contact diagram \mathcal{M}_c involves the 4-point massive spin-2 vertex and contributes the following wave function overlap integrals:

$$a_{klmn} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi e^{-2A} \psi_k \psi_l \psi_m \psi_n,$$

$$b_{k'l'mn} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi e^{-4A} (\partial_\varphi \psi_k) (\partial_\varphi \psi_l) \psi_m \psi_n.$$

As shown explicitly in [9–11,16] and references therein, and can be argued from general power counting arguments originating from external

polarization and the tensor structures, the helicity-zero contribution to the \mathcal{M}_c diagrams diverge as fast as $\mathcal{O}(s^5)$.

- (ii) The spin-2 mediated diagrams $\mathcal{M}_{j,X}$ involve 3-point spin-2 vertices and contribute the following wave function overlap integrals:

$$a_{lmn} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi e^{-2A} \psi_l \psi_m \psi_n,$$

$$b_{l'm'n} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi e^{-4A} (\partial_\varphi \psi_l) (\partial_\varphi \psi_m) \psi_n.$$

Just as the contact diagrams [9–11,16], the helicity-zero contribution to the sum of the $\mathcal{M}_{j,X}$ diagrams diverges as $\mathcal{O}(s^5)$.

- (iii) The spin-0 mediated diagrams $\mathcal{M}_{(i),X}$ involve 3-point scalar-(spin-2)-(spin-2) couplings and contribute the following wave function overlap integrals:

$$a_{l'm'(n)} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi e^{-2A} (\partial_\varphi \psi_l) (\partial_\varphi \psi_m) \gamma_n.$$

Due to the structure of the Lagrangian, no corresponding “ $b_{l'm'(n)}$ ” or “ $b_{lm'(n)}$ ” is generated. In this case [9–11,16] the scalar-exchange diagrams grow more slowly, where each $\mathcal{M}_{(i),X}$ diverges like $\mathcal{O}(s^3)$.

In general, (i) couplings labeled with an “ a ” have an e^{-2A} weight factor, whereas those labeled with a “ b ” involve e^{-4A} ; (ii) the subscript KK indices indicate the relevant wave functions to include in each integral (remembering that parentheses indicate scalar modes); and (iii) a subscript KK index with a prime denotes that the corresponding mode number’s wave function should be differentiated with respect to the extradimensional coordinate φ .

We can reduce the number of coupling integrals present by using the properties of the KK wave functions; namely, the mode equation and completeness relations, Eqs. (7) and (9). For example, in prior work [11,12] we showed how the spin-2 mode equation (7) and the corresponding completeness relation (9) relate some of the a and b couplings:

$$b_{l'm'n} = \frac{1}{2} [\mu_l^2 + \mu_m^2 - \mu_n^2] a_{lmn}, \quad b_{n'n'nn} = \frac{\mu_n^2}{3} a_{nnnn}. \quad (18)$$

We use these relations and eliminate all b -type overlap integrals in favor of a -type integrals. Doing so, we may write the sum rules entirely using a -type overlap integrals.

B. Sum rules ensuring consistency of scattering amplitudes

By requiring the scattering amplitude to grow no faster than $\mathcal{O}(s)$ in the GW model [16], we previously determined the following general sum rules should be satisfied:

$$\sum_{j=0}^{\infty} a_{nnj}^2 = a_{nnnn}, \quad (19)$$

$$\sum_{j=0}^{\infty} \mu_j^2 a_{nnj}^2 = \frac{4}{3} \mu_n^2 a_{nnnn}, \quad (20)$$

$$\sum_{j=0}^{\infty} \mu_j^4 a_{nnj}^2 = \frac{4}{15} \mu_n^4 (4a_{nnnn} - 3a_{nn0}^2) + \frac{36}{5} \sum_{i=0}^{\infty} a_{n'n'(i)}^2, \quad (21)$$

$$\sum_{j=0}^{\infty} \mu_j^6 a_{nnj}^2 = -4\mu_n^6 a_{nn0}^2 + 9 \sum_{i=0}^{\infty} (4\mu_n^2 - \mu_{(i)}^2) a_{n'n'(i)}^2. \quad (22)$$

The first two sum rules, Eqs. (19) and (20), ensure that the contributions to the scattering amplitudes growing like $\mathcal{O}(s^5)$ and $\mathcal{O}(s^4)$ vanish. They follow directly from the Sturm-Liouville form of the spin-2 KK mode equation (7) and the corresponding completeness relation (9)—so the proofs given in [10–12] apply to any model producing a geometry defined by a warp function $A(y)$. However, the sum rules in Eqs. (21) and (22), which ensure cancellation of the contributions to the amplitude growing like $\mathcal{O}(s^3)$ and $\mathcal{O}(s^2)$, involve the scalar tower present in the GW model.

By combining the last two sum rules, Eqs. (21) and (22), to eliminate the common sum $\sum_i a_{n'n'(i)}^2$, we find a mixed rule:

$$\sum_{j=0}^{\infty} [5\mu_n^2 - \mu_j^2] \mu_j^4 a_{nnj}^2 = \frac{16}{3} \mu_n^6 a_{nnnn} + 9 \sum_{i=0}^{\infty} \mu_{(i)}^2 a_{n'n'(i)}^2. \quad (23)$$

As we show now, this combined sum rule can be expressed in a way that depends only on the spin-2 wave functions. The only scalar tower sum $\sum_{i=0}^{\infty} \mu_{(i)}^2 a_{n'n'(i)}^2$ remaining in this particular combination of the $\mathcal{O}(s^3)$ and $\mathcal{O}(s^2)$ sum rules can be eliminated using the spin-0 completeness relation Eq. (16). Since the spin-2 wave functions satisfy Neumann boundary conditions, $(\partial_\varphi \psi_n) = 0$ at $\varphi = 0$ and π , we find

$$\sum_{i=0}^{\infty} \mu_{(i)}^2 a_{n'n'(i)}^2 = \frac{1}{6} \left\{ \int d\varphi (\partial_\varphi \phi_0)^2 e^{-8A} (\partial_\varphi \psi_n)^4 \right\}. \quad (24)$$

Hence the combination of sum rules in Eq. (23) does not depend on the explicit form of the scalar wave functions $\gamma_i(\varphi)$, but only on the spin-2 wave functions $\psi_n(\varphi)$, the exponentiated warp factor $e^{A(\varphi)}$, and (the derivative of) the background scalar field ϕ_0 .

In Appendix B 5 a we show, by applying only the *spin-2* mode equation (7), Neumann boundary conditions, and completeness relations (9), that

$$\sum_{j=0}^{\infty} [5\mu_n^2 - \mu_j^2] \mu_j^4 a_{nnj}^2 = \frac{16}{3} \mu_n^6 a_{nnnn} + 18 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi (\partial_\varphi^2 A) e^{-8A} (\partial_\varphi \psi_n)^4 \right\}. \quad (25)$$

In the GW model we have the Einstein equation [via Eq. (10)] $(\partial_\varphi^2 A) = (\partial_\varphi \phi_0)^2 / 12 + \sum_i V_i r_c \delta_i(\varphi) / 3$. The Dirac delta terms vanish because $(\partial_\varphi \psi_n) = 0$ at the boundaries, and hence

$$\sum_{j=0}^{\infty} [5\mu_n^2 - \mu_j^2] \mu_j^4 a_{nnj}^2 = \frac{16}{3} \mu_n^6 a_{nnnn} + \frac{3}{2} \left\{ \int d\varphi (\partial_\varphi \phi_0)^2 e^{-8A} (\partial_\varphi \psi_n)^4 \right\}. \quad (26)$$

Applying Eq. (24) we immediately obtain Eq. (23). Hence Eq. (23) depends nontrivially on the dynamics of the GW model—in particular, on the Einstein equations for the warp factor and on the scalar completeness relation which follows from the mode equation and the scalar mode normalization condition.

Using the spin-2 completeness relations and the relations between the a and b couplings in Eq. (18) and the sum rule in Eq. (19), we find

$$\sum_{j=0}^{\infty} \mu_j^4 a_{nnj}^2 = 4c_{n'n'n'n'} + \frac{4}{3} \mu_n^4 a_{nnnn}. \quad (27)$$

Here we have defined the quantity

$$c_{n'n'n'n'} = \sum_{j=0}^{\infty} b_{n'n'j}^2 \equiv \frac{1}{\pi} \int d\varphi e^{-6A} (\partial_\varphi \psi_n)^4, \quad (28)$$

which depends only on the spin-2 wave functions. Plugging this into Eq. (26) we find

$$\sum_{j=0}^{\infty} \mu_j^6 a_{nnj}^2 = \frac{4}{3} \mu_n^6 a_{nnnn} + 20\mu_n^2 c_{n'n'n'n'} - \frac{3}{2} \left\{ \int d\varphi (\partial_\varphi \phi_0)^2 e^{-8A} (\partial_\varphi \psi_n)^4 \right\}, \quad (29)$$

which depends only on the spin-2 wave functions and the background scalar-field configuration ϕ_0 .

Finally, having demonstrated that one linear combination of Eqs. (21) and (22) is determined entirely by the spin-2 sector of the GW model, we can also use Eq. (27) to isolate the contribution from the GW scalar sector

$$\sum_{i=0}^{\infty} a_{n'n'(i)}^2 = \frac{5}{9} c_{n'n'n'n'} + \frac{1}{9} \mu_n^4 a_{nn0}^2 + \frac{1}{27} \mu_n^4 a_{nnnn}, \quad (30)$$

which succinctly summarizes the necessary (but unproven) relationship between the scalar and spin-2 couplings (and hence wave functions) which must be satisfied in order for the spin-2 scattering amplitudes to grow no faster than $\mathcal{O}(s)$.

C. Sum-rule summary

In summary, the spin-2 coupling relations

$$\sum_{j=0} a_{nnj}^2 = a_{nnnn}, \quad (19 \text{ revisited})$$

$$\sum_{j=0} \mu_j^2 a_{nnj}^2 = \frac{4}{3} \mu_n^2 a_{nnnn}, \quad (20 \text{ revisited})$$

$$\sum_{j=0}^{+\infty} \mu_j^4 a_{nnj}^2 = 4c_{n'n'n'n'} + \frac{4}{3} \mu_n^4 a_{nnnn}, \quad (27 \text{ revisited})$$

$$\begin{aligned} \sum_{j=0}^{+\infty} \mu_j^6 a_{nnj}^2 &= \frac{4}{3} \mu_n^6 a_{nnnn} + 20\mu_n^2 c_{n'n'n'n'} \\ &\quad - \frac{3}{2} \left\{ \int d\varphi (\partial_\varphi \phi_0)^2 e^{-8\Lambda} (\partial_\varphi \psi_n)^4 \right\} \end{aligned} \quad (29 \text{ revisited})$$

follow from the form of the spin-2 mode equations, the spin-2 wave function completeness, the Einstein equations for the warp factor, and the scalar completeness relation which follows from the scalar mode equation and the scalar mode normalization condition. Analytic derivations of all of these relations are given above or in Appendix B 5 a. With respect to guaranteeing that the massive spin-2 scattering amplitudes grow no faster than $\mathcal{O}(s)$, these relations show that the first two sum rules Eqs. (19) and (20) and one combination of the sum rules in Eqs. (21) and (22) are always satisfied.¹⁰

Separately, the sum rule

$$\sum_{i=0}^{+\infty} a_{n'n'(i)}^2 = \frac{5}{9} c_{n'n'n'n'} + \frac{1}{9} \mu_n^4 a_{nn0}^2 + \frac{1}{27} \mu_n^4 a_{nnnn}, \quad (30 \text{ revisited})$$

which depends on the spin-0 GW scalar couplings must also be satisfied in order for the spin-2 scattering amplitudes to grow no faster than $\mathcal{O}(s)$. With the methods discussed here, we have been unable to prove analytically that this scalar sum rule is satisfied in general.¹¹

¹⁰Generalizations of the proven rules (and their proofs) as well as additional unproven rules necessary for $\mathcal{O}(s)$ growth of the inelastic amplitude $(k, l) \rightarrow (m, n)$ are provided in Appendix B.

¹¹While this paper was under review, an analytic proof of this remaining sum rule has been developed [35] by reframing the problem in conformal coordinates to reveal a hidden $N = 2$ supersymmetry structure of the mode equations.

In [16], we demonstrated that these sum rules in Eqs. (19)–(22) were satisfied in the “flat-stabilized” model—a slight deformation of a flat extradimensional model in which the radion is massive and the size of the extra dimension is stable. We now demonstrate numerically that these relations are satisfied in the presence of the warping required to produce the hierarchy between the weak and Planck scales in the Randall-Sundrum model.

V. PERTURBATIVE ANALYSIS OF A WARPED STABILIZED MODEL

In the original formulation of the RS1 model [3,4], Randall and Sundrum constructed a consistent solution to the Einstein field equations by choosing a warp factor $A(\varphi) = kr_c |\varphi|$ sourced by brane and bulk cosmological constants. Once the Goldberger-Wise mechanism is implemented, such a simple functional form becomes unavailable: the background geometry is augmented, the Einstein field equations are changed, and the warp factor $A(\varphi)$ is made more complicated. The background Einstein field equations of the stabilized model [Eqs. (A26), (A27), (A29), and (A30)] are coupled nonlinear equations with respect to $A(\varphi)$ that depend on the derivative of the scalar background $\phi'_0(\varphi)$ and are generally difficult to solve. DeWolfe, Freedman, Gubser, and Karch have constructed a specific class of exactly solvable potentials (the DFGK model) [36] which make calculations in the stabilized RS1 model feasible.

In this section, we review the DFGK class of solutions to set notation, and we detail a specific DFGK model which enables perturbative expansion around the (solved) warped unstabilized RS1 model [16]. Subsequently, for the physical spin-0 and spin-2 towers, we perturbatively compute the wave functions and masses using the Sturm-Liouville equations described in Sec. III. Finally, we demonstrate that the sum rules defined in Sec. IV are numerically satisfied at second order in the expansion parameter, the lowest order necessary to generate a nonzero radion mass.

A. The DFGK model

A key strategy employed in the DFGK model [36] is the introduction of a superpotential-inspired function $W[\hat{\phi}]$ which is used to simplify the stabilized model’s background field equations [Eqs. (A26), (A27), (A29), and (A30)]. In particular, it is assumed that the scalar bulk and brane potentials may be parametrized as

$$V r_c^2 = \frac{1}{8} \left(\frac{dW}{d\hat{\phi}} \right)^2 - \frac{W^2}{24}, \quad (31)$$

$$\begin{aligned} V_1 r_c &= +\frac{W}{2} + \beta_1^2 [\hat{\phi}(\varphi) - \phi_1]^2, \\ V_2 r_c &= -\frac{W}{2} + \beta_2^2 [\hat{\phi}(\varphi) - \phi_2]^2. \end{aligned} \quad (32)$$

In this case, the background scalar and Einstein equations are solved if

$$(\partial_\varphi A) = \frac{W}{12} \Big|_{\hat{\phi}=\phi_0} \text{sign}(\varphi), \quad (\partial_\varphi \phi_0) = \frac{dW}{d\hat{\phi}} \Big|_{\hat{\phi}=\phi_0} \text{sign}(\varphi), \quad (33)$$

where $\phi_1 \equiv \hat{\phi}(0)$ and $\phi_2 \equiv \hat{\phi}(\pi)$. Reference [36] introduces a convenient $W[\hat{\phi}]$ with the following specific form¹²:

$$W[\hat{\phi}(\varphi)] = 12kr_c - \frac{1}{2}\hat{\phi}(\varphi)^2 ur_c. \quad (34)$$

Plugging this into Eq. (33) we find solutions for the bulk scalar vacuum and the warp factor:

$$\phi_0(\varphi) = \phi_1 e^{-ur_c|\varphi|}, \quad (35)$$

$$A(\varphi) = kr_c|\varphi| + \frac{1}{48}\phi_1^2[e^{-2ur_c|\varphi|} - 1]. \quad (36)$$

In the limit that the parameter u vanishes, this reduces to the usual unstabilized RS1, with the bulk field acquiring a constant vacuum expectation value. The parameters u , ϕ_1 , and ϕ_2 are related according to

$$ur_c = \frac{1}{\pi} \log \frac{\phi_1}{\phi_2}. \quad (37)$$

We next define the small- u limit carefully to facilitate a perturbative analysis.

B. The perturbative DFGK model

The forms of $\phi_0(\varphi)$ and $A(\varphi)$ in Eqs. (35) and (36) are useful for solving the background equations, but it remains difficult to find general analytic solutions (i.e., the KK wave functions and masses) to the differential equations defined in Sec. III. Therefore, we define a limit of the model so that we may perturbatively expand ϕ_0 and A around the unstabilized background (for which analytic solutions are well known), take the stiff wall limit [with $\check{V}_{1,2} \rightarrow \infty$ so that the scalar boundary conditions in Eq. (13) reduce to Neumann conditions at $\varphi = 0, \pi$], and develop solutions for the Sturm-Liouville problems order-by-order in perturbation theory.¹³ Details of the perturbation theory can be found in Appendix C. Here, we now proceed to

introduce the effective warp parameter \tilde{k} and perturbation parameter ϵ [16].

Suppose we series expand $A(\varphi)$ with respect to the unitless quantity ur_c . Expanding around $ur_c = 0$ yields

$$A(\varphi) = kr_c|\varphi| - \left[\frac{\phi_1^2}{24}|\varphi|\right](ur_c) + \left[\frac{\phi_1^2}{24}|\varphi|^2\right](ur_c)^2 + \mathcal{O}((ur_c)^3) \quad (38)$$

$$= \left[k - \frac{\phi_1^2}{24}\right]r_c|\varphi| + \left[\frac{\phi_1^2(ur_c)^2}{24}\right]|\varphi|^2 + \mathcal{O}((ur_c)^3). \quad (39)$$

The first term in the second line demonstrates that, when ur_c is sufficiently small, the stabilized model is a small deformation of an unstabilized Randall-Sundrum I model [3,4]. Because we intend to work in the $ur_c \rightarrow 0$ limit, we will eliminate the actual warp parameter k in favor of the effective warp parameter,

$$\tilde{k} \equiv k - \phi_1^2 u/24, \quad (40)$$

that applies in that limit.¹⁴

To simplify various factors that would otherwise be present in multiple equations, we will also replace ur_c (and its role as our expansion parameter) with the rescaled dimensionless perturbative parameter $\epsilon \equiv \phi_1(ur_c)/\sqrt{24}$. This definition of ϵ simplifies $A(\varphi)$ at $\mathcal{O}(\epsilon^2)$,

$$A(\varphi) = \tilde{k}r_c|\varphi| + \frac{\phi_1^2}{48} \left[\exp\left(-\frac{4\sqrt{6}}{\phi_1}\epsilon|\varphi|\right) - 1 \right] + \frac{\phi_1}{2\sqrt{6}}\epsilon|\varphi| \quad (41)$$

$$= \tilde{k}r_c|\varphi| + \epsilon^2|\varphi|^2 + \mathcal{O}(\epsilon^3), \quad (42)$$

and yields, to all orders in ϵ ,

$$W[\hat{\phi}(\varphi)] = 12\tilde{k}r_c + \frac{\sqrt{6}}{\phi_1}[\phi_1^2 - \hat{\phi}(\varphi)^2]\epsilon, \quad (43)$$

$$\phi_0(\varphi) = \phi_1 \exp\left(-\frac{2\sqrt{6}}{\phi_1}\epsilon|\varphi|\right) = \phi_1 \exp(-\alpha\epsilon|\varphi|), \quad (44)$$

where $\alpha = \frac{2\sqrt{6}}{\phi_1}$. It is the form of the warp factor shown in Eq. (41) that we use in subsequent perturbative computations.

¹⁴In [16] we discussed the properties of the ‘‘flat-stabilized model’’ with $\tilde{k} = 0$. DS: Here we discuss the phenomenologically relevant limit of large $\tilde{k}r_c$.

¹²Here u is a parameter, and not the \hat{u} field of Eq. (A3).

¹³Specifically, we will solve this problem numerically in two limits, both of which are phenomenologically relevant and interesting. One corresponds to large values of $\tilde{k}r_c$ that connects the 4D Planck scale to TeV scale physics in the context of RS models. The second limit corresponds to small values of ur_c that give rise to small values of the radion mass and allow for us to solve the relevant equations perturbatively.

1. Spin-2 wave functions and masses in perturbation theory

To order $\mathcal{O}(\epsilon^2)$, the spin-2 mode equation (7) becomes¹⁵

$$\partial_\varphi [e^{-4\tilde{k}r_c\varphi - 4\epsilon^2\varphi^2} \partial_\varphi \psi_n] = -\mu_n^2 e^{-2\tilde{k}r_c\varphi - 2\epsilon^2\varphi^2} \psi_n. \quad (45)$$

The form of this equation, along with the Neumann boundary conditions $\partial_\varphi \psi_n = 0$ at $\varphi = 0, \pi$, ensures that the zero mode (the graviton) is massless and has a wave function which is constant in φ . Expanding this equation in powers of ϵ^2 , we solve Eq. (7) using perturbation theory as described in Appendix C. The perturbative expressions for the spin-2 wave functions and masses are quite lengthy, and to simplify them, we restrict ourselves to the limit when $\tilde{k}r_c$ is large. The expressions for the spin-2 mass and wave functions are given to order ϵ^2 in the large- $\tilde{k}r_c$ limit in Appendix C 2.

To illustrate the effects of the geometry on the spin-2 masses, we calculate the masses for Eq. (7) using the Wentzel–Kramers–Brillouin approximation. The asymptotic formula for the masses is given by¹⁶

$$\mu_n = m_n r_c = \frac{n\pi}{l}, \quad l = \int_0^\pi d\varphi e^A. \quad (46)$$

The equations above show how the eigenvalues are positive and form an infinite tower of states. Using the form of $A(\varphi)$ from Eq. (36), we find the mass of spin-2 KK modes for a large mode number to be

$$\mu_n|_{n \gg 1} = \frac{(n\pi)\tilde{k}r_c}{e^{\pi\tilde{k}r_c} - 1} - (n\pi) \frac{e^{\pi\tilde{k}r_c} [\pi\tilde{k}r_c (\pi\tilde{k}r_c - 2) + 2] - 2}{\tilde{k}r_c (e^{\pi\tilde{k}r_c} - 1)^2} \epsilon^2 + \mathcal{O}(\epsilon^3). \quad (47)$$

In the limit when $\tilde{k}r_c \gg 1$, which is the phenomenologically interesting limit, this expression further simplifies to

$$\mu_n|_{n \gg 1} \simeq (n\pi)\tilde{k}r_c e^{-\pi\tilde{k}r_c} \left\{ 1 - \frac{[\pi\tilde{k}r_c (\pi\tilde{k}r_c - 2) + 2]}{\tilde{k}^2 r_c^2} \epsilon^2 \right\}. \quad (48)$$

We can see how the effect of the on the geometry from the bulk scalar field reduces the masses of the massive spin-2

¹⁵Here, and in subsequent equations related to the perturbative model in this paper, we expand $A(\varphi)$ to $\mathcal{O}(\epsilon^2)$ using Eq. (41), but we do not record the expansion of the exponents in powers of ϵ to retain succinctness of expressions when writing. However, these exponents are actually expanded during our calculations, e.g., when we define the perturbed differential equations in (C4) and (C5).

¹⁶There are $\mathcal{O}(1)$ corrections to $\mu_n^2 \ell^2$ that do not grow with n , due to the Neumann boundary conditions. These effects can be included, but do not affect the analysis given here.




FIG. 2. We show for each spin-2 KK mode n , the ratio of mass correction to leading order mass times ϵ^2 (dots). The mass correction $\delta\mu_n^2$ is calculated in the large $\tilde{k}r_c$ limit to order ϵ^2 , as shown in Eq. (C29). The line represents the asymptotic form of the expression given in Eq. (49), which is valid for the large spin-2 KK number.

modes. Further, the square of the ratio of the correction to the mass ($\delta\mu_n$) to the leading order mass ($\mu_n^{(0)}$) is

$$\frac{\delta\mu_n^2}{(\mu_n^{(0)})^2} \Big|_{n \gg 1; \tilde{k}r_c \gg 1} \simeq \left[-2\pi^2 + \frac{4\pi}{\tilde{k}r_c} - \frac{4}{\tilde{k}^2 r_c^2} + \dots \right] \epsilon^2. \quad (49)$$

We can therefore conclude that the perturbation theory we use is valid when $|\epsilon| \ll 1/(\sqrt{2}\pi)$. In Fig. 2, we compare the full expression for the mass corrections [given in Eq. (C29) of Appendix C] to the asymptotic form shown here. Here we see that the full form of the mass, represented by the blue dots, approaches the asymptotic value, represented by the bold red line.

While the asymptotic formula is simple and provides a convenient cross-check of our calculations, it is insufficient for our present purposes. To demonstrate cancellations and verify sum rules, we need to be able to evaluate the $\mathcal{O}(\epsilon^2)$ wave functions and masses without approximation. We provide exact expressions to order ϵ^2 in the perturbation theory within Appendix C 2. These expressions are consistent with the large $\tilde{k}r_c$ limit results. Full expressions (which are valid for arbitrary values of $\tilde{k}r_c$) are provided as supplementary *Mathematica* files [28].

2. Spin-0 wave functions and masses in perturbation theory

Equation (44) implies that

$$(\phi'_0)^2 = 24\epsilon^2 e^{-2\alpha\epsilon\varphi} \quad (50)$$

in the bulk. To simplify our analysis, we consider the perturbative solution for the scalar tower in the “stiff-wall”

limit, $\check{V}_{1,2} \rightarrow +\infty$, so that Eq. (13) reduces to Neumann conditions

$$\partial_\varphi \gamma_i|_{\varphi=0,\pi} = 0. \quad (51)$$

Using this and the expansion of $A(\varphi)$ to order $\mathcal{O}(\epsilon^2)$ from Eq. (41), the spin-0 mode equation (12) becomes (after multiplying through by $24\epsilon^2$)

$$\begin{aligned} \partial_\varphi [e^{(2\tilde{k}r_c\varphi + 2a\epsilon\varphi + 2\epsilon^2\varphi^2)} \partial_\varphi \gamma_i] - 4\epsilon^2 e^{(2\tilde{k}r_c\varphi + 2\epsilon^2\varphi^2)} \gamma_i \\ = -\mu_i^2 e^{(4\tilde{k}r_c\varphi + 2a\epsilon\varphi + 4\epsilon^2\varphi^2)} \gamma_i, \end{aligned} \quad (52)$$

while the normalization conditions to this order are

$$\begin{aligned} \delta_{mn} = \frac{1}{4\pi\epsilon^2} \int_{-\pi}^{+\pi} d\varphi [e^{(2\tilde{k}r_c\varphi + 2a\epsilon\varphi + 2\epsilon^2\varphi^2)} \gamma'_m \gamma'_n \\ + 4\epsilon^2 e^{(2\tilde{k}r_c\varphi + 2\epsilon^2\varphi^2)\varphi} \gamma_m \gamma_n]. \end{aligned} \quad (53)$$

Expanding the spin-0 mode equation (52) in powers of ϵ , we solve the differential Eq. (12) using the perturbation theory described in Appendix C. In particular, using Eq. (C12), we find that the radion [identified as the zero mode of the KK expansion in Eq. (11)] acquires a mass-squared at order ϵ^2 in perturbation theory

$$\mu_{(0)}^2 = \frac{8\epsilon^2}{1 + e^{2\pi\tilde{k}r_c}} + \mathcal{O}(\epsilon^3), \quad (54)$$

while the radion wave function evaluated to order ϵ^2 is

$$\begin{aligned} \gamma_{(0)} = \sqrt{\frac{\tilde{k}r_c\pi}{e^{2\tilde{k}r_c\pi} - 1}} + C(\tilde{k}r_c) \{2\tilde{k}r_c\varphi \\ + \operatorname{sech}(\pi\tilde{k}r_c) \sinh[\tilde{k}r_c(\pi - 2\varphi)] \\ - \tanh(\pi\tilde{k}r_c)\} \epsilon^2 + \mathcal{O}(\epsilon^3). \end{aligned} \quad (55)$$

Here $C(\tilde{k}r_c)$ is determined through normalization; the normalized radion wave function is provided in Appendix C in Eq. (C34). In the limit where ϵ vanishes, the gravitational degrees of freedom and the bulk scalar cease mixing. This results in a massless radion, an unstabilized extra dimension, and an entirely separate tower of scalar states. Note that the leading order radion wave function is flat, signaling that it is massless at that order.

The GW scalar wave function to order ϵ is given in Eq. (C36). As expected, the wave functions are composed of Bessel functions. Due to the normalization condition in Eq. (53), the massive GW scalar mode wave functions have no ϵ^0 terms and start at order ϵ . Since the sum rules in Eq. (19) through Eq. (23) have only products of GW scalar wave functions, when trying to verify them to order ϵ^2 , we only need expressions of the wave functions to order ϵ .

We therefore only provide expressions for the GW scalar wave function and masses to leading order.¹⁷

C. Numerical verification of sum rules

We now verify the sum rules summarized in Sec. IV C for the warped-stabilized model using our perturbative computations. We can substitute expressions for wave functions and masses calculated in the DFGK model and provided in Appendix C 2 and evaluate the overlap integrals numerically. The sum rules in Eqs. (19), (20), (27), (29), and (30) can thereby be evaluated order-by-order in ϵ . We know that the sum rules are satisfied for the unstabilized RS model [11], and therefore these expressions agree to leading order (ϵ^0). Using our perturbative expressions, we verify here that the sum rules are satisfied to leading nontrivial order, $\mathcal{O}(\epsilon^2)$. Equivalently, we show that the $\mathcal{O}(\epsilon^2)$ contributions on the left- and right-hand sides of Eqs. (19), (20), (27), (29), and (30) agree.

Note that the left-hand sides of these expressions are given as infinite sums over different overlap integrals. It is therefore not possible to perform the entire sum. Instead, we perform the sum up a ‘‘cutoff’’ KK number and show that the relative error in the $\mathcal{O}(\epsilon^2)$ contributions to the left-hand side (LHS) converge to the $\mathcal{O}(\epsilon^2)$ contributions to the right-hand side (RHS) of each expression as the number of KK modes included in the sum increases. For example, to numerically verify Eq. (19), we take the coefficient of the ϵ^2 piece in $\sum_j a_{nnj}$, referred to as ΔLHS and divide by the coefficient of the ϵ^2 piece in a_{nnnn} , referred to as ΔRHS . We examine how the relative error $\log_{10} |1 - \Delta\text{LHS}/\Delta\text{RHS}|$ scales as we increase the number of KK modes in the sum of ΔLHS .

1. Spin-2 sum rules and completeness

We begin with Eqs. (19), (20), (27), and (29). The result of this exercise is shown in Fig. 3, in the case $n = 1$ (e.g., for elastic scattering of spin-2 modes at KK level 1) and for $\tilde{k}r_c = 40/\pi = 12.73$. We see that each of the series converges nicely with the relative error reducing with the addition of terms to the sum on the LHS of the equation. As described in detail in Sec. IV, Eqs. (19), (20), and (27) can be proven directly using the completeness properties of the solutions of the spin-2 mode equation (7). Their numerical verification demonstrates that accuracy of our perturbative analysis. The first two of these equations demonstrate that the $\mathcal{O}(s^5)$ and $\mathcal{O}(s^4)$ contributions to helicity-0 spin-2 elastic scattering vanish to this order in perturbation theory.

¹⁷Note that the limit $\tilde{k}r_c = 0$, dubbed as the flat stabilized model, was studied previously in our work [16], where we show that the sum rules required for the scattering amplitudes to grow only as $\mathcal{O}(s)$ were satisfied to $\mathcal{O}(\epsilon^2)$.




FIG. 3. Verification of sum rules [Eqs. (19), (20), (27), and (29), reproduced at the bottom of the diagram] for elastic scattering of KK mode number one ($n = 1$). The index j , shown on the x -axis, is the number of spin-2 modes included in the sum of the left-hand side of the sum rules. The y -axis indicates the log of relative error ($\log_{10}(1 - \Delta(\text{LHS})/\Delta(\text{RHS}))$) between the $\mathcal{O}(\epsilon^2)$ corrections to the LHS and the RHS of the relevant equations. This has been evaluated in the large $\tilde{k}r_c$ limit.

We also see that the sum rule in Eq. (29) converges as well. As discussed above, Eq. (29) depends nontrivially on the dynamics of the Goldberger-Wise model implemented here—in particular on the Einstein equation coupling the scalar potential to the curvature of the extra dimension Eq. (10) and on the completeness conditions of the spin-0 modes Eq. (16). As discussed in Sec. IV C, this verifies that one linear combination of Eqs. (21) and (22) is also satisfied.

Finally, in all cases we see that the series only converges rapidly once we have included the $j = 2$ term. This is because the overlap integral defining the coupling between two spin-2 level-1 states and a spin-2 state at level j is largest for $j = 2$ —which can be understood as a remnant of the “discrete” KK momentum conservation, which would be present in a flat extra dimension.

2. The spin-0 sum rule

Finally, we examine the sum rule in Eq. (30) for which we have no analytic proof, and which depends on the couplings of the individual spin-0 states to the massive spin-2 KK modes. The result of this exercise is shown in Fig. 4, in the case $n = 1, 5, 11$ (e.g., for scalar-exchange contributions to elastic scattering of spin-2 modes at KK levels 1, 5, and 11) and for $\tilde{k}r_c = 12.73$. Again, what is plotted here is the agreement between the $\mathcal{O}(\epsilon^2)$ contributions to the left- and right-hand sides of Eq. (30)—which, at this order, depends explicitly on the forms of the scalar wave functions that solve Eq. (12) and which are subject to the normalization conditions of Eq. (15).




FIG. 4. Verification of the scalar sum rule in Eq. (30) for elastic scattering of spin-2 KK mode number $n = 1$ (red), $n = 5$ (blue), and $n = 11$ (green). Here n represents the KK mode number of the spin-2 particles in the external legs. On the x -axis we show the number of GW-scalar modes included in the sum (i) of the left-hand side of the sum rules in Eq. (30) versus the log of the relative error ($\log_{10}(1 - \Delta(\text{LHS})/\Delta(\text{RHS}))$) between the $\mathcal{O}(\epsilon^2)$ contributions to both sides of the relevant equation. This has been evaluated in the large $\tilde{k}r_c$ limit.

The nonzero difference between the left- and right-hand sides of the $\mathcal{O}(\epsilon^2)$ corrections to Eq. (30) at $i = 1$ demonstrate the need to include the tower of Goldberger-Wise scalar states for the spin-2 scattering amplitudes to have the proper high-energy behavior. Again we see that the largest single contribution to the $\mathcal{O}(\epsilon^2)$ corrections comes from the exchange of the GW scalar state whose mode number is twice that of the incoming particles— $i = 2, 10,$ and 20 , respectively, for incoming modes 1, 5, and 10 spin-2 states. However, the continued convergence when adding additional states is also clear and, formally, the entire tower is needed for the sum rule to be satisfied.

VI. CONCLUSION

In this paper we have presented a thorough analysis of the scattering of massive spin-2 Kaluza-Klein excitations in phenomenologically realistic models based on a warped geometry [3,4] stabilized via the Goldberger-Wise [17,18] mechanism. These results significantly extend the work presented in [9–11] on the unstabilized RS1 model and the results in [16] on the “flat-stabilized” model ($\tilde{k}r_c = 0$). We briefly recap our findings here:

- (i) We provided a complete and self-contained derivation of the mode expansions for the spin-2 and spin-0 states and their interactions. Generalizing the presentations in [24–26], our computations are given in de Donder gauge for massless gravitons—allowing us to consistently compute scattering amplitudes involving intermediate off-shell states.

- (ii) In previous work [16] we had demonstrated the extended sum-rule relationships between spin-2 and spin-0 modes, and their masses and couplings, which must be satisfied in order for elastic massive spin-2 KK scattering to grow no faster than $\mathcal{O}(s)$. Here, we have provided an analytic proof for one combination of these sum rules and showed its relation to both the Einstein and the scalar background field equations implementing the Goldberger-Wise dynamics and also to the properties of the mode equations for the physical scalar fields (fields which are admixtures of bulk scalar and gravitational modes in the original theory).
- (iii) We have provided, in Appendix B to this work, a complete list of the sum-rule relations which must be satisfied if all $2 \rightarrow 2$ massive spin-2 scattering amplitudes, elastic or inelastic, are to grow no faster than $\mathcal{O}(s)$ —completing the analyses begun in [9–12].
- (iv) Finally, using a version of the DFGK model [36] in which the Goldberger-Wise dynamics can be treated perturbatively [16], we have checked numerically that the sum rules which enforce the proper high-energy behavior of massive spin-2 scattering continue to be satisfied in the case of the large warping that would be required to produce the hierarchy between the weak and Planck scales. These numerical computations demonstrate that, in models with a massive radion, proper cancellation is achieved only after including the contributions from the tower of scalar states present in the Goldberger-Wise model.

In future work we will also explore the phenomenological consequences of the fact that all spin-2 scattering amplitudes in models of compactified gravity can grow no faster than $\mathcal{O}(s)$; specifically, we will study the implications for the computation of relic abundances of dark matter particles in KK graviton-portal theories and related theories.

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation under Grant No. PHY-1915147. The work of D. S. for this paper was supported in part by the University of Adelaide and the Australian Research Council through the Centre of Excellence for Dark Matter Particle Physics (CE200100008).

APPENDIX A: THE DERIVATION OF THE SCALAR KALUZA-KLEIN MODES

The canonical quadratic Lagrangian for the Kaluza-Klein modes in a Goldberger-Wise model has been derived in Refs. [24–26]. Here, for completeness and

as a guide to the interested reader, we present our derivation of the canonical quadratic Lagrangian. In addition, since our computation includes diagrams with off-shell gravitons, we are careful to derive our results in de Donder gauge.¹⁸ We include all the necessary details needed to reproduce our results: deriving the background equations of motion for a Goldberger-Wise-stabilized Randall-Sundrum I model, showing how the 5D scalar field \hat{r} and 5D tensor field \hat{h} decouple, motivating the gauge condition relating the fluctuation field \hat{f} and the 5D scalar field \hat{r} , describing how the linear equations of motion inspire the Kaluza-Klein decomposition of the 5D scalar field \hat{r} , and demonstrating that the 5D scalar field $\hat{r}(x, y)$ generates a tower of canonical spin-0 fields $\hat{r}^{(i)}(x)$ with masses $m_{(i)} \equiv \mu_{(i)}/r_c$.

1. The Lagrangian

The Goldberger-Wise-stabilized Randall-Sundrum I Lagrangian is constructed from several elements, including the spacetime metric. Focusing our attention on the spin-2 ($\hat{h}_{\mu\nu}$) and scalar (\hat{r}) fluctuations about a geometry determined by the warp factor $A(y)$, we use the following parametrization of the metric G_{MN} and its 4D projection \tilde{G}_{MN} :

$$[G_{MN}] = \begin{pmatrix} wg_{\mu\nu} & 0 \\ 0 & -v^2 \end{pmatrix}, \quad [\tilde{G}_{\mu\nu}] = wg_{\mu\nu}, \quad (\text{A1})$$

where (taking our parametrization from [32])

$$w = e^{-2[A(y)+\hat{u}(x,y)]}, \quad v = 1 + 2\hat{u}(x, y), \quad (\text{A2})$$

$$g_{\mu\nu} = \eta_{\mu\nu} + \kappa \hat{h}_{\mu\nu}(x, y), \quad \hat{u} = \frac{e^{2A(y)}}{2\sqrt{6}} \kappa \hat{r}(x, y). \quad (\text{A3})$$

The compact extra dimension is parametrized on a circle of radius r_c such that $y \in (-\pi r_c, +\pi r_c]$ and the 5D coordinates $X^M = (x^\mu, y)$ define a 4D Minkowski spacetime “slice” at each fixed value of y . We impose an orbifold invariance $y \leftrightarrow -y$ on the infinitesimal spacetime interval $G_{MN}dX^M dX^N$, and identify $y = 0$ and $y = \pi r_c$ as orbifold fixed points. For our spacetime signatures, we use the mostly minus convention, i.e., $\eta_{\mu\nu} \equiv \text{Diag}(+1, -1, -1, -1)$. Lowercase Greek letters (e.g., μ, ν, ρ, σ) denote 4D indices, uppercase Latin letters (e.g., M, N, R, S) denote 5D indices, and lowercase Latin letters (e.g., m, n, i, j) denote Kaluza-Klein mode numbers.

¹⁸If one is only concerned with external gravitons, and not doing scattering computations, one can impose the transverse-traceless conditions on all of the fields to simplify the computations of the interactions.

We then define the Lagrangian as

$$\mathcal{L}_{5D} \equiv \mathcal{L}_{\text{EH}} + \mathcal{L}_{\Phi\Phi} + \mathcal{L}_{\text{pot}} + \mathcal{L}_{\text{GHY}} + \Delta\mathcal{L}, \quad (\text{A4})$$

where the Einstein-Hilbert (EH), Gibbons-Hawking-York (GHY), scalar kinetic terms ($\Phi\Phi$), and scalar potential terms (pot) are defined as

$$\mathcal{L}_{\text{EH}} \equiv -\frac{2}{\kappa^2} \sqrt{GR}, \quad (\text{A5})$$

$$\begin{aligned} \mathcal{L}_{\text{GHY}} &\equiv -\frac{4}{\kappa^2} \partial_y [\sqrt{G}K] \\ &= \frac{2}{\kappa^2} \partial_y \left[\frac{w^2}{v} \sqrt{-\det g} \left(\tilde{g}^{\mu\nu} \partial_y (g_{\mu\nu}) + 4 \frac{\partial_y(w)}{w} \right) \right], \end{aligned} \quad (\text{A6})$$

$$\mathcal{L}_{\Phi\Phi} \equiv \sqrt{G} \left[\frac{1}{2} \tilde{G}^{MN} (\partial_M \hat{\Phi}) (\partial_N \hat{\Phi}) \right], \quad (\text{A7})$$

$$\mathcal{L}_{\text{pot}} \equiv -\frac{4}{\kappa^2} \left[\sqrt{G} V[\hat{\Phi}] + \sum_{i=1,2} \sqrt{G} V_i[\hat{\Phi}] \delta_i(\varphi) \right], \quad (\text{A8})$$

respectively. Here R is the Ricci scalar and K is the extrinsic curvature at the boundaries. G and \tilde{G} are the determinants of the metric and the induced metric, respectively. The Dirac deltas on the branes are defined in the limit as we approach the branes from within the $y \in [0, \pi r_c]$ half of the bulk: $\delta_1(\varphi) \equiv \delta(\varphi - 0^+)$ and $\delta_2(\varphi) \equiv \delta(\varphi - \pi^-)$, where $\varphi = y/r_c$.

Note that no dynamics are given to describe the physics of the branes themselves, which are assumed to arise through unspecified dynamics at some higher scale (e.g., string physics)—and, in particular, no modes arising from that physics (fluctuations of the branes themselves) are included. More on this below, when we discuss the effect of taking the so-called “stiff-wall” limit on the scalar mode expansions.

Meanwhile, the contribution $\Delta\mathcal{L}$ in \mathcal{L}_{5D} is a total derivative we add for convenience which generalizes a total derivative from our unstabilized analysis [11,12]. It cancels terms in the Lagrangian at linear order, eliminates mixing between tensor and scalar 5D fields at quadratic order, and simplifies the vertices relevant to this paper.¹⁹ Explicitly, we define it as

¹⁹If such a term were not introduced here, we would recover its effects (at least to quadratic order) as additional total derivative terms needed to make the Lagrangian explicitly canonical. This $\Delta\mathcal{L}$ naively generalized the $\Delta\mathcal{L}$ introduced in Refs. [11,12], and (unlike the latter) does not eliminate Dirac deltas or twice-differentiated quantities to all orders.

$$\Delta\mathcal{L} \equiv \frac{2}{\kappa^2} \partial_y \left[\frac{w^2}{v} \sqrt{-\det g} \left(-3 \frac{\partial_y(w)}{w} \right) \right]. \quad (\text{A9})$$

The perturbative expansion of the gravitational contributions as series in κ proceeds as usual. For convenience, we rewrite the bulk scalar field $\hat{\Phi}(x, y)$ such that it is perturbed about the background $\phi_0(\varphi)/\kappa$ by an amount \hat{f}/κ , i.e., $\hat{\Phi} \equiv \hat{\phi}/\kappa \equiv (\phi_0 + \hat{f})/\kappa$. This rewrite allows us to expand the bulk and brane potentials in \mathcal{L}_{pot} about $\hat{\phi} = \phi_0$ with respect to the dimensionless scalar fluctuation field \hat{f} like so:

$$V[\hat{\Phi}] = V + \dot{V} \hat{f} + \frac{1}{2} \ddot{V} (\hat{f})^2 + \mathcal{O}((\hat{f})^3), \quad (\text{A10})$$

$$V_i[\hat{\Phi}] = V_i + \dot{V}_i \hat{f} + \frac{1}{2} \ddot{V}_i (\hat{f})^2 + \mathcal{O}((\hat{f})^3), \quad (\text{A11})$$

where dots denote $\hat{\phi}$ functional derivatives, and V and V_i (and their $\hat{\phi}$ derivatives) are set to $\hat{\Phi} = \phi_0(y)/\kappa$ when their functional arguments are unspecified.

The path forward is, in principle, clear: find the appropriate solutions for the background fields $A(y)$ and $\phi_0(y)$, compute the Lagrangian that describes the dynamics of fluctuations about these background fields, and then diagonalize the quadratic terms in this Lagrangian to establish the (canonically normalized) physical modes and their interactions. In practice this is difficult because of the complicated algebraic structures involved in the mixing between the scalar components of the metric and Goldberger-Wise scalar field in the presence of a nontrivial scalar background. Some simplification results from the fact that diffeomorphism invariance implies that only one linear combination of the fields \hat{r} and \hat{f} are physical, and we can set the gauge of the calculation such that these fields are related according to

$$\phi'_0 \hat{f} = \sqrt{6} e^{2A} \hat{r}'. \quad (\text{A12})$$

We begin our analysis in general, without imposing this gauge condition, and only use the gauge constraint Eq. (A12) to identify the physical scalar modes after deriving the background equations, which we turn to now.

2. The Lagrangian to quadratic order

After weak field expanding the Lagrangian Eq. (A4) without applying the gauge condition Eq. (A12) or any background field equations, we obtain

$$\begin{aligned} \mathcal{L}_{5D} &\equiv \mathcal{L}_{5D, \text{bkgd}} + \mathcal{L}_{5D, h} + \mathcal{L}_{5D, r}^* + \mathcal{L}_{5D, f}^* + \mathcal{L}_{5D, hr}^* + \mathcal{L}_{5D, hf}^* \\ &\quad + \mathcal{L}_{5D, rf}^* + \mathcal{L}_{5D, hh} + \mathcal{L}_{5D, rr}^* + \mathcal{L}_{5D, ff}^* + \mathcal{O}(\kappa) \end{aligned} \quad (\text{A13})$$

to all orders in the background fields and up to quadratic order in the fluctuations. The background-only terms in the Lagrangian are

$$\mathcal{L}_{5D,\text{bkgd}} \equiv \frac{e^{-4A}}{2r_c^2\kappa^2} \left[24A'' - 48(A')^2 - (\phi'_0)^2 - 8 \left(Vr_c^2 + \sum_{i=1,2} V_i r_c \delta_i \right) \right]. \quad (\text{A14})$$

The terms linear in the fluctuations are

$$\mathcal{L}_{5D,h} \equiv \frac{e^{-4A}}{4r_c^2\kappa} \left[24A'' - 48(A')^2 - (\phi'_0)^2 - 8 \left(Vr_c^2 + \sum_{i=1,2} V_i r_c \delta_i \right) \right] \hat{h}, \quad (\text{A15})$$

$$\mathcal{L}_{5D,r}^* \equiv \frac{\sqrt{6}e^{-2A}}{r_c^2\kappa} [\hat{r}'' - 2A'\hat{r}'] + \frac{e^{-2A}}{2\sqrt{6}r_c^2\kappa} \left[-48A'' + 48(A')^2 + 3(\phi'_0)^2 + 8 \left(Vr_c^2 + 2 \sum_{i=1,2} V_i r_c \delta_i \right) \right] \hat{r}, \quad (\text{A16})$$

$$\mathcal{L}_{5D,f}^* \equiv -\frac{e^{-4A}}{r_c^2\kappa} \phi'_0 \hat{f}' - \frac{4e^{-4A}}{r_c^2\kappa} \left[\dot{V}r_c^2 + \sum_{i=1,2} \dot{V}_i r_c \delta_i \right] \hat{f}, \quad (\text{A17})$$

where $\hat{h} = \eta^{\mu\nu} \hat{h}_{\mu\nu}$. At quadratic order in the fluctuations, “off-diagonal” (mode-mixing) quadratic terms are

$$\mathcal{L}_{5D,hr}^* \equiv \frac{\kappa}{2} \mathcal{L}_{5D,r}^* \hat{h}, \quad (\text{A18})$$

$$\mathcal{L}_{5D,hf}^* \equiv \frac{\kappa}{2} \mathcal{L}_{5D,f}^* \hat{h}, \quad (\text{A19})$$

$$\mathcal{L}_{5D,rf}^* \equiv \sqrt{\frac{3}{2}} \frac{e^{-2A}}{r_c^2} \phi'_0 \hat{r} \hat{f}' + \sqrt{\frac{8}{3}} \frac{e^{-2A}}{r_c^2} \left[\dot{V}r_c^2 + 2 \sum_{i=1,2} \dot{V}_i r_c \delta_i \right] \hat{r} \hat{f}, \quad (\text{A20})$$

and the “on-diagonal” quadratic terms are given by

$$\begin{aligned} \mathcal{L}_{5D,hh} &\equiv e^{-2A} \left[(\partial^\nu \hat{h}_{\mu\nu})(\partial^\mu \hat{h}) - (\partial^\nu \hat{h}_{\mu\nu})^2 + \frac{1}{2} (\partial_\mu \hat{h}_{\nu\rho})^2 - \frac{1}{2} (\partial_\mu \hat{h})^2 \right] + \frac{e^{-4A}}{2r_c^2} [(\hat{h}')^2 - (\hat{h}'_{\mu\nu})^2] \\ &+ \frac{e^{-4A}}{16r_c^2} \left[24A'' - 48(A')^2 - (\phi'_0)^2 - 8 \left(Vr_c^2 + \sum_{i=1,2} V_i r_c \delta_i \right) \right] [(\hat{h})^2 - 2(\hat{h}_{\mu\nu})^2], \end{aligned} \quad (\text{A21})$$

$$\mathcal{L}_{5D,rr}^* \equiv \frac{1}{2} e^{+2A} (\partial_\mu \hat{r})^2 - \frac{1}{r_c^2} [2(\hat{r}')^2 + 3\hat{r}\hat{r}''] - \frac{1}{12r_c^2} \left[-48A'' + 5(\phi'_0)^2 + 16 \sum_{i=1,2} V_i r_c \delta_i \right] (\hat{r})^2, \quad (\text{A22})$$

$$\mathcal{L}_{5D,ff}^* \equiv \frac{1}{2} e^{-2A} (\partial_\mu \hat{f})^2 - \frac{e^{-4A}}{2r_c^2} (\hat{f}')^2 - \frac{2e^{-4A}}{r_c^2} \left[\dot{V}r_c^2 + \sum_{i=1,2} \dot{V}_i r_c \delta_i \right] (\hat{f})^2. \quad (\text{A23})$$

Here we use an asterisk to denote that we have not yet applied a gauge condition relating \hat{r} and \hat{f} .

The first line of terms in Eq. (A21) will yield the usual canonical spin-2 Lagrangians after Kaluza-Klein decomposition. As we will soon demonstrate, the other terms in Eq. (A21) will be canceled when the background fields satisfy their equations of motion. However, as close as $\mathcal{L}_{5D,hh}$ is to the desired spin-2 result, the quadratic analysis overall is complicated by the presence of the mixing terms $\mathcal{L}_{5D,hr}^*$ and $\mathcal{L}_{5D,hf}^*$, which seemingly imply kinetic mixing between the tensor field \hat{h} and the scalar fields \hat{f} and \hat{r} . To eliminate these mixing terms, we must derive the equations

of motion for the background fields and for the fluctuations, which we discuss next.

3. Equations of motion

The Einstein field equations derived from \mathcal{L}_{5D} equal

$$\begin{aligned} G_{MN} - V[\hat{\Phi}] G_{MN} - \left[V_1[\hat{\Phi}] \frac{\delta_1(\varphi)}{r_c} + V_2[\hat{\Phi}] \frac{\delta_2(\varphi)}{r_c} \right] \frac{\sqrt{\bar{G}}}{\sqrt{G}} \bar{G}_{MN} \\ = \frac{\kappa^2}{4} T_{MN}, \end{aligned} \quad (\text{A24})$$

where $\mathcal{G}_{MN} = R_{MN} - \frac{1}{2}G_{MN}R$ is the Einstein tensor, R_{MN} and $R = \tilde{G}^{AB}R_{AB}$ are the Ricci tensor and Ricci scalar, respectively, and the stress-energy tensor equals

$$T_{MN} = 2 \frac{\delta \mathcal{L}_{\Phi\Phi}}{\delta \tilde{G}^{MN}} - G_{MN} \mathcal{L}_{\Phi\Phi} = (\partial_M \hat{\Phi})(\partial_N \hat{\Phi}) - G_{MN} \left[\frac{1}{2} \tilde{G}^{AB} (\partial_A \hat{\Phi})(\partial_B \hat{\Phi}) \right]. \quad (\text{A25})$$

Recall that $\hat{\Phi} \equiv \hat{\phi}/\kappa \equiv (\phi_0 + \hat{f})/\kappa$. We will discuss the Einstein field equations in terms of their decomposition as $(M, N) = \{(\mu, \nu), (\mu, 5), (5, 5)\}$, to the first two orders in κ .

a. Background equations of motion

To lowest order in κ , in which no fluctuation fields are present, only the (μ, ν) and $(5, 5)$ Einstein field equations are nontrivial (because of the Lorentz invariance of the constant- y subspaces, the (μ, ν) components of the curvature are proportional to $\eta_{\mu\nu}$), and they imply

$$A'' = 2(A')^2 + \frac{1}{24}(\phi'_0)^2 + \frac{1}{3} \left[Vr_c^2 + \sum_{i=1,2} V_i r_c \delta_i \right],$$

$$Vr_c^2 = -6(A')^2 + \frac{1}{8}(\phi'_0)^2, \quad (\text{A26})$$

respectively, for the background fields. The first of these equations implies the boundary conditions [integrating over the end points using an S^1/Z_2 orbifold construction where $A(y)$ is assumed to be “even” under orbifold reflection]

$$V_1 r_c \delta_1 = +6A' \delta_1, \quad V_2 r_c \delta_2 = -6A' \delta_2. \quad (\text{A27})$$

By combining the equations of (A26), we may also write

$$A'' = \frac{1}{12} \left[(\phi'_0)^2 + 4 \sum_{i=1,2} V_i r_c \delta_i \right]. \quad ((10) \text{ revisited})$$

Note that Eq. (A26) ensures $\mathcal{L}_{5D, \text{bkgd}}$ and $\mathcal{L}_{5D, h}$ from Eqs. (A14) and (A15) vanish, and ensures $\mathcal{L}_{5D, hh}$ yields canonical spin-2 Lagrangians after Kaluza-Klein decomposition. Equation (A26) also simplifies the various pieces of the Lagrangian, including the linear \hat{r} terms:

$$\mathcal{L}_{5D, r}^* = \frac{\sqrt{6}}{r_c^2 \kappa} \partial_\varphi [e^{-2A} \hat{r}]. \quad (\text{A28})$$

While the mixing terms $\mathcal{L}_{5D, hr}^*$ and $\mathcal{L}_{5D, hf}^*$ remain at this point, these will vanish once we have analyzed the scalar sector, which we discuss now.

We obtain another background equation by considering the Euler-Lagrange equation of the scalar field. The terms independent of the fluctuations yield

$$\phi''_0 = 4A' \phi'_0 + 4\dot{V}r_c^2 + 4 \sum_{i=1,2} \dot{V}_i r_c \delta_i, \quad (\text{A29})$$

which implies its own boundary conditions (again, assuming the background scalar field configuration is even under the orbifold projection)

$$\dot{V}_1 r_c \delta_1 = +\frac{1}{2} \phi'_0 \delta_1, \quad \dot{V}_2 r_c \delta_2 = -\frac{1}{2} \phi'_0 \delta_2. \quad (\text{A30})$$

This simplifies $\mathcal{L}_{5D, f}^*$, such that

$$\mathcal{L}_{5D, f}^* = -\frac{1}{r_c^2 \kappa} \partial_\varphi [e^{-4A} \phi'_0 \hat{f}]. \quad (\text{A31})$$

This completes our derivation of background equations.

Recall that whenever we write a quantity multiplying $\delta_1(\varphi)$ or $\delta_2(\varphi)$, it is understood that the quantity is evaluated *in the limit* as φ approaches the appropriate orbifold fixed point from inside the $[0, \pi r_c]$ half of the bulk. This implies, for example, via Eq. (A29),

$$\phi''_0 \delta_i = (\phi''_0)_{\text{bulk}} \delta_i \equiv [4A' \phi'_0 + 4\dot{V}r_c^2] \delta_i. \quad (\text{A32})$$

This also ensures quantities such as $A'(\varphi)\delta_i(\varphi)$ in Eqs. (A27) and (A32) are written unambiguously, despite $A'(\varphi)$ being orbifold odd across the orbifold fixed points.

b. Lagrangian at quadratic order: Mode equations

Next, we examine the equations of motion derived from considering terms in the Lagrangian that are quadratic or lower in the fluctuations. These will give us the equations which will define the mode expansions—the Kaluza-Klein decomposition—for the fluctuating fields. As mentioned in the previous subsection, we will be ignoring the spin-2 fields—they will ultimately decouple from the scalar fields after having performed the correct scalar-field mode expansions.

We begin with the scalar fields in the metric. Simplifying the expressions using the background equations (A26)–(A29), the (μ, ν) , $(\mu, 5)$, and $(5, 5)$ Einstein field equations at $\mathcal{O}(\kappa)$ are satisfied only if, respectively,

$$0 = [\partial_\varphi - 4A'] [\sqrt{6} e^{2A} \hat{r}' - \phi'_0 \hat{f}], \quad (\text{A33})$$

$$0 = \partial_\mu [\sqrt{6} e^{2A} \hat{r}' - \phi'_0 \hat{f}], \quad (\text{A34})$$

$$\begin{aligned} \partial_\varphi \left[\frac{e^{-2A}}{\sqrt{6}} \phi'_0 \hat{f} \right] &= e^{2A} r_c^2 (\square \hat{r}) + 2A' \left\{ 2\hat{r}' + \left[\frac{e^{-2A}}{\sqrt{6}} \phi'_0 \hat{f} \right] \right\} \\ &+ \frac{8\dot{V}r_c^2}{\phi'_0} \left[\frac{e^{-2A}}{\sqrt{6}} \phi'_0 \hat{f} \right] + \frac{1}{6} (\phi'_0)^2 \hat{r}' \\ &+ 2(\delta_1 - \delta_2) \left[\frac{e^{-2A}}{\sqrt{6}} \phi'_0 \hat{f} \right], \end{aligned} \quad (\text{A35})$$

where the final equation has also utilized the jump conditions of Eq. (A30). As noted in Ref. [23], integrating Eq. (A35), we end up with a tautology and end up with boundary terms that provide no additional physical information. By moving the Dirac deltas of Eq. (A35) to the LHS and evaluating the derivatives, we derive an alternative form of the equation which lacks Dirac deltas (implicit or explicit):

$$\frac{e^{-2A}}{\sqrt{6}}\phi'_0\hat{f}' = e^{2A}r_c^2(\square\hat{r}) + 4A'\hat{r}' + \frac{1}{6}(\phi'_0)^2\hat{r} + \sqrt{\frac{8}{3}}e^{-2A}\dot{V}r_c^2\hat{f}. \quad (\text{A36})$$

We may also consider the Euler-Lagrangian equation of the fluctuation field \hat{f} at this order, which yields

$$\begin{aligned} \hat{f}'' &= e^{2A}r_c^2(\square\hat{f}) + 4A'\hat{f}' + 4\ddot{V}r_c^2\hat{f} \\ &+ \sqrt{\frac{3}{2}}e^{2A}\phi'_0\hat{r}' + \sqrt{\frac{2}{3}}e^{2A}[4\dot{V}r_c^2 + 3A'\phi'_0]\hat{r} \\ &+ \left[4\ddot{V}_1r_c\hat{f} + \sqrt{\frac{2}{3}}e^{2A}\phi'_0\hat{r}\right]\delta_1 \\ &+ \left[4\ddot{V}_2r_c\hat{f} - \sqrt{\frac{2}{3}}e^{2A}\phi'_0\hat{r}\right]\delta_2. \end{aligned} \quad (\text{A37})$$

This equation implies, via the orbifold construction, the boundary conditions

$$\begin{aligned} \hat{f}'\delta_1 &= \left[2\ddot{V}_1r_c\hat{f} + \frac{1}{\sqrt{6}}e^{2A}\phi'_0\hat{r}\right]\delta_1, \\ \hat{f}'\delta_2 &= -\left[2\ddot{V}_2r_c\hat{f} - \frac{1}{\sqrt{6}}e^{2A}\phi'_0\hat{r}\right]\delta_2. \end{aligned} \quad (\text{A38})$$

Multiply these jump conditions by $e^{-2A}\phi'_0/\sqrt{6}$ and use Eq. (A36) to get

$$\left\{e^{2A}r_c^2(\square\hat{r}) + 4A'\hat{r}' + \left[\frac{4\dot{V}r_c^2}{\phi'_0} - 2\ddot{V}_1r_c\right]\left[\frac{e^{-2A}}{\sqrt{6}}\phi'_0\hat{f}\right]\right\}\delta_1 = 0, \quad (\text{A39})$$

$$\left\{e^{2A}r_c^2(\square\hat{r}) + 4A'\hat{r}' + \left[\frac{4\dot{V}r_c^2}{\phi'_0} + 2\ddot{V}_2r_c\right]\left[\frac{e^{-2A}}{\sqrt{6}}\phi'_0\hat{f}\right]\right\}\delta_2 = 0. \quad (\text{A40})$$

Using Eq. (A32), we may instead write the jump conditions Eq. (A38) as

$$\begin{aligned} &\left\{e^{2A}r_c^2(\square\hat{r}) + 4A'\hat{r}' + \left[\frac{\phi''_0}{\phi'_0} - 4A' - 2\ddot{V}_1r_c\right]\left[\frac{e^{-2A}}{\sqrt{6}}\phi'_0\hat{f}\right]\right\}\delta_1 \\ &= 0, \\ &\left\{e^{2A}r_c^2(\square\hat{r}) + 4A'\hat{r}' + \left[\frac{\phi''_0}{\phi'_0} - 4A' + 2\ddot{V}_2r_c\right]\left[\frac{e^{-2A}}{\sqrt{6}}\phi'_0\hat{f}\right]\right\}\delta_2 \\ &= 0. \end{aligned} \quad (\text{A41})$$

This form is more common in the literature.

The linear field equations (A33)–(A35) and (A37) describe the scalar modes of the theory. Note, in particular, the recurring quantity $\sqrt{6}e^{2A}\hat{r}' - \phi'_0\hat{f}'$. This will vanish once we impose the gauge condition (A12), which is our next focus.

c. The gauge condition

The form of the metric specified by (A1)–(A3) does not completely fix the “gauge” for this calculation: we have access to various five-dimensional diffeomorphism transformations which maintain the form of the metric and with which we can choose to simplify our computations. In particular, as shown in [25], we can always perform a change of coordinate to impose the gauge condition introduced previously,

$$\sqrt{6}e^{2A}\hat{r}' = \phi'_0\hat{f}'. \quad (\text{A12})$$

One immediate consequence of this gauge choice is that the *sum* of the mixing terms $\mathcal{L}_{5D,hr}^*$ $\mathcal{L}_{5D,hf}^*$ vanishes, eliminating (as promised) any problematic mixing between the scalar and spin-2 mode sectors.

The physical implication of the gauge condition (A12) is that one combination of the scalar fields is a gauge artifact, and does not correspond to a propagating degree of freedom.²⁰ Note that the “mixing” of the scalar degree of freedom in the five-dimensional metric \hat{r} with the bulk scalar field \hat{f} only occurs in the presence of a y -dependent scalar background field configuration ($\phi'_0 \neq 0$). It is precisely this mixing between the two sectors that enables the dynamics which stabilize the size of the extra dimension in the Goldberger-Wise mechanism [17,18] and which simultaneously give rise to a “radion” mass. One advantage of working in this “unitary” gauge and eliminating the fluctuations of the scalar field \hat{f} in favor of scalar fluctuations of the metric \hat{r} is that all couplings linear in the physical scalar fields have the same algebraic form as couplings linear in the (massless) radion within the unstabilized model—simplifying the required coupling computations.

²⁰The precise combination of Lagrangian fields which is physical and the corresponding form of its interactions depend on the gauge choice—although all physical amplitudes are gauge-invariant.

Having imposed this gauge condition, the (μ, ν) and $(\mu, 5)$ linear Einstein field equations (A33) and (A34) vanish, the $(5, 5)$ linear Einstein field equation (A35) simplifies to

$$\begin{aligned} \hat{r}'' &= e^{2A} r_c^2 (\square \hat{r}) + \left[6A' + \frac{8\dot{V} r_c^2}{\phi_0'} \right] \hat{r}' \\ &+ \frac{1}{6} (\phi_0')^2 \hat{r} + 2(\delta_1 - \delta_2) \hat{r}', \end{aligned} \quad (\text{A42})$$

and the jump conditions in Eq. (A41) reduce to

$$\begin{aligned} &\left\{ e^{2A} r_c^2 (\square \hat{r}) - \left[2\dot{V}_1 r_c - \frac{\phi_0''}{\phi_0'} \right] \hat{r}' \right\} \delta_1 \\ &= \left\{ e^{2A} r_c^2 (\square \hat{r}) + \left[2\dot{V}_2 r_c + \frac{\phi_0''}{\phi_0'} \right] \hat{r}' \right\} \delta_2 = 0. \end{aligned} \quad (\text{A43})$$

These equations of motion for the field \hat{r} at quadratic order will define the Kaluza-Klein decomposition of the \hat{r} field. Note that, being careful about Dirac deltas,²¹

$$\begin{aligned} \partial_\varphi \left[\frac{e^{2A}}{(\phi_0')^2} \hat{r}' \right] &= \frac{2e^{2A}}{(\phi_0')^2} A' \hat{r}' - \frac{2e^{2A}}{(\phi_0')^3} (\phi_0'')_{\text{bulk}} \hat{r}' + \frac{e^{2A}}{(\phi_0')^2} \hat{r}'' \\ &= \frac{e^{2A}}{(\phi_0')^2} \left\{ - \left[6A' + \frac{8\dot{V} r_c^2}{\phi_0'} \right] \hat{r}' + \hat{r}'' \right\}, \end{aligned} \quad (\text{A44})$$

such that Eq. (A42) may also be written in a more conventional form:

$$\begin{aligned} \partial_\varphi \left[\frac{e^{2A}}{(\phi_0')^2} \hat{r}' \right] &- \frac{e^{2A}}{6} \hat{r} + 2[\delta_2(\varphi) - \delta_1(\varphi)] \frac{e^{2A}}{(\phi_0')^2} \hat{r}' \\ &= \frac{e^{4A}}{(\phi_0')^2} r_c^2 (\square \hat{r}). \end{aligned} \quad (\text{A45})$$

In the next subsection, we use Eqs. (A45) and (A43) to define the Kaluza-Klein decomposition of the 5D scalar field. [Note again that the jump conditions of the field \hat{r} are trivial in this form of the equation, and boundary conditions in Eq. (A43) are required.]

4. Kaluza-Klein decomposition of the scalar field

Next, we assume we can decompose the 5D scalar field $\hat{r}(x, y)$ into a tower of 4D fields $\hat{r}_i(x)$ and extradimensional wave functions $\gamma_i(\varphi)$:

$$\hat{r}(x, y) = \frac{1}{\sqrt{\pi r_c}} \sum_{i=0}^{+\infty} \hat{r}^{(i)}(x) \gamma_i(\varphi), \quad (\text{A46})$$

²¹Refer to the discussion after Eq. (A65) for more details. In short, the quantity $1/(\phi_0')^2$ cannot generate Dirac deltas upon differentiation.

where the states are arranged in order of increasing mass and $\varphi = y/r_c$. We will show that if the γ_i satisfies the Sturm-Liouville-like equation [compare to Eq. (A45)]:

$$\begin{aligned} \partial_\varphi \left[\frac{e^{2A}}{(\phi_0')^2} (\partial_\varphi \gamma_i) \right] &- \frac{e^{2A}}{6} \gamma_i + 2[\delta_2(\varphi) - \delta_1(\varphi)] \frac{e^{2A}}{(\phi_0')^2} (\partial_\varphi \gamma_i) \\ &= -\mu_{(i)}^2 \frac{e^{4A}}{(\phi_0')^2} \gamma_i, \end{aligned} \quad (\text{A47})$$

with boundary conditions [compare to Eq. (A43)]

$$\begin{aligned} (\partial_\varphi \gamma_i)|_{\varphi=0+} &= - \left[2\dot{V}_1 r_c - \frac{\phi_0''}{\phi_0'} \right]^{-1} \mu_{(i)}^2 e^{2A} \gamma_i|_{\varphi=0+}, \\ (\partial_\varphi \gamma_i)|_{\varphi=\pi-} &= + \left[2\dot{V}_2 r_c + \frac{\phi_0''}{\phi_0'} \right]^{-1} \mu_{(i)}^2 e^{2A} \gamma_i|_{\varphi=\pi-}, \end{aligned} \quad (\text{A48})$$

the $\hat{r}^{(i)}(x)$ are the properly normalized scalar Kaluza-Klein fields.

These scalar boundary conditions can alternatively be enforced (recalling that \hat{r} and hence γ_i are orbifold-even) using the equation introduced in the body of the paper:

$$\begin{aligned} \partial_\varphi \left[\frac{e^{2A}}{(\phi_0')^2} (\partial_\varphi \gamma_i) \right] &- \frac{e^{2A}}{6} \gamma_i = -\mu_{(i)}^2 \frac{e^{4A}}{(\phi_0')^2} \gamma_i \\ &\times \left\{ 1 + \frac{2\delta(\varphi)}{[2\dot{V}_1 r_c - \frac{\phi_0''}{\phi_0'}]} + \frac{2\delta(\varphi - \pi)}{[2\dot{V}_2 r_c + \frac{\phi_0''}{\phi_0'}]} \right\}. \end{aligned} \quad (12)$$

In this form, the Sturm-Liouville nature of the problem is manifest [24,25,33,34]. We will choose to normalize the wave functions such that

$$\begin{aligned} \delta_{m,n} &= \frac{6\mu_n^2}{\pi} \int_{-\pi}^{+\pi} d\varphi \gamma_m \gamma_n \frac{e^{4A}}{(\phi_0')^2} \\ &\times \left\{ 1 + \frac{2\delta(\varphi)}{[2\dot{V}_1 r_c - \frac{\phi_0''}{\phi_0'}]} + \frac{2\delta(\varphi - \pi)}{[2\dot{V}_2 r_c + \frac{\phi_0''}{\phi_0'}]} \right\} \end{aligned} \quad (\text{A49})$$

$$= \frac{6}{\pi} \int_{-\pi}^{+\pi} d\varphi \left[\frac{e^{2A}}{(\phi_0')^2} \gamma_m' \gamma_n' + \frac{e^{2A}}{6} \gamma_m \gamma_n \right], \quad (\text{A50})$$

where the second line follows by applying the differential Eq. (12) and integration by parts on the periodic doubled ‘‘orbifold.’’ We will show that this normalization will yield properly normalized scalar Kaluza-Klein modes.²² For our numerical investigations, we consider the ‘‘stiff-wall’’ limit $\dot{V}_{1,2} \rightarrow \infty$, in which case the eigenmodes γ_i satisfy Neumann boundary conditions. While the stiff-wall limit is (ultimately) unphysical, it is consistent with the simplification we made in ignoring the dynamics of the brane itself—and we can expect the results of our analysis correctly

²²Note that this choice is consistent since we have no massless physical scalar modes in this model.

describe low-energy properties of the system. Outside of numerical investigations, we do not take the stiff-wall limit.

To facilitate manipulations at the 5D level, define the following useful auxillary field:

$$\hat{z} \equiv \frac{1}{\sqrt{\pi r_c}} \sum_{i=0}^{+\infty} \mu_{(i)}^2 \hat{r}^{(i)}(x) \gamma_i(\varphi). \quad (\text{A51})$$

Using \hat{z} and the decomposition in (A46), the wave function differential equation becomes

$$\hat{r}'' = -e^{2A} \hat{z} + \left[6A' + \frac{8\dot{V} r_c^2}{\phi_0'} \right] \hat{r}' + \frac{1}{6} (\phi_0')^2 \hat{r} + 2(\delta_1 - \delta_2) \hat{r}', \quad (\text{A52})$$

such that

$$(\hat{r}'')_{\text{bulk}} \equiv -e^{2A} \hat{z} + \left[6A' + \frac{8\dot{V} r_c^2}{\phi_0'} \right] \hat{r}' + \frac{1}{6} (\phi_0')^2 \hat{r}, \quad (\text{A53})$$

and the wave function boundary conditions imply

$$\left\{ -e^{2A} \hat{z} + \left[4A' + \frac{4\dot{V} r_c^2}{\phi_0'} \right] \hat{r}' \right\} \delta_1 = \{ +2\ddot{V}_1 r_c \hat{r}' \} \delta_1, \\ \left\{ -e^{2A} \hat{z} + \left[4A' + \frac{4\dot{V} r_c^2}{\phi_0'} \right] \hat{r}' \right\} \delta_2 = \{ -2\dot{V}_2 r_c \hat{r}' \} \delta_2. \quad (\text{A54})$$

These boundary conditions are written in such a way to most easily replace away \dot{V}_i for future convenience. Let us now return to the Lagrangian.

5. The canonical scalar mode expansion

After applying the background equations of motion Eqs. (A26), (A27), (A29), and (A30), as well as the gauge condition (A12), we find the collections of Lagrangian terms Eqs. (A14)–(A23) are simplified. Most contributions now explicitly vanish:

$$\mathcal{L}_{5\text{D},\text{bkgd}} = \mathcal{L}_{5\text{D},h} = \mathcal{L}_{5\text{D},r}^* + \mathcal{L}_{5\text{D},f}^* = \mathcal{L}_{5\text{D},hr}^* + \mathcal{L}_{5\text{D},hf}^* = 0. \quad (\text{A55})$$

The tensor quadratic Lagrangian is now of the desired form to yield a tower of canonical spin-2 states after Kaluza-Klein decomposition:

$$\mathcal{L}_{5\text{D},hh} \equiv e^{-2A} \left[(\partial^\nu \hat{h}_{\mu\nu})(\partial^\mu \hat{h}) - (\partial^\nu \hat{h}_{\mu\nu})^2 + \frac{1}{2} (\partial_\mu \hat{h}_{\nu\rho})^2 - \frac{1}{2} (\partial_\mu \hat{h})^2 \right] + \frac{e^{-4A}}{2r_c^2} [(\hat{h}')^2 - (\hat{h}'_{\mu\nu})^2]. \quad (\text{A56})$$

The scalar quadratic Lagrangian, however, remains quite complicated. We organize the terms from each part of the quadratic scalar Lagrangian as follows:

$$\mathcal{L}_{5\text{D},rr} = \mathcal{L}_{5\text{D},rr}^* + \mathcal{L}_{5\text{D},rf}^* + \mathcal{L}_{5\text{D},ff}^* \quad (\text{A57}) \\ = \mathcal{L}_{\text{EH},rr} + \mathcal{L}_{\text{GHY},rr} + \mathcal{L}_{\Phi\Phi,rr} + \mathcal{L}_{\text{pot},rr} + \Delta\mathcal{L}_{rr}, \quad (\text{A58})$$

where

$$\mathcal{L}_{\text{EH},rr} = -\frac{1}{6} e^{2A} (\partial_\mu \hat{r})^2 - \frac{2}{3} e^{2A} \hat{r} (\square \hat{r}) - \frac{3}{r_c^2} (\hat{r}')^2 - \frac{4}{r_c^2} \hat{r} \hat{r}'' + \frac{8}{3r_c^2} A' \hat{r} \hat{r}' + \frac{16}{3r_c^2} A'' \hat{r}^2, \quad (\text{A59})$$

$$\mathcal{L}_{\text{GHY},rr} = \frac{4}{r_c^2} (\hat{r}')^2 + \frac{4}{r_c^2} \hat{r} \hat{r}'' - \frac{32}{3r_c^2} A' \hat{r} \hat{r}' - \frac{16}{3r_c^2} A'' \hat{r}^2, \quad (\text{A60})$$

$$\mathcal{L}_{\Phi\Phi,rr} = \frac{3}{(\phi_0')^2} e^{2A} (\partial_\mu \hat{r}')^2 - \frac{3}{r_c^2 (\phi_0')^4} \left[\phi_0' \hat{r}'' - \phi_0'' \hat{r}' + 2A' (\phi_0') \hat{r}' - \frac{1}{2} (\phi_0')^3 \hat{r} \right]^2 + \frac{(\phi_0')^2}{3r_c^2} \hat{r}^2, \quad (\text{A61})$$

$$\mathcal{L}_{\text{pot},rr} = \frac{4\dot{V}}{\phi_0'} \hat{r} \hat{r}' - \frac{12\ddot{V}}{(\phi_0')^2} (\hat{r}')^2 - \sum_{i=1,2} \left[-\frac{4V_i}{3r_c} \hat{r}^2 + \frac{8\dot{V}_i}{r_c \phi_0'} \hat{r} \hat{r}' - \frac{12\ddot{V}_i}{r_c (\phi_0')^2} (\hat{r}')^2 \right] \delta_i, \quad (\text{A62})$$

$$\Delta\mathcal{L}_{rr} = -\frac{3}{r_c^2} (\hat{r}')^2 - \frac{3}{r_c^2} \hat{r} \hat{r}'' + \frac{8}{r_c^2} A' \hat{r} \hat{r}' + \frac{4}{r_c^2} A'' \hat{r}^2. \quad (\text{A63})$$

For ease of comparison, we present these results without yet applying the background equations of motion or integration-by-parts. Note that the squared quantity in $\mathcal{L}_{\Phi\Phi,rr}$ is not singular because the delta functions in $\phi_0' \hat{r}'' - \phi_0'' \hat{r}'$ cancel.

Rather than consider this quadratic scalar Lagrangian directly, we first add a convenient total derivative (which we determined through trial and error). Generally, adding a total derivative to a Lagrangian reorganizes how

information is stored in the fields, but ultimately does not change the physics described by the Lagrangian; a classic example of this is the Gibbons-Hawking-York total derivative, which is used to make the Einstein-Hilbert Lagrangian of 4D gravity into a Lagrangian which only depends on fields and their first derivatives [30]. The total derivative we add to $\mathcal{L}_{5D,rr}$ [in addition to the total derivative $\Delta\mathcal{L}$ defined in Eq. (A9), which is already folded into $\mathcal{L}_{5D,rr}$] is

$$\bar{\Delta}\mathcal{L}_{rr} = \frac{1}{r_c^2} \partial_\varphi \left\{ \hat{r}\hat{r}' - \frac{3e^{2A}}{(\phi'_0)^2} \hat{z}\hat{r}' + \frac{12}{(\phi'_0)^4} [A'(\phi'_0)^2 + V'r_c^2](\hat{r}')^2 \right\} \quad (\text{A64})$$

such that we consider the combination

$$\mathcal{L}_{5D,rr} + \bar{\Delta}\mathcal{L}_{rr}. \quad (\text{A65})$$

The φ -derivative in $\bar{\Delta}\mathcal{L}_{rr}$ must be evaluated with care, lest we generate spurious singularities. In particular, Dirac delta-function singularities will be generated whenever φ -differentiating a discontinuity in a function's slope. In the present calculation, such a discontinuity only ever happens at the orbifold fixed points. The standard example

of this from Randall-Sundrum models is the twice-differentiated quantity $|\varphi|'' = \partial_\varphi(|\varphi|') = \partial_\varphi(\text{sign } \varphi)$, which equals $2(\delta_1 - \delta_2)$ in our scheme. If we are not careful when taking φ -derivatives more generally, we can accidentally generate spurious Dirac deltas which contradict our scheme. Consider φ -differentiating a quantity which is a square of a φ -differentiated quantity, such as $1/(\phi'_0)^2$. Naively, we attain $-2\phi''_0/(\phi'_0)^3$, which generates nonzero Dirac deltas through the ϕ''_0 . These Dirac deltas are spurious. First, note that ϕ_0 is a function of $|\varphi|$, which means ϕ'_0 is proportional to $|\varphi|' = \text{sign}(\varphi)$. Thus $1/(\phi'_0)^2 \propto 1/(\text{sign } \varphi)^2 = 1$ and $1/(\phi_0)^2$ lacks the overall factor of $|\varphi|'$ necessary to generate Dirac deltas upon φ -differentiation. While naive differentiation yields $-2\phi''_0/(\phi'_0)^3$, careful analysis reveals the φ -derivative of $1/(\phi'_0)^2$ is actually the Dirac delta-free quantity $-2(\phi''_0)_{\text{bulk}}/(\phi'_0)^3$.

For these reasons, evaluation of the φ -derivative present in the total derivative $\bar{\Delta}\mathcal{L}_{rr}$ yields fewer Dirac deltas than naively expected. Namely, they are only generated upon differentiating \hat{r}' in the first two terms and A' and V' in the third term. Explicitly, we thus calculate²³

$$\bar{\Delta}\mathcal{L}_{rr} = \frac{1}{r_c^2} \left\{ (\hat{r}')^2 + \hat{r}\hat{r}'' + \frac{6e^{2A}}{(\phi'_0)^3} (\phi''_0)_{\text{bulk}} \hat{z}\hat{r}' - \frac{6e^{2A}}{(\phi'_0)^2} A' \hat{z}' \hat{r}' - \frac{3e^{2A}}{(\phi'_0)^2} \hat{z}\hat{r}'' - 48 \frac{(\phi''_0)_{\text{bulk}}}{(\phi'_0)^5} [A'(\phi'_0)^2 + V'r_c^2](\hat{r}')^2 + \frac{24}{(\phi'_0)^4} [A'(\phi'_0)^2 + V'r_c^2] \hat{r}' (\hat{r}'')_{\text{bulk}} + \frac{12}{(\phi'_0)^4} [A''(\phi'_0)^2 + 2A'\phi'_0(\phi''_0)_{\text{bulk}} + V''r_c^2](\hat{r}')^2 \right\}. \quad (\text{A66})$$

Having evaluated $\bar{\Delta}\mathcal{L}_{rr}$ as above, we next consider the quadratic scalar terms $\mathcal{L}_{5D,rr} + \bar{\Delta}\mathcal{L}_{rr}$ after performing the following sequence of manipulations:

- (1) Use 4D integration-by-parts to eliminate any 4D d'Alembertian operators $\square = \partial_t^2 - \vec{\nabla}^2$, e.g., taking $\hat{r}(\square\hat{r})$ to $-(\partial_\mu\hat{r})^2$.
- (2) Eliminate all instances of \hat{r}'' , A'' , V , and ϕ''_0 (and their Dirac delta-free bulk forms) via Eqs. (A52), (A26), and (A29), respectively. Having done so, all Dirac deltas in the original weak field expanded Lagrangian have been made explicit.
- (3) Eliminate V_i and \dot{V}_i via the background equations (A27) and (A30), respectively.
- (4) Eliminate \dot{V}_i (which always multiplies an \hat{r}') via the boundary conditions, Eq. (A54).

- (5) Eliminate all instances of \ddot{V} , V' , and V'' , and in favor of \dot{V} and \dot{V}' via chain rule relations, i.e.,

$$V'' = \dot{V}'\phi'_0 + \dot{V}\phi''_0, \quad V' = \dot{V}\phi'_0, \quad \ddot{V} = \frac{\dot{V}'}{\phi'_0}, \quad (\text{A67})$$

where ϕ''_0 is then replaced by Eq. (A29), as done earlier. With this, all Dirac deltas in $\bar{\Delta}\mathcal{L}$ are also explicit.

After performing these replacements, we find all \dot{V} and \dot{V}' terms cancel, all Dirac deltas cancel, and we are left with very few terms²⁴

²³Technically these same considerations are important when calculating, for example, $\Delta\mathcal{L}$; however, $\Delta\mathcal{L}_{rr} = \partial_\varphi[(4A'\hat{r} - 3\hat{r}')\hat{r}]/r_c^2$, and naive differentiation yields the correct result.

²⁴An alternate way of deriving the canonical quadratic Lagrangian is to start with the expression on the right-hand side of Eq. (A68), and substituting \hat{z} from the (5, 5) Einstein equation (A36), as well as a similar expression for \hat{z}' derived from the Euler-Lagrange equation (A37). The resulting expression can be shown to be equal to the combination $\mathcal{L}_{5D,rr} + \bar{\Delta}\mathcal{L}_{rr}$. It is useful, when performing these manipulations, to remove explicit Dirac delta terms by using the background equations of motion given in Eqs. (A26) and (A29).

$$\begin{aligned} \mathcal{L}_{5D,rr} + \Delta\mathcal{L}_{rr} &= \frac{e^{2A}}{2} \left[(\partial_\mu \hat{r})^2 - \frac{\hat{z} \hat{r}}{r_c^2} \right] \\ &+ \frac{3e^{2A}}{(\phi'_0)^2} \left[(\partial_\mu \hat{r}')^2 - \frac{\hat{z}' \hat{r}'}{r_c^2} \right]. \end{aligned} \quad (\text{A68})$$

Upon Kaluza-Klein decomposition via Eqs. (A46) and (A51), $\mathcal{L}_{5D,rr} + \Delta\mathcal{L}_{rr}$ immediately yields

$$\begin{aligned} \sum_{m,n=0}^{+\infty} \frac{1}{2} [(\partial_\mu \hat{r}^{(m)})(\partial^\mu \hat{r}^{(n)}) - \mu_{(m)}^2 \hat{r}^{(m)} \hat{r}^{(n)}] \\ \cdot \frac{6}{\pi} \int_{-\pi}^{+\pi} d\varphi \left[\frac{e^{2A}}{(\phi'_0)^2} \gamma'_m \gamma'_n + \frac{e^{2A}}{6} \gamma_m \gamma_n \right]. \end{aligned} \quad (\text{A69})$$

Recall that the scalar state wave functions are normalized according to Eq. (A50), such that the integral on the right above (including the $6/\pi$) equals $\delta_{m,n}$. Consequently, we finally achieve our desired result:

$$\mathcal{L}_{5D,rr} + \Delta\mathcal{L}_{rr} = \sum_{n=0}^{+\infty} \left\{ \frac{1}{2} (\partial^\mu \hat{r}^{(n)})^2 - \frac{1}{2} \mu_{(n)}^2 (\hat{r}^{(n)})^2 \right\}. \quad (\text{A70})$$

That is, the 5D field $\hat{r}(x, y)$ generates a scalar tower of canonical 4D scalar states $\{\hat{r}^{(n)}(x)\}$, each having mass $m_{(n)} \equiv \mu_{(n)}/r_c$, where $\mu_{(n)}$ is determined by solving the differential equation problem for the wave functions $\{\gamma_n\}$ laid out between Eqs. (A47) and (A48).

APPENDIX B: THE INELASTIC SUM RULES RELATING COUPLINGS AND MASSES

This section derives and summarizes relationships between couplings and mass spectra that are relevant to ensuring at most $\mathcal{O}(s)$ growth of tree-level inelastic 2-to-2 helicity-zero massive spin-2 KK mode scattering amplitudes [i.e., the process $(k, l) \rightarrow (m, n)$] in the Goldberger-Wise-stabilized Randall-Sundrum I model. We briefly consider the implications of completeness before deriving a means of expressing all cubic and quartic (spin-2 exclusive) B-type couplings in terms of A-type couplings and special objects $B_{(kl)(mn)}$. These B-to-A formulas reduce the problem of finding amplitude-relevant formulas to the problem of simplifying sums of the form $\sum_j \mu_j^{2p} a_{klj} a_{mnj}$. The relevant (inelastic and elastic) sum rules are derived and then summarized in their own subsections. The final subsection describes the remaining set of (unproven) sum rules necessary for at-most $\mathcal{O}(s)$ growth in the fully inelastic process.

This appendix is written as a stand-alone report of the sum-rule relationships needed to ensure that all *inelastic* scattering amplitudes (all $2 \rightarrow 2$ scattering amplitudes with massive spin-2 fields of arbitrary mode number in the external states) grow no faster than $\mathcal{O}(s)$ and report which we have succeeded in proving—completing the program

begun in [9–12]. Section B 5 a derives relationships used in Sec. IV B of the main body of this paper and can be read independently.

1. Definitions

It is convenient to define generalized “couplings” to be overlap integrals of spin-2 and spin-0 wave functions of the form

$$x_{(k' \dots l') \dots (m' \dots n')}^{(p)} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \varepsilon^p (\partial_\varphi \gamma_k) \cdots \gamma_l \cdots (\partial_\varphi \psi_m) \cdots \psi_n, \quad (\text{B1})$$

where $A(\varphi)$ is the warp factor, $\varepsilon \equiv \exp(-A)$, and we add an additional factor of $(\partial_\varphi A)/kr_c$ to the integrand if only an odd number of differentiated wave functions are present in the integrand otherwise. The most common integrands appearing in the 4D effective Lagrangian carry powers ε^{-2} and ε^{-4} and are given special symbols:

$$a_{(k' \dots l') \dots (m' \dots n')} \equiv x_{(k' \dots l') \dots (m' \dots n')}^{(-2)}, \quad b_{(k' \dots l') \dots (m' \dots n')} = x_{(k' \dots l') \dots (m' \dots n')}^{(-4)}. \quad (\text{B2})$$

We will also encounter the label “c,” which is associated with $p = -6$. In particular, we encounter this integral often:

$$\begin{aligned} c_{k'l'm'n'} &\equiv x_{k'l'm'n'}^{(-6)} \\ &= \frac{1}{\pi} \int d\varphi \varepsilon^{-6} (\partial_\varphi \psi_k) (\partial_\varphi \psi_l) (\partial_\varphi \psi_m) (\partial_\varphi \psi_n). \end{aligned} \quad (\text{B3})$$

Another object that will be useful throughout the rest of this document is the symbol $\mathcal{D} \equiv \varepsilon^{-4} \partial_\varphi$, which is a combination of quantities that is often present as a result of the spin-2 Sturm-Liouville equation. When desperate for space, we will nest the notation even further, utilizing $\mathcal{D}_n \equiv \varepsilon^{-4} (\partial_\varphi \psi_n)$.

We will ultimately derive sum rules that allow us to rewrite certain useful sums of intermediate masses and couplings in terms of just the quartic A-type coupling a_{klmn} , three $B_{(kl)(mn)}$ objects (of which any two fix the value of the third), and integrals $c_{k'l'm'n'}$ and $x_{\phi'_0 \phi'_0 n' n' n' n'}^{(-8)}$.

2. Applications of completeness

The spin-2 mode completeness relation is

$$\delta(\varphi_2 - \varphi_1) = \frac{1}{\pi} \varepsilon(\varphi_1)^{-2} \sum_{j=0}^{+\infty} \psi_j(\varphi_1) \psi_j(\varphi_2), \quad (\text{B4})$$

where $\varepsilon(\varphi) \equiv e^{A(\varphi)}$ and the wave functions ψ_n satisfy $\partial_\varphi \mathcal{D} \psi_n = \partial_\varphi [\varepsilon^{-4} (\partial_\varphi \psi_n)] = -\mu_n^2 \varepsilon^{-2} \psi_n$. Spin-2 mode completeness allows us to collapse certain sums of cubic coupling products into a single quartic coupling, e.g.,

$$a_{klmn} = \sum_j a_{klj} a_{mnj} = \sum_j a_{kmj} a_{lnj} = \sum_j a_{knj} a_{lmj}, \quad (\text{B5})$$

$$b_{k'l'mn} = \sum_j b_{k'l'j} a_{mnj}. \quad (\text{B6})$$

Furthermore, by combining cubic B-type couplings in this same way, we arrive at

$$c_{k'l'm'n'} = \sum_j b_{k'l'j} b_{m'n'j} = \sum_j b_{k'm'j} b_{l'n'j} = \sum_j b_{k'n'j} b_{l'm'j}.$$

This is as far as direct applications of completeness can get us for now.

3. B-to-A formulas

This subsection details how to eliminate all B-type couplings (e.g., $b_{l'm'n}$ and $b_{k'l'mn}$) in favor of A-type couplings (e.g., a_{lmn} and a_{klmn}) and new structures $B_{(kl)(mn)}$. To begin, we note we can absorb a factor of μ^2 into A-type couplings with help from the Sturm-Liouville equation. A standard application of this technique proceeds as follows:

$$\mu_n^2 a_{lmn} = \frac{1}{\pi} \int d\varphi \varepsilon^{-2} \psi_l \psi_m [\mu_n^2 \psi_n] \quad (\text{B7})$$

$$= \frac{1}{\pi} \int d\varphi \varepsilon^{-2} \psi_l \psi_m [-\varepsilon^{+2} \partial_\varphi (\mathcal{D}\psi_n)] \quad (\text{B8})$$

$$= \frac{1}{\pi} \int d\varphi \partial_\varphi [\psi_l \psi_m] (\mathcal{D}\psi_n) \quad (\text{B9})$$

$$= \frac{1}{\pi} \int d\varphi \varepsilon^{-4} (\partial_\varphi \psi_l) \psi_m (\partial_\varphi \psi_n) \\ + \frac{1}{\pi} \int d\varphi \varepsilon^{-4} \psi_l (\partial_\varphi \psi_m) (\partial_\varphi \psi_n) \quad (\text{B10})$$

$$= b_{l'mn'} + b_{lm'n'}, \quad (\text{B11})$$

where integration by parts was utilized between Eqs. (B8) and (B9); because $(\mathcal{D}\psi_n)$ vanishes on the boundaries, there is no surface term. This and the equivalent calculation with the quartic A-type coupling yield

$$\mu_n^2 a_{lmn} = b_{l'mn'} + b_{lm'n'}, \quad (\text{B12})$$

$$\mu_n^2 a_{klmn} = b_{k'l'mn'} + b_{k'l'mn'} + b_{klm'n'}. \quad (\text{B13})$$

By considering different permutations of KK indices, each of these equations corresponds to three and four unique constraints, respectively. Because there are only three unique cubic B-type couplings with KK indices l , m , and n (specifically, $b_{l'm'n}$, $b_{l'mn'}$, and $b_{lm'n'}$), Eq. (B12) can be inverted to yield

$$\boxed{b_{l'm'n} = \frac{1}{2} [\mu_l^2 + \mu_m^2 - \mu_n^2] a_{lmn}} \quad (\text{B14})$$

with which we can eliminate all cubic B-type couplings in favor of the cubic A-type coupling.

There are *six* unique quartic B-type couplings with KK indices k , l , m , and n . We first halve this set by rewriting each quartic B-type coupling $b_{k'l'mn}$ in terms of new objects $B_{(kl)(mn)}$. These new objects are motivated as follows: note that Eq. (B13) implies

$$\frac{1}{2} [\mu_k^2 + \mu_l^2 - \mu_m^2 - \mu_n^2] a_{klmn} = b_{k'l'mn} - b_{klm'n'}. \quad (\text{B15})$$

Equivalently, we may write this as

$$b_{k'l'mn} + \frac{1}{2} [\mu_m^2 + \mu_n^2] a_{klmn} = b_{klm'n'} + \frac{1}{2} [\mu_k^2 + \mu_l^2] a_{klmn}. \quad (\text{B16})$$

In other words, the quantity on the LHS possesses a symmetry under the pair swap $(k, l) \leftrightarrow (m, n)$. Furthermore, this symmetry is maintained under the addition of any quantity $\tilde{B}_{(kl)(mn)}$ which is also symmetric under this pair swap. Inspired by Eq. (B16), we define

$$B_{(kl)(mn)} \equiv b_{k'l'mn} + \frac{1}{2} [\mu_m^2 + \mu_n^2] a_{klmn} + \tilde{B}_{(kl)(mn)}. \quad (\text{B17})$$

We will choose the quantity $\tilde{B}_{(kl)(mn)}$ momentarily. Because the B-type couplings satisfy Eq. (B13), the sum of all unique B objects satisfies

$$B_{(kl)(mn)} + B_{(km)(ln)} + B_{(kn)(lm)} \\ = \tilde{\mu}^2 a_{klmn} + \tilde{B}_{(kl)(mn)} + \tilde{B}_{(km)(ln)} + \tilde{B}_{(kn)(lm)}, \quad (\text{B18})$$

where $\tilde{\mu}^2 \equiv \mu_k^2 + \mu_l^2 + \mu_m^2 + \mu_n^2$. That is, we can ensure the convenient property

$$B_{(kl)(mn)} + B_{(km)(ln)} + B_{(kn)(lm)} \doteq 0 \quad (\text{B19})$$

as long as we choose $\tilde{B}_{(kl)(mn)}$ such that

$$\tilde{B}_{(kl)(mn)} + \tilde{B}_{(km)(ln)} + \tilde{B}_{(kn)(lm)} = -\tilde{\mu}^2 a_{klmn}. \quad (\text{B20})$$

One immediate choice (and the choice we take now) is to set each \tilde{B} equal to one-third of $-\tilde{\mu}^2 a_{klmn}$,

$$\tilde{B}_{(kl)(mn)} \doteq -\frac{1}{3} a_{klmn}. \quad (\text{B21})$$

This yields (as a replacement rule for $b_{k'l'mn}$ and definition of $B_{(kl)(mn)}$)

$$\boxed{b_{k'l'mn} = B_{(kl)(mn)} + \frac{1}{6} [2(\mu_k^2 + \mu_l^2) - (\mu_m^2 + \mu_n^2)] a_{klmn}}, \quad (\text{B22})$$

where B is symmetric within each pair and between pairs

$$B_{(kl)(mn)} = B_{(mn)(kl)} = B_{(mn)(lk)} \quad (\text{B23})$$

and satisfies the additional constraint

$$B_{(kl)(mn)} + B_{(km)(ln)} + B_{(kn)(lm)} = 0 \quad (\text{B24})$$

such that only two among $\{B_{(kl)(mn)}, B_{(km)(ln)}, B_{(kn)(lm)}\}$ are linearly independent. Note that $B_{(kl)(mn)}$ has the same symmetry properties as $\sum_j \mu_j^{2p} a_{klj} a_{mnj}$. Because Eq. (B22) reduces B-type couplings to A-type couplings as much as possible, we refer to it as the quartic B-to-A rule. This and Eq. (B14) comprise the desired B-to-A formulas.

The above rules are sufficient as is for reducing the sum $\sum_j \mu_j^2 a_{klj} a_{mnj}$ and yielding the first nontrivial sum rule. Using the cubic coupling Eq. (B12) with completeness yields

$$b_{k'l'mn} = \frac{1}{2} [\mu_k^2 + \mu_l^2] a_{klmn} - \frac{1}{2} \sum_{j=0} \mu_j^2 a_{klj} a_{mnj}. \quad (\text{B25})$$

Meanwhile, the LHS can be simplified via Eq. (B22). Solving for the undetermined sum then gives us

$$\sum_j \mu_j^4 a_{klj} a_{mnj} = 4c_{k'l'm'n'} - 2(\vec{\mu}^2) B_{(kl)(mn)} + \left[\frac{1}{3} (\vec{\mu}^2)^2 - (\mu_k^2 + \mu_l^2)(\mu_m^2 + \mu_n^2) \right] a_{klmn} \quad (\text{B30})$$

as desired. Deriving the $\sum_j \mu_j^6 a_{klj} a_{mnj}$ relation requires significantly more work.

5. The μ_j^6 sum rule

a. Elastic

As a warm-up to the inelastic case, let us first derive the μ_j^6 sum rule (and review the other sum rules) as they appear

$$b_{n'l'n} = \frac{1}{2} [2\mu_n^2 - \mu_j^2] a_{nnj}, \quad b_{j'n'n} = \frac{1}{2} \mu_j^2 a_{nnj}, \quad b_{n'n'nn} = \frac{1}{3} \mu_n^2 a_{nnnn}, \quad (\text{B31})$$

which allow us to rewrite any spin-2 exclusive B-type couplings in terms of A-type couplings. Using these in combination with completeness, we find

$$\begin{aligned} a_{nnnn} &= \sum_j a_{nnj}^2, \\ b_{n'n'nn} &= \sum_j b_{n'n'j} a_{nnj} = \mu_n^2 \sum_j a_{nnj}^2 - \frac{1}{2} \sum_j \mu_j^2 a_{nnj}^2, \\ c_{n'n'n'n'} &\equiv \sum_j b_{n'n'j}^2 = \mu_n^4 \sum_j a_{nnj}^2 - \mu_n^2 \sum_j \mu_j^2 a_{nnj}^2 + \frac{1}{4} \sum_j \mu_j^4 a_{nnj}^2, \end{aligned}$$

$$\sum_{j=0} \mu_j^2 a_{klj} a_{mnj} = -2B_{(kl)(mn)} + \frac{1}{3} \vec{\mu}^2 a_{klmn}, \quad (\text{B26})$$

where $\vec{\mu}^2 \equiv \mu_k^2 + \mu_l^2 + \mu_m^2 + \mu_n^2$. We next turn our attention to $\sum_j \mu_j^4 a_{klj} a_{mnj}$ and then $\sum_j \mu_j^6 a_{klj} a_{mnj}$.

4. The μ_j^4 sum rule

The $\sum_j \mu_j^4 a_{klj} a_{mnj}$ relation is relatively straightforward. As defined in Eq. (B3), we can rewrite $c_{k'l'm'n'}$ in terms of B-type cubic couplings, to which we can then apply the B-to-A formulas:

$$c_{k'l'm'n'} = \sum_{j=0} b_{k'l'j} b_{m'n'j} \quad (\text{B27})$$

$$= \frac{1}{4} \sum_j [\mu_k^2 + \mu_l^2 - \mu_j^2] [\mu_m^2 + \mu_n^2 - \mu_j^2] a_{klj} a_{mnj} \quad (\text{B28})$$

$$\begin{aligned} &= \frac{1}{4} (\mu_k^2 + \mu_l^2) (\mu_m^2 + \mu_n^2) a_{klmn} \\ &\quad - \frac{1}{4} (\vec{\mu}^2) \sum_j \mu_j^2 a_{klj} a_{mnj} + \frac{1}{4} \sum_j \mu_j^4 a_{klj} a_{mnj} \quad (\text{B29}) \end{aligned}$$

such that, using Eq. (B26) and solving for the undetermined sum $\sum_j \mu_j^4 a_{klj} a_{mnj}$,

in the elastic case, i.e., when $k = l = m = n$. This will provide the general flow of the argument which is made more complicated in the inelastic case. Definitions for x , a , b , etc., are included in Sec. B 1.

Using the spin-2 completeness relation and differential equation alone, we have previously derived many elastic coupling relations [9–12]. For example, there are the elastic B-to-A formulas

which imply various expressions for sums of the form $\sum_j \mu_j^{2p} a_{nnj}^2$:

$$\sum_{j=0}^{+\infty} a_{nnj}^2 = a_{nnnn}, \quad (\text{B32})$$

$$\sum_{j=0}^{+\infty} \mu_j^2 a_{nnj}^2 = \frac{4}{3} \mu_n^2 a_{nnnn}, \quad (\text{B33})$$

$$\sum_{j=0}^{+\infty} \mu_j^4 a_{nnj}^2 = 4c_{n'n'n'n'} + \frac{4}{3} \mu_n^2 a_{nnnn}. \quad (\text{B34})$$

These relations allow us to quickly rewrite various sums between B-type couplings, including

$$\sum_{j=0}^{+\infty} b_{n'n'j} b_{j'n'n} = \frac{1}{3} \mu_n^4 a_{nnnn} - c_{n'n'n'n'}. \quad (\text{B35})$$

As discussed in the main text, a combination of the GW model sum rules ensuring cancellation of the $\mathcal{O}(s^3)$ and $\mathcal{O}(s^2)$ contributions to the amplitude may be written as

$$\begin{aligned} \text{Stabilized RS1: } \sum_{j=0}^{+\infty} [5\mu_n^2 - \mu_j^2] \mu_j^4 a_{nnj}^2 &= \frac{16}{3} \mu_n^6 a_{nnnn} \\ &+ 9 \sum_{i=0}^{+\infty} \mu_{(i)}^2 a_{n'n'(i)}^2. \quad (23) \end{aligned}$$

We can use our existing relations to rewrite this expression as

$$\sum_{j=0}^{+\infty} \mu_j^6 a_{nnj}^2 = 20\mu_n^2 c_{n'n'n'n'} + \frac{4}{3} \mu_n^6 a_{nnnn} + 9 \sum_{i=0}^{+\infty} \mu_{(i)}^2 a_{n'n'(i)}^2. \quad (\text{B36})$$

It is this variant we now seek to prove.

To begin, note that the B-to-A formulas relate the sum $\sum_j \mu_j^6 a_{nnj}^2$ to the sum $\sum_j \mu_j^2 b_{n'n'j}^2$ like so:

$$\sum_{j=0}^{+\infty} \mu_j^6 a_{nnj}^2 = 16\mu_n^2 c_{n'n'n'n'} + 4 \sum_{j=0}^{+\infty} \mu_j^2 b_{n'n'j}^2. \quad (\text{B37})$$

Thus, if we determine a means of rewriting $\sum_j \mu_j^2 b_{n'n'j}^2$ in terms of $c_{n'n'n'n'}$ and a_{nnnn} , then we will have a means of doing the same for the desired sum.

We will arrive at the desired form by considering two integrals of total derivatives, each of which vanishes because $(\partial_\phi \psi_n)$ vanishes at the orbifold fixed points $\phi \in \{0, \pi\}$.

Integral 1: To begin, consider the following trivial integral:

$$\frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi \partial_\phi \{ [(\partial_\phi \psi_j) - 6(\partial_\phi A) \psi_j] \varepsilon^{-6} (\partial_\phi \psi_n)^2 \} = 0. \quad (\text{B38})$$

By evaluating the net derivative and using the spin-2 mode differential equation to simplify second derivatives of spin-2 wave functions,²⁵ we attain

$$\begin{aligned} 0 &= -12 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi (\partial_\phi A)^2 \varepsilon^{-6} (\partial_\phi \psi_n)^2 \psi_j \right\} \\ &+ 12(kr_c) \mu_n^2 x_{n'n'j}^{(-4)} - 2\mu_n^2 b_{j'n'n} \\ &- 6 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi (\partial_\phi^2 A) \varepsilon^{-6} (\partial_\phi \psi_n)^2 \psi_j \right\} - \mu_j^2 b_{n'n'j}. \quad (\text{B39}) \end{aligned}$$

We can then construct an instance of $\sum_j \mu_j^2 b_{n'n'j}^2$ within this by multiplying it by $b_{n'n'j}$ and summing over j . This yields

$$\begin{aligned} 0 &= -12 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi (\partial_\phi A)^2 \varepsilon^{-8} (\partial_\phi \psi_n)^4 \right\} \\ &+ 12(kr_c) \mu_n^2 x_{n'n'n'n}^{(-6)} - 2\mu_n^2 \sum_{j=0}^{+\infty} b_{n'n'j} b_{j'n'n} \\ &- 6 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi (\partial_\phi^2 A) \varepsilon^{-8} (\partial_\phi \psi_n)^4 \right\} - \sum_{j=0}^{+\infty} \mu_j^2 b_{n'n'j}^2. \quad (\text{B40}) \end{aligned}$$

Integral 2: Next consider the following trivial integral:

$$\frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi \partial_\phi \left\{ \left[\frac{3}{2} (\partial_\phi A) \varepsilon^{-2} (\partial_\phi \psi_n) - \mu_n^2 \psi_n \right] \varepsilon^{-6} (\partial_\phi \psi_n)^3 \right\} = 0. \quad (\text{B41})$$

Evaluating this derivative in the same way as we did with the first integral, we find

$$\begin{aligned} 0 &= 12 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi (\partial_\phi A)^2 \varepsilon^{-8} (\partial_\phi \psi_n)^4 \right\} \\ &- 12(kr_c) \mu_n^2 x_{n'n'n'n}^{(-6)} - \mu_n^2 c_{n'n'n'n'} + 3\mu_n^4 b_{n'n'nn} \\ &+ \frac{3}{2} \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\phi (\partial_\phi^2 A) \varepsilon^{-8} (\partial_\phi \psi_n)^4 \right\}. \quad (\text{B42}) \end{aligned}$$

Combining: Summing Eqs. (B40) and (B42) and then solving for $\sum_{j=0}^{+\infty} \mu_j^2 b_{n'n'j}^2$ immediately yields

²⁵It is useful to repackage each $(\partial_\phi \psi)$ instead as $\varepsilon^{+4}(\mathcal{D}\psi)$ where $\mathcal{D} \equiv \varepsilon^{-4} \partial_\phi$ because then the spin-2 mode equation may be utilized more directly in the form $\partial_\phi \mathcal{D}\psi_n = -\mu_n^2 \varepsilon^{-2} \psi_n$.

$$\sum_{j=0}^{+\infty} \mu_j^2 b_{n'n'j}^2 = -\mu_n^2 c_{n'n'n'n'} + 3\mu_n^4 b_{n'n'nn} - 2\mu_n^2 \sum_{j=0}^{+\infty} b_{n'n'j} b_{j'n'n} - \frac{9}{2} \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi (\partial_\varphi^2 A) \varepsilon^{-8} (\partial_\varphi \Psi_n)^4 \right\}. \quad (\text{B43})$$

We already know how to rewrite $b_{n'n'nn}$ and $\sum_j b_{n'n'j} b_{j'n'n}$ in terms of $c_{n'n'n'n'}$ and a_{nnnn} ; namely, Eqs. (B31) and (B35). Using these, Eq. (B43) becomes

$$\sum_{j=0}^{+\infty} \mu_j^2 b_{n'n'j}^2 = \mu_n^2 c_{n'n'n'n'} + \frac{1}{3} \mu_n^6 a_{nnnn} - \frac{9}{2} \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi (\partial_\varphi^2 A) \varepsilon^{-8} (\partial_\varphi \Psi_n)^4 \right\}. \quad (\text{B44})$$

Finally, applying this result to Eq. (B37) gives us an expression for the desired “ μ_j^6 ” sum:

$$\sum_{j=0}^{+\infty} \mu_j^6 a_{nnj}^2 = 20\mu_n^2 c_{n'n'n'n'} + \frac{4}{3} \mu_n^6 a_{nnnn} - 18 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi (\partial_\varphi^2 A) \varepsilon^{-8} (\partial_\varphi \Psi_n)^4 \right\}. \quad (\text{B45})$$

Using the “ μ_j^4 ” sum in Eq. (B34) we obtain the result quoted in the text:

$$\sum_{j=0}^{+\infty} [5\mu_n^2 - \mu_j^2] \mu_j^4 a_{nnj}^2 = \frac{16}{3} \mu_n^6 a_{nnnn} + 18 \left\{ \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi (\partial_\varphi^2 A) \varepsilon^{-8} (\partial_\varphi \Psi_n)^4 \right\}. \quad (25)$$

This is all that is required for the elastic case discussed in the main text. The inelastic case covered in the next subsection is logically similar, but involves longer expressions and significantly more algebra.

b. Inelastic

Before beginning the derivation of the inelastic μ_j^6 sum rule, it is advantageous to define a well-organized polynomial basis with which we can write our results succinctly. In particular, because the sums we wish to simplify ($\sum_j \mu_j^{2p} a_{klj} a_{mnj}$) and relevant quartic degrees of freedom

all have (at least) the symmetries of $B_{(kl)(mn)}$, it is useful to define a symmetrization operation that forms quantities with symmetries identical to $B_{(kl)(mn)}$:

$$\langle f_{klmn} \rangle \equiv \{ [f_{klmn} + (k \leftrightarrow l)] + (m \leftrightarrow n) \} + (kl \leftrightarrow mn) \quad (\text{B46})$$

$$= f_{klmn} + f_{lkmn} + f_{klnm} + f_{lknm} + f_{mnlk} + f_{nmkl} + f_{nmkl} + f_{nmlk}. \quad (\text{B47})$$

This allows us to quickly construct a finite basis for polynomials of $\mu^2 \in \{\mu_k^2, \mu_l^2, \mu_m^2, \mu_n^2\}$ having the aforementioned symmetry structures. For a single power of μ^2 , there is only one basis element:

$$\alpha_{(kl)(mn)}^{(1,1)} \equiv \langle \mu_k^2 \rangle = 2\bar{\mu}^2 \equiv 2(\mu_k^2 + \mu_l^2 + \mu_m^2 + \mu_n^2). \quad (\text{B48})$$

For two powers of μ^2 , there are three:

$$\alpha_{(kl)(mn)}^{(2,1)} \equiv \langle \mu_k^4 \rangle, \quad \alpha_{(kl)(mn)}^{(2,2)} \equiv \langle \mu_l^2 \mu_k^2 \rangle, \quad \alpha_{(kl)(mn)}^{(2,3)} \equiv \langle \mu_m^2 \mu_k^2 \rangle, \quad (\text{B49})$$

and for three powers of μ^2 , there are four:

$$\alpha_{(kl)(mn)}^{(3,1)} \equiv \langle \mu_k^6 \rangle, \quad \alpha_{(kl)(mn)}^{(3,2)} \equiv \langle \mu_l^2 \mu_k^4 \rangle, \quad \alpha_{(kl)(mn)}^{(3,3)} \equiv \langle \mu_m^2 \mu_k^4 \rangle, \quad \alpha_{(kl)(mn)}^{(3,4)} \equiv \langle \mu_m^2 \mu_l^2 \mu_k^2 \rangle. \quad (\text{B50})$$

With these, we can generically construct any polynomial of the squared masses (up to cubic degree) having the aforementioned symmetry properties:

$$M_{(kl)(mn)}^{(1)}(c_1) = 2c_1 \bar{\mu}^2, \quad (\text{B51})$$

$$M_{(kl)(mn)}^{(2)}(c_1, c_2, c_3) = \sum_{i=1}^3 c_i \alpha_{(kl)(mn)}^{(2,i)}, \quad (\text{B52})$$

$$M_{(kl)(mn)}^{(3)}(c_1, c_2, c_3, c_4) = \sum_{i=1}^4 c_i \alpha_{(kl)(mn)}^{(3,i)}. \quad (\text{B53})$$

Note that these symbols are intentionally linear in their c_i arguments. In this language, Eq. (B30) may be rewritten as

$$\sum_j \mu_j^4 a_{klj} a_{mnj} = 4c_{k'l'm'n'} - 2\bar{\mu}^2 B_{(kl)(mn)} + \frac{1}{6} M_{(kl)(mn)}^{(2)}(1, 1, -1) a_{klmn}. \quad (\text{B54})$$

We now proceed to the $\sum_j \mu_j^6 a_{klj} a_{mnj}$ rule.

As in the previous subsection, we begin our derivation by applying the B-to-A formulas to a sum of cubic B-type couplings, and then apply existing sum rules:

$$\begin{aligned} \sum_j \mu_j^2 b_{k'l'j} b_{m'n'j} &= \frac{1}{4} \sum_j \mu_j^6 a_{klj} a_{mnj} - \vec{\mu}^2 c_{k'l'm'n'} - \frac{1}{12} \vec{\mu}^2 (\mu_k^2 + \mu_l^2 - \mu_m^2 - \mu_n^2)^2 a_{klmn} \\ &\quad + \frac{1}{2} [(\vec{\mu}^2)^2 - (\mu_k^2 + \mu_l^2)(\mu_m^2 + \mu_n^2)] B_{(kl)(mn)}. \end{aligned} \quad (\text{B55})$$

On the RHS, only the desired sum $\sum_j \mu_j^6 a_{klj} a_{mnj}$ remains undetermined. However, unlike the previous subsection, we do not yet have a simplification of the LHS of this expression. To find such a simplification, we concoct a vanishing combination of two integrals [namely, (I1)_{(kl)j} and (I2)_{k(lmn)} of Eqs. (B56) and (B62)], each of which vanishes independently because their integrands are total derivatives.

Integral 1: The first integral yields a vanishing combination of cubic quantities and is defined as

$$(\text{I1})_{(kl)j} \equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \partial_\varphi \left\{ \left[\frac{1}{2} (\partial_\varphi \psi_j) - 3(\partial_\varphi A) \psi_j \right] \varepsilon^{-6} (\partial_\varphi \psi_k) (\partial_\varphi \psi_l) \right\} \quad (\text{B56})$$

$$= \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \partial_\varphi \left\{ \frac{1}{2} \varepsilon^{+6} \mathcal{D}_j \mathcal{D}_k \mathcal{D}_l - 3A' \varepsilon^{+2} \psi_j \mathcal{D}_k \mathcal{D}_l \right\}, \quad (\text{B57})$$

where $\mathcal{D}_x \equiv \mathcal{D}\psi_x \equiv \varepsilon^{-4} (\partial_\varphi \psi_x)$. By explicitly applying the differentiation and using the wave function of the spin-2 modes ($\partial_\varphi \mathcal{D}_x = -\mu_x^2 \varepsilon^{-2} \psi_x$), we attain

$$\begin{aligned} (\text{I1})_{(kl)j} &\equiv \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \left\{ [-3A'' - 6(A')^2] \varepsilon^{+2} \mathcal{D}_k \mathcal{D}_l \psi_j - \frac{1}{2} \varepsilon^{+4} [\mu_k^2 \psi_k \mathcal{D}_l + \mu_l^2 \mathcal{D}_k \psi_l] \mathcal{D}_j \right. \\ &\quad \left. + 3A' [\mu_k^2 \psi_k \mathcal{D}_l + \mu_l^2 \mathcal{D}_k \psi_l] \psi_j - \frac{1}{2} \varepsilon^{+4} \mathcal{D}_k \mathcal{D}_l \mu_j^2 \psi_j \right\}. \end{aligned} \quad (\text{B58})$$

Next, to attain the desired index structure, we multiply by $b_{m'n'j}$ and sum over all j :

$$\begin{aligned} \sum_{j=0}^{+\infty} (\text{I1})_{(kl)j} b_{m'n'j} &= \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \{ [-3A'' - 6(A')^2] \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \} - \frac{1}{2} \sum_{j=0}^{+\infty} \mu_j^2 b_{k'l'j} b_{m'n'j} \\ &\quad - \frac{1}{2} \sum_{j=0}^{+\infty} [\mu_k^2 b_{k'l'j} b_{m'n'j} + \mu_l^2 b_{k'l'j} b_{m'n'j}] + \frac{3}{\pi} \int_{-\pi}^{+\pi} d\varphi \{ A' \varepsilon^{+6} [\mu_k^2 \psi_k \mathcal{D}_l + \mu_l^2 \mathcal{D}_k \psi_l] \mathcal{D}_m \mathcal{D}_n \}. \end{aligned} \quad (\text{B59})$$

Because (I1)_{(kl)j} vanishes, this sum vanishes too, as does the following combination:

$$\begin{aligned} (\text{II}) &\equiv \sum_{j=0}^{+\infty} (\text{I1})_{(kl)j} b_{m'n'j} + b_{k'l'j} (\text{II})_{(mn)j} \quad (\text{B60}) \\ &= \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \{ [-6A'' - 12(A')^2] \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \} - \sum_{j=0}^{+\infty} \mu_j^2 b_{k'l'j} b_{m'n'j} \\ &\quad - \frac{1}{2} \sum_{j=0}^{+\infty} [\mu_k^2 b_{k'l'j} b_{m'n'j} + \mu_l^2 b_{k'l'j} b_{m'n'j} + \mu_m^2 b_{k'l'j} b_{mn'j} + \mu_n^2 b_{k'l'j} b_{m'n'j}] \\ &\quad + \frac{3}{\pi} \int_{-\pi}^{+\pi} d\varphi \{ A' \varepsilon^{+6} [(\mu_k^2 \psi_k) \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_k (\mu_l^2 \psi_l) \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_k \mathcal{D}_l (\mu_m^2 \psi_m) \mathcal{D}_n + \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m (\mu_n^2 \psi_n)] \}. \end{aligned} \quad (\text{B61})$$

This completes our manipulations of the first integral quantity.

Integral 2: The second integral directly yields a vanishing combination of quartic quantities and is defined as

$$(I2)_{k(lmn)} = \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \partial_\varphi \left\{ \left[\frac{3}{8} (\partial_\varphi A) \varepsilon^{-2} (\partial_\varphi \psi_k) - \mu_k^2 \psi_k \right] \varepsilon^{-6} (\partial_\varphi \psi_l) (\partial_\varphi \psi_m) (\partial_\varphi \psi_n) \right\} \quad (B62)$$

$$= \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \partial_\varphi \left\{ \frac{3}{8} A' \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n - \mu_k^2 \varepsilon^{+6} \psi_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \right\}. \quad (B63)$$

As the previous integral, we next carry out the differentiation while making use of the spin-2 mode wave function ($\partial_\varphi \mathcal{D}_x = -\mu_x^2 \varepsilon^{-2} \psi_x$) as we go, and thereby attain

$$(I2)_{k(lmn)} = \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \left\{ \frac{1}{4} \left[\frac{3}{2} A'' + 12(A')^2 - \mu_k^2 \varepsilon^{+2} \right] \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \right. \\ \left. + \frac{1}{4} \varepsilon^{+4} (\mu_k^2 \psi_k) [(\mu_l^2 \psi_l) \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_l (\mu_m^2 \psi_m) \mathcal{D}_n + \mathcal{D}_l \mathcal{D}_m (\mu_n^2 \psi_n)] \right. \\ \left. - \frac{3}{8} A' \varepsilon^{+6} [5(\mu_k^2 \psi_k) \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_k (\mu_l^2 \psi_l) \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_k \mathcal{D}_l (\mu_m^2 \psi_m) \mathcal{D}_n + \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m (\mu_n^2 \psi_n)] \right\}. \quad (B64)$$

Next, we symmetrize over the indices, as to attain all unique combinations of indices:

$$(I2) \equiv (I2)_{k(lmn)} + (I2)_{l(kmn)} + (I2)_{m(kln)} + (I2)_{n(klm)} \quad (B65)$$

$$= \frac{1}{\pi} \int_{-\pi}^{+\pi} d\varphi \left\{ \left[\frac{3}{2} A'' + 12(A')^2 \right] \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \right\} - \frac{1}{4} \bar{\mu}^2 c_{k'l'm'n'} \\ + \frac{1}{2} [\mu_k^2 \mu_l^2 b_{klm'n'} + \mu_k^2 \mu_m^2 b_{kl'mn'} + \mu_k^2 \mu_n^2 b_{kl'm'n} + \mu_l^2 \mu_m^2 b_{k'lmn'} + \mu_l^2 \mu_n^2 b_{k'l'm'n} + \mu_m^2 \mu_n^2 b_{k'l'mn}] \\ - \frac{3}{\pi} \int_{-\pi}^{+\pi} d\varphi \{ A' \varepsilon^{+6} [(\mu_k^2 \psi_k) \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_k (\mu_l^2 \psi_l) \mathcal{D}_m \mathcal{D}_n + \mathcal{D}_k \mathcal{D}_l (\mu_m^2 \psi_m) \mathcal{D}_n + \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m (\mu_n^2 \psi_n)] \}. \quad (B66)$$

Because $(I2)_{k(lmn)}$ vanishes, $(I2)$ vanishes as well. This completes our manipulations of the second integral.

Combining: We finally add (I1) from Eq. (B61) and (I2) from Eq. (B66) to attain a new quantity that, of course, also equals zero. Doing so, we attain

$$0 = (I1) + (I2) \\ = -\frac{9}{2\pi} \int_{-\pi}^{+\pi} d\varphi \{ A'' \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \} - \frac{1}{4} \bar{\mu}^2 c_{k'l'm'n'} - \sum_{j=0}^{+\infty} \mu_j^2 b_{k'l'j} b_{m'n'j} \\ - \frac{1}{2} \sum_{j=0}^{+\infty} [\mu_k^2 b_{k'l'j} b_{m'n'j} + \mu_l^2 b_{k'l'j} b_{m'n'j} + \mu_m^2 b_{k'l'j} b_{m'n'j} + \mu_n^2 b_{k'l'j} b_{m'n'j}] \\ + \frac{1}{2} [\mu_k^2 \mu_l^2 b_{klm'n'} + \mu_k^2 \mu_m^2 b_{kl'mn'} + \mu_k^2 \mu_n^2 b_{kl'm'n} + \mu_l^2 \mu_m^2 b_{k'lmn'} + \mu_l^2 \mu_n^2 b_{k'l'm'n} + \mu_m^2 \mu_n^2 b_{k'l'mn}]. \quad (B67)$$

The two integrals were intentionally weighted so as to ensure all terms containing A' exactly cancel between the two expressions. The resulting expression possesses a couple of important features.

First, Eq. (B67) contains the desired sum $\sum_{j=0}^{+\infty} \mu_j^2 b_{k'l'j} b_{m'n'j}$, which we have already demonstrated generates our ultimate target $\sum_{j=0}^{+\infty} \mu_j^6 a_{klj} a_{mnj}$ via the B-to-A formulas, as made explicit in Eq. (B55).

Second, nearly all other terms in Eq. (B67) can be expressed in terms of a_{klmn} , the $B_{(kl)(mn)}$ and $c_{k'l'm'n'}$ using our existing relations. The only exceptional term is

$$-\frac{9}{2\pi} \int_{-\pi}^{+\pi} d\varphi A'' \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n, \quad (B68)$$

which (as we describe now) contains important information about the radion and the tower of Goldberger-Wise scalars.

In the *unstabilized* model, A'' is a sum of brane-localized Dirac deltas (specifically, $A'' = (kr_c|\varphi|)'' = 2kr_c[\delta(\varphi) - \delta(\varphi - \pi)]$) and thus—because $\mathcal{D}_x = \varepsilon^{-4}\partial_\varphi\psi_x$ vanishes at the branes for each spin-2 state—Eq. (B68) vanishes in the absence of a stabilization mechanism. This contrasts the Goldberger-Wise-stabilized Randall-Sundrum I model, where the equations of motion demand

$$(\partial_\varphi^2 A) = \frac{1}{12}(\partial_\varphi\phi_0)^2 + \frac{1}{3}V_1 r_c \delta(\varphi) + \frac{1}{3}V_2 r_c \delta(\varphi - \pi) \quad (\text{B69})$$

such that Eq. (B68) yields a nonzero contribution directly originating from the nonconstant profile of ϕ_0 through the bulk. That is, a nonzero contribution from Eq. (B68) to our calculations directly reflects the stabilization of the radion.

We may use the equation of motion Eq. (B69) to simplify Eq. (B68). In particular,

$$\begin{aligned} & -\frac{9}{2\pi} \int_{-\pi}^{+\pi} d\varphi A'' \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \\ & = -\frac{3}{8\pi} \int_{-\pi}^{+\pi} d\varphi (\phi_0')^2 \varepsilon^{+8} \mathcal{D}_k \mathcal{D}_l \mathcal{D}_m \mathcal{D}_n \equiv -\frac{3}{8} x_{\phi_0' \phi_0' k' l' m' n'}^{(-8)}. \end{aligned} \quad (\text{B70})$$

To finish rewriting Eq. (B67), all that remains now is the application of the B-to-A formulas, the sum relations, Eq. (B55), and a lot of algebra. The next section presents the results of this process and summarizes the other inelastic sum relations we have derived. The section

thereafter reduces those results to the equivalent elastic rules. Finally, the last section of these Supplemental Material [28] reviews additional unproven rules necessary to cancel any “bad” high-energy behavior from the tree-level helicity-zero $(k, l) \rightarrow (m, n)$ matrix element in the stabilized Randall-Sundrum model.

6. Summary of proven sum rules (inelastic)

All B-type couplings $\{b_{l'm'n}, b_{k'l'mn}\}$ can be eliminated in favor of A-type couplings $\{a_{lmn}, a_{klmn}\}$ and new $B_{(kl)(mn)}$ objects via the B-to-A formulas

$$\begin{aligned} b_{l'm'n} &= \frac{1}{2}[\mu_l^2 + \mu_m^2 - \mu_n^2] a_{lmn}, \\ b_{k'l'mn} &= B_{(kl)(mn)} + \frac{1}{6}[2(\mu_k^2 + \mu_l^2) - (\mu_m^2 + \mu_n^2)] a_{klmn}, \end{aligned} \quad (\text{B71})$$

where the $B_{(kl)(mn)}$ are constrained such that $B_{(km)(ln)} + B_{(kn)(lm)} + B_{(kl)(mn)} = 0$, and are symmetric in each individual pair (k, l) and (m, n) as well as with respect to the pair swap replacement $(k, l) \leftrightarrow (m, n)$. These sums are

$$\sum_{j=0} a_{klj} a_{mnj} = a_{klmn}, \quad (\text{B72})$$

$$\sum_{j=0} \mu_j^2 a_{klj} a_{mnj} = -2B_{(kl)(mn)} + \frac{1}{3} \vec{\mu}^2 a_{klmn}, \quad (\text{B73})$$

$$\sum_{j=0} \mu_j^4 a_{klj} a_{mnj} = 4c_{k'l'm'n'} - 2\vec{\mu}^2 B_{(kl)(mn)} + \frac{1}{6} M_{(kl)(mn)}^{(2)}(1, 1, -1) a_{klmn}, \quad (\text{B74})$$

$$\begin{aligned} \sum_{j=0} \mu_j^6 a_{klj} a_{mnj} &= 5\vec{\mu}^2 c_{k'l'm'n'} + M_{(km)(ln)}^{(2)}(1, 1, 1) B_{(km)(ln)} + M_{(kn)(lm)}^{(2)}(1, 1, 1) B_{(kn)(lm)} \\ &\quad - M_{(kl)(mn)}^{(2)}(0, 1, 0) B_{(kl)(mn)} + \frac{1}{6} M_{(kl)(mn)}^{(3)}(1, 4, -4, 0) a_{klmn} - \frac{3}{2} x_{\phi_0' \phi_0' k' l' m' n'}^{(-8)}, \end{aligned} \quad (\text{B75})$$

where $\vec{\mu}^2 \equiv \mu_k^2 + \mu_l^2 + \mu_m^2 + \mu_n^2$ and

$$c_{k'l'm'n'} \equiv \frac{1}{\pi} \int d\varphi \varepsilon^{-6} (\partial_\varphi \psi_k) (\partial_\varphi \psi_l) (\partial_\varphi \psi_m) (\partial_\varphi \psi_n), \quad (\text{B76})$$

$$x_{\phi_0' \phi_0' k' l' m' n'}^{(-8)} \equiv \frac{1}{\pi} \int d\varphi \varepsilon^{-8} (\partial_\varphi \phi_0')^2 (\partial_\varphi \psi_k) (\partial_\varphi \psi_l) (\partial_\varphi \psi_m) (\partial_\varphi \psi_n). \quad (\text{B77})$$

The last two sum rules can be combined so as to cancel all factors of $c_{k'l'm'n'}$ and thereby yield

$$\begin{aligned} \sum_{j=0} \mu_j^4 \left(\mu_j^2 - \frac{5}{4} \vec{\mu}^2 \right) a_{klj} a_{mnj} &= \frac{1}{4} M_{(kl)(mn)}^{(2)}(5, 1, 10) B_{(kl)(mn)} + M_{(km)(ln)}^{(2)}(1, 1, 1) B_{(km)(ln)} + M_{(kn)(lm)}^{(2)}(1, 1, 1) B_{(kn)(lm)} \\ &\quad - \frac{1}{24} M_{(kl)(mn)}^{(3)}(1, -1, 16, 0) a_{klmn} - \frac{3}{2} x_{\phi_0' \phi_0' k' l' m' n'}^{(-8)}. \end{aligned} \quad (\text{B78})$$

These equations extend and generalize the sum rules derived in [11].

7. Summary of proven sum rules (elastic)

Oftentimes, we are particularly interested in the *elastic* massive spin-2 KK mode scattering process, wherein $k = l = m = n (\neq 0)$ and the relations of the previous subsections simplify. Consider, for example, the B 's constraint in this context:

$$B_{(km)(ln)} + B_{(kn)(lm)} + B_{(kl)(mn)} = 0 \xrightarrow{\text{elastic}} B_{(nn)(nn)} = 0 \quad (\text{B79})$$

such that all of the B 's become identical and vanish. The relevant B-to-A formulas become

$$\begin{aligned} b_{n'n'j} &= \frac{1}{2} [2\mu_n^2 - \mu_j^2] a_{nnj}, & b_{j'n'n} &= \frac{1}{2} \mu_j^2 a_{nmj}, \\ b_{n'n'nn} &= \frac{1}{3} \mu_n^2 a_{nnnn}, \end{aligned} \quad (\text{B80})$$

whereas the sum rules reduce to

$$\sum_j a_{nnj}^2 = a_{nnnn}, \quad (\text{B81})$$

$$\sum_j \mu_j^2 a_{nnj}^2 = \frac{4}{3} \mu_n^2 a_{nnnn}, \quad (\text{B82})$$

$$\sum_j \mu_j^4 a_{nnj}^2 = 4c_{n'n'n'n'} + \frac{4}{3} \mu_n^4 a_{nnnn}, \quad (\text{B83})$$

$$\sum_j \mu_j^6 a_{nnj}^2 = 20\mu_n^2 c_{n'n'n'n'} + \frac{4}{3} \mu_n^6 a_{nnnn} - \frac{3}{2} x_{\phi_0 \phi_0}^{(-8)}, \quad (\text{B84})$$

with the last two expressions combining to yield

$$\sum_j [\mu_j^2 - 5\mu_n^2] \mu_j^4 a_{nnj}^2 = -\frac{16}{3} \mu_n^6 a_{nnnn} - \frac{3}{2} x_{\phi_0 \phi_0}^{(-8)}. \quad (\text{B85})$$

8. Unproven rules

The aforementioned rules are insufficient on their own for ensuring cancellations of the $(k, l) \rightarrow (m, n)$ matrix element [which naively contains $\mathcal{O}(s^5)$ terms] down to $\mathcal{O}(s)$ growth. This is true in both the fully elastic ($k = l = m = n$) and more general cases.

In the fully elastic case, only one additional rule is required:

$$3 \left[9 \sum_{i=0}^{+\infty} a_{n'n'n'(i)}^2 - \mu_n^2 \mu_n^2 a_{nn0}^2 \right] = 15c_{n'n'n'n'} + 2\mu_n^4 a_{nnnn}. \quad (\text{B86})$$

The inelastic case provides a generalization of this rule, as well as two additional rules we have yet to prove analytically. These analytic rules have been attained by calculating the full $(k, l) \rightarrow (m, n)$ matrix element (a nontrivial task), asymptotically series expanding that matrix element in s down to $\mathcal{O}(s^{3/2})$ (also nontrivial; note odd powers of s automatically vanish for this particular process), applying the sum rules we previously derived, and demanding coefficients of any s^σ for $\sigma > 1$ vanish.

Having done so, we find cancellations of “bad” high-energy behavior additionally require

$$\begin{aligned} 6B_{(kl)(mn)} &= (\mu_k^2 - \mu_m^2)(\mu_l^2 - \mu_n^2) \sum_{j>0} \frac{a_{kmj} a_{lnj}}{\mu_j^2} \\ &+ (\mu_k^2 - \mu_n^2)(\mu_l^2 - \mu_m^2) \sum_{j>0} \frac{a_{knj} a_{lmj}}{\mu_j^2} \end{aligned} \quad (\text{B87})$$

to cancel $\mathcal{O}(s^4)$ growth and, noting the KK indices (k, l, m) are cycled through from term to term,

$$\begin{aligned} 0 &= (\mu_k^2 - \mu_l^2)(\mu_m^2 - \mu_n^2) \sum_{j>0} \frac{a_{klj} a_{mnj}}{\mu_j^2} \\ &+ (\mu_l^2 - \mu_m^2)(\mu_k^2 - \mu_n^2) \sum_{j>0} \frac{a_{lmj} a_{knj}}{\mu_j^2} \\ &+ (\mu_m^2 - \mu_k^2)(\mu_l^2 - \mu_n^2) \sum_{j>0} \frac{a_{mkj} a_{lnj}}{\mu_j^2} \end{aligned} \quad (\text{B88})$$

to cancel $\mathcal{O}(s^3)$ growth.

To simplify writing expressions such as those above, define

$$L_{kl;mn} = (\mu_k^2 - \mu_l^2)(\mu_m^2 - \mu_n^2) \sum_{j>0} \frac{a_{klj} a_{mnj}}{\mu_j^2}. \quad (\text{B89})$$

$L_{kl;mn}$ is antisymmetric under $k \leftrightarrow l$ and $m \leftrightarrow n$, and is symmetric under $kl \leftrightarrow mn$. The previously listed new sum rules can thus be written succinctly as

$$6B_{(kl)(mn)} = L_{km;ln} + L_{kn;lm}, \quad (\text{B90})$$

$$0 = L_{kl;mn} + L_{lm;kn} + L_{mk;ln}. \quad (\text{B91})$$

This latter sum rule is mathematically distinct from the defining constraint of $B_{(kl)(mn)}$ (i.e., that the sum of all unique B vanishes). Note that $L_{nn;nn} = 0$, thus explaining the absence of these relations when deriving our elastic sum rules.

Last, the $\mathcal{O}(s^3)$ cancellations also necessitate the following generalization of the elastic radion rule [Eq. (B86)]:

$$\begin{aligned}
12 \left[9 \sum_{i=0}^{+\infty} a_{k'l'(i)} a_{m'n'(i)} - \mu_k^2 \mu_m^2 a_{kl0} a_{mn0} \right] &= 60 c_{k'l'm'n'} + \frac{1}{2} M_{(kl)(mn)}^{(2)}(4, -8, 5) a_{klmn} \\
&- 3(\mu_k^2 - \mu_l^2)^2 (\mu_m^2 - \mu_n^2)^2 \sum_{j>0} \frac{a_{klj} a_{mnj}}{\mu_j^4} + 2\vec{\mu}^2 [L_{km;ln} + L_{kn;lm}] \\
&- 3[(\mu_k^2 + \mu_l^2)(\mu_m^2 - \mu_n^2)^2 + (\mu_k^2 - \mu_l^2)^2 (\mu_m^2 + \mu_n^2)] \sum_{j>0} \frac{a_{klj} a_{mnj}}{\mu_j^2}, \quad (\text{B92})
\end{aligned}$$

where $M_{(nn)(nn)}^{(2)}(4, -8, 5) = 8\mu_n^4$. In the unstabilized inelastic calculation, an identical rule is attained, but with the sum $\sum_i a_{(i)k'l'} a_{(i)m'n'}$ replaced simply by $a_{(0)k'l'} a_{(0)m'n'}$. This completes the rules necessary to ensure cancellations down to $\mathcal{O}(s)$ for 2-to-2 spin-2 mode scattering in the stabilized Randall-Sundrum model.

APPENDIX C: EIGENVALUES AND EIGENFUNCTIONS OF SPIN-2 AND SPIN-0 MODES

The solutions to the SL problem for the spin-0 and the spin-2 parts of the stabilized RS model determine the eigenvalues and eigenfunctions. Here we outline two related methods of computing the eigenvalues and eigenfunctions for both the spin-0 and the spin-2 SL problem in perturbation theory. In the following we introduced the standard Rayleigh-Schrödinger perturbation theory in the context of a SL problem. Here, the perturbed wave functions are expressed as an infinite series in unperturbed wave functions. On the other hand, being able to have closed form expressions for the perturbed wave functions is extremely useful, especially for the numerical part of our analysis. This leads us to solving the perturbed SL problem directly, by solving an inhomogeneous differential equation. In the end the normalized wave functions derived in either of these methods are identical, and calculating the wave function using these two methods serves as a cross-check of our results.

1. Perturbation theory and a general Sturm-Liouville problem

Here we discuss the application of Rayleigh-Schrödinger perturbation theory to a general Sturm-Liouville problem, including one in which the weight function is also perturbed. We compute the first-order shifts to the eigenvalues and eigenfunctions, and we demonstrate that completeness holds to the appropriate order. In Sec. C 1 a we show how the perturbed eigenfunctions can be calculated as a linear combination of unperturbed eigenfunctions. In practice performing an infinite sum of wave functions to determine the perturbed wave function is not computationally efficient, so in Sec. C 1 b we outline an equivalent method of determining the perturbed wave functions as closed form

expressions and show how it is related to Rayleigh-Schrödinger perturbation theory.

a. Mass corrections

Consider a generic SL problem for the Kaluza-Klein modes, which is of the form

$$\tilde{\mathcal{L}}\psi_n = -\lambda_n \tilde{\rho} \psi_n, \quad (\text{C1})$$

where $\tilde{\mathcal{L}}$ is the SL operator (given appropriate boundary conditions) acting on eigenfunctions ψ_n with eigenvalues $\tilde{\lambda}_n$ and a weight factor $\tilde{\rho}$. The solutions to the SL problem are orthogonal with respect to the weight factor $\tilde{\rho}$

$$\frac{1}{\pi} \int_{-\pi}^{\pi} d\varphi \tilde{\rho}(\varphi) \psi_k(\varphi) \psi_l(\varphi) = \delta_{k,l}. \quad (\text{C2})$$

These solutions then satisfy the completeness relation²⁶

$$\sum_{\ell} \tilde{\rho}(\varphi) \psi_{\ell}(\varphi) \psi_{\ell}(\varphi') = \pi \delta(\varphi - \varphi'). \quad (\text{C3})$$

Depending on the nature of the SL problem, the boundary conditions can be Dirichlet or Neumann as pointed out in the main body of the paper. For the rest of this appendix, we will drop the argument φ in wave functions and weight factors for simplicity.

In perturbation theory, the SL operator and weight function can be expanded as $\tilde{\mathcal{L}} = \mathcal{L} + \delta\mathcal{L}$ and $\tilde{\rho} = \rho + \delta\rho$, while we expand the eigenvalue $\tilde{\lambda}_n = \lambda_n^{(0)} + \lambda_n^{(1)} + \lambda_n^{(2)} + \dots$. Here both \mathcal{L} and $\delta\mathcal{L}$ are of Sturm-Liouville form:

$$\mathcal{L} = \frac{d}{d\varphi} \left[p \frac{d}{d\varphi} \right] + q, \quad (\text{C4})$$

$$\delta\mathcal{L} = \frac{d}{d\varphi} \left[\delta p \frac{d}{d\varphi} \right] + \delta q. \quad (\text{C5})$$

In our problems, the perturbations δp , δq , and $\delta\rho$ come from expanding Eqs. (45) and (52), respectively, in powers

²⁶The symmetry of the δ -function implies that the argument of ρ in the sum could be either φ or φ' .

of ϵ .²⁷ We then expand the eigenfunction in perturbation theory as

$$\psi_n = \psi_n^{(0)} + \psi_n^{(1)} + \psi_n^{(2)} + \dots \quad (\text{C6})$$

As usual in perturbation theory, we expand the first-order perturbed wave function as a sum of unperturbed wave functions,

$$\psi_n^{(1)} = \sum_{m=0}^{\infty} C_{nm} \psi_m^{(0)}. \quad (\text{C7})$$

The coefficients C_{nm} can be determined using perturbation theory, as will be described later. Here, as usual in Rayleigh-Schrödinger perturbation theory, we assume that $\psi_n^{(1)}$ is chosen to be orthogonal to $\psi_n^{(0)}$.

To the lowest order, the SL equation reads

$$\mathcal{L}\psi_n^{(0)} = -\lambda_n^{(0)} \rho \psi_n^{(0)}, \quad (\text{C8})$$

where $\lambda_n^{(0)}$ is the lowest order (unperturbed) eigenvalue. Additionally, $\psi_n^{(0)}$ is the lowest order eigenfunction.

Expanding the SL equation to first order, and using the fact the lowest order SL problem satisfies Eq. (C8), we obtain, to first order,

$$\mathcal{L}\psi_n^{(1)} + \delta\mathcal{L}\psi_n^{(0)} = -(\lambda_n^{(0)} \delta\rho \psi_n^{(0)} + \lambda_n^{(1)} \rho \psi_n^{(0)} + \lambda_n^{(0)} \rho \psi_n^{(1)}). \quad (\text{C9})$$

Multiplying the first-order perturbed equation by ψ_n^0 and integrating,

$$\int d\varphi \psi_n^{(1)} \mathcal{L}\psi_n^{(0)} + \int d\varphi \delta\mathcal{L}(\psi_n^{(0)})^2 = - \int d\varphi (\lambda_n^{(0)} \delta\rho (\psi_n^{(0)})^2 + \lambda_n^{(1)} \rho (\psi_n^{(0)})^2 + \lambda_n^{(0)} \rho \psi_n^{(1)} \psi_n^{(0)}). \quad (\text{C10})$$

In the first term in the equation above, we have used the fact that the operator \mathcal{L} is self-adjoint. Using Eq. (C8) and rearranging, we obtain

$$\lambda_n^{(1)} = -\frac{1}{\pi} \left(\int_{-\pi}^{\pi} d\varphi \delta\rho (\psi_n^{(0)})^2 + \int_{-\pi}^{\pi} d\varphi \psi_n^{(0)} \delta\mathcal{L}\psi_n^{(0)} \right). \quad (\text{C11})$$

Since $\delta\mathcal{L} = \frac{d}{dx}[\delta p] + \delta q$, we can integrate the above equation by parts to obtain

$$\lambda_n^{(1)} = -\frac{1}{\pi} \left[- \int_{-\pi}^{\pi} d\varphi \delta p \left(\frac{d\psi_n^{(0)}}{d\varphi} \right)^2 + \int_{-\pi}^{\pi} d\varphi \delta q (\psi_n^{(0)})^2 + \int_{-\pi}^{\pi} d\varphi \delta\rho (\psi_n^{(0)})^2 \right]. \quad (\text{C12})$$

Now that we have the perturbed eigenvalue, we can proceed to calculate the perturbed eigenfunctions. We describe two methods to do this. The first one involves directly solving the nonhomogenous differential equation in Eq. (C9). The second one makes use of standard Rayleigh-Schrödinger perturbation theory. In the end, both methods lead to the same eigenfunctions, and the use of these two methods serves as a cross-check of our results.

b. Solving the inhomogeneous differential equation using variation of parameters

In this first method, one simply solves the nonhomogeneous differential equation that is derived by substituting into Eq. (C9) the unperturbed eigenvalue and eigenfunction

$$\left[\frac{d}{d\varphi} \left(p \frac{d}{d\varphi} \right) + q - \lambda_n^{(0)} \rho \right] \psi_n^{(1)} = \left[-\frac{d}{d\varphi} \left(\delta p \frac{d}{d\varphi} \right) - \delta q + \lambda_n^{(0)} \delta\rho + \lambda_n^{(1)} \rho \right] \psi_n^{(0)}. \quad (\text{C13})$$

To solve this equation for the spin-2 KK modes, we have used the method of variation of parameters. Using the solution $\psi_n^{(1)}$ found using this method, the wave function $\psi_n^{(0)} + \psi_n^{(1)}$ must then be normalized with respect to $\rho + \delta\rho$ as follows:

$$\tilde{\psi}_n^{(\text{normalized})} = \psi_n^{(0)} + \left[\psi_n^{(1)} - \frac{\psi_n^{(0)}}{\pi} \int d\varphi' \left(\rho \psi_n^{(0)} \psi_n^{(1)} + \frac{1}{2} \delta\rho (\psi_n^{(0)})^2 \right) \right]. \quad (\text{C14})$$

²⁷Note that the spin-2 system in Eq. (45) yields a perturbation expansion in ϵ^2 , whereas that for the spin-0 system in Eq. (52) gives an expansion in powers of ϵ .

In the following we describe how to determine the same wave function using the Rayleigh-Schrödinger perturbation theory. We reiterate that the wave functions determined in either method are identical and that the advantage of determining the wave function in this way is that we get closed form solutions for the perturbed wave functions and masses.

c. Wave functions in Rayleigh-Schrödinger perturbation theory

The perturbed wave function is determined as a sum over the unperturbed wave function as shown in Eq. (C7). To determine the coefficients C_{nm} , we multiply the perturbed SL equation (C9) by $\psi_m^{(0)}$ ($m \neq n$) and integrate

$$-\lambda_m^{(0)} \pi C_{nm} + \int_{-\pi}^{\pi} d\varphi \psi_m^{(0)} (\delta\mathcal{L}\psi_m^{(0)}) = -\left(\lambda_n^{(0)} \int_{-\pi}^{\pi} d\varphi \psi_m^{(0)} \delta\rho \psi_n^{(0)} + \lambda_n^{(0)} \pi C_{nm} \right), \quad (\text{C15})$$

leading to

$$C_{nm} = -\frac{1}{\pi} \frac{(\lambda_n^{(0)} \int_{-\pi}^{\pi} d\varphi \psi_m^{(0)} \delta\rho \psi_n^{(0)} + \int_{-\pi}^{\pi} d\varphi \psi_m^{(0)} (\delta\mathcal{L}\psi_m^{(0)}))}{\lambda_n^{(0)} - \lambda_m^{(0)}}. \quad (\text{C16})$$

Using the definition of $\delta\mathcal{L}$ and integrating by parts, we get

$$C_{nm} = -\frac{1}{\pi} \left[\frac{-\int_{-\pi}^{\pi} d\varphi \delta\rho \frac{d\psi_n^{(0)}}{d\varphi} \frac{d\psi_m^{(0)}}{d\varphi} + \int_{-\pi}^{\pi} d\varphi \delta q \psi_n^{(0)} \psi_m^{(0)} + \lambda_n^{(0)} \int_{-\pi}^{\pi} d\varphi \delta\rho \psi_n^{(0)} \psi_m^{(0)} \right]_{\lambda_n^{(0)} - \lambda_m^{(0)}}. \quad (\text{C17})$$

In Rayleigh-Schrödinger perturbation theory, we usually assume that the first-order perturbed solutions are orthogonal to the lowest order solution so that $C_{nn} = 0$. However, in the presence of a perturbation to the weight function, that is no longer the case. Here, to obtain the coefficient C_{nn} , we use the normalization condition

$$\int_{-\pi}^{\pi} d\varphi \tilde{\rho} \tilde{\psi}_n^2 = \int_{-\pi}^{\pi} d\varphi (\rho + \delta\rho) \left(\psi_n^{(0)} + \sum_m C_{nm} \psi_m^{(0)} \right)^2 = \pi, \quad (\text{C18})$$

which, to first order, implies

$$C_{nn} = -\frac{1}{2\pi} \int_{-\pi}^{\pi} d\varphi \delta\rho (\psi_n^{(0)})^2. \quad (\text{C19})$$

We have checked, numerically, that the wave functions derived using Eqs. (C7) and (C17) are identical to the ones derived using Eqs. (C13) and (C14).

2. Wave function and masses of KK modes in the DFGK model

Here we present the wave functions and mass corrections for the spin-2 as well as the scalar sector for the DFGK model in the stiff-wall limit. These expressions are derived by solving the differential equations described in Sec. C and specifically using Eqs. (C12) and (C13). The expressions presented here are relevant for the large kr_c limit. The general expressions, valid for all kr_c , are quite cumbersome and are provided in supplementary *Mathematica* files on GitHub [28].

a. Spin-2 mass and wave function corrections

To verify sum rules to order ϵ^2 we need the spin-2 wave function and masses to order ϵ^2 . We start by expanding the spin-2 Sturm-Liouville equation in (7) up to order ϵ^2 . We also expand the wave function and masses, as described earlier, to order ϵ^2

$$\psi_n = \psi_n^{(0)} + \psi_n^{(2)} + \dots, \quad (\text{C20})$$

$$\mu_n^2 = (\mu_n^{(0)})^2 + \delta\mu_n^2 + \dots. \quad (\text{C21})$$

We have dropped the $\psi_n^{(1)}$ term since the corrections to the spin-2 Sturm-Liouville problem start at order ϵ^2 as can be seen from its expanded form below

$$0 = [\partial_\varphi^2 - 4\tilde{k}r_c \partial_\varphi + (\mu_n^{(0)})^2 e^{2\tilde{k}r_c \varphi}] \psi_n^{(0)} + \epsilon^2 \{ [-8\varphi \partial_\varphi + e^{2\tilde{k}r_c \varphi} (\delta\mu_n^2 + 2\varphi^2 (\mu_n^{(0)})^2)] \psi_n^{(0)} + [\partial_\varphi^2 - 4\tilde{k}r_c \partial_\varphi + e^{2\tilde{k}r_c \varphi} (\mu_n^{(0)})^2] \psi_n^{(2)} \} + \mathcal{O}(\epsilon^3). \quad (\text{C22})$$

Here we see that the leading term for $\psi_n^{(0)}$ is the usual one that we encounter in the unstabilized limit. Solutions to the leading order differential equation are well known and can be found in Ref. [11]. We reproduce some of these results here later. After solving for $\psi_n^{(0)}$, we then proceed to solve the above differential equation at order ϵ^2 .

b. The massless graviton to order ϵ^2

The massless graviton is the easiest, since it does not acquire a mass, and its wave function is derived by setting

$\mu_0^{(0)} = \delta\mu_0 = 0$. The wave function is a constant, and any ϵ dependence comes from the normalization condition (8). The normalized wave function for the massless graviton to order ϵ^2 is of the form

$$\begin{aligned} \psi_0 &= \left[\frac{1}{\pi} \int d\varphi e^{-2A} \right]^{-1/2} \\ &= e^{\pi\tilde{k}r_c} \sqrt{\frac{\pi\tilde{k}r_c}{e^{2\pi\tilde{k}r_c} - 1}} \\ &\quad + \epsilon^2 \frac{\sqrt{\pi} e^{\pi\tilde{k}r_c} [-2\pi^2\tilde{k}^2 r_c^2 - 2\pi\tilde{k}r_c + e^{2\pi\tilde{k}r_c} - 1]}{8r_c^{3/2} [\tilde{k}(e^{2\pi\tilde{k}r_c} - 1)]^{3/2}} \\ &\quad + \mathcal{O}(\epsilon^3). \end{aligned} \quad (C23)$$

c. Massive spin-2 modes to order ϵ^2

Before we present results on the perturbed wave functions and masses of the massive spin-2 modes, we remind the reader about the unperturbed wave functions and masses that are identical to the unstabilized case. Since the full expressions can be quite lengthy, we present only results that are valid in the large $\tilde{k}r_c$ limit and provide the full expressions in supplementary *Mathematica* files on GitHub [28]. The wave function for the massive spin-2 modes to order ϵ^0 is the same as those derived in the unstabilized RS model and in the large $\tilde{k}r_c$ limit is of the form

$$\begin{aligned} \psi_n^{(0)} &= e^{2\varphi\tilde{k}r_c} \sqrt{\frac{\pi r_c \tilde{k}}{e^{2\tilde{k}r_c \pi} J_2(j_{1,n})^2 - J_2(e^{-\pi\tilde{k}r_c} j_{1,n})^2}} \\ &\quad \times J_2(e^{\tilde{k}(\varphi r_c - \pi r_c)} j_{1,n}). \end{aligned} \quad (C24)$$

Here J_i are Bessel-J functions. The masses of the spin-2 modes are determined by solving a transcendental equation that is derived from the Neumann boundary condition satisfied by the spin-2 modes:

$$J_1\left(\frac{e^{\pi\tilde{k}r_c} \mu_n}{\tilde{k}r_c}\right) Y_1\left(\frac{\mu_n}{\tilde{k}r_c}\right) - J_1\left(\frac{\mu_n}{\tilde{k}r_c}\right) Y_1\left(\frac{e^{\pi\tilde{k}r_c} \mu_n}{\tilde{k}r_c}\right) = 0. \quad (C25)$$

In the large $\tilde{k}r_c$ limit, the solution to the transcendental equation reduces to a simple form:

$$m_n = \frac{\mu_n^{(0)}}{r_c} = \tilde{k} e^{-\pi\tilde{k}r_c} j_{1,n}. \quad (C26)$$

Here $j_{1,n}$ are the roots of J_1 .

Substituting the above form of the leading order spin-2 wave function and masses into Eq. (C22), we end up with a nonhomogeneous differential equation which can be solved using the method of variation of parameters. Alternately, the same differential equation can also be derived from Eq. (C13) and in the large $\tilde{k}r_c$ limit is of the form given below:

$$[\partial_\varphi^2 - 4\tilde{k}r_c \partial_\varphi + (\mu_n^{(0)})^2 e^{2\tilde{k}r_c \varphi}] \psi_n^{(2)} = 2c_1 e^{3\varphi\tilde{k}r_c} \left[8\mu_n^{(0)} \varphi J_1\left(\frac{e^{\tilde{k}r_c \varphi} \mu_n^{(0)}}{\tilde{k}r_c}\right) - e^{\tilde{k}r_c \varphi} (\delta\mu_n^2 + 2(\mu_n^{(0)})^2 \varphi^2) J_2\left(\frac{e^{\tilde{k}r_c \varphi} \mu_n^{(0)}}{\tilde{k}r_c}\right) \right]. \quad (C27)$$

Here c_1 corresponds to the normalization of the leading order wave function $\psi_n^{(0)}$. Below we write down the perturbation of the spin-2 wave function $\psi_n^{(2)}$. The resulting expressions at order ϵ^2 are quite lengthy and are also provided in supplementary *Mathematica* files on GitHub [28]:

$$\begin{aligned} \psi_n^{(2)} &= \frac{c_1 Y_2(z)}{384\mu_n^4} \epsilon^2 \left\{ \mu_n^2 \pi z^6 {}_3F_4\left(\frac{3}{2}, 2, 2; 1, 3, 3, 4; -z^2\right) [1 - 4 \log(\beta z)] \right. \\ &\quad + 2\pi z^6 {}_2F_3\left(\frac{3}{2}, 2; 1, 3, 4; -z^2\right) [2 \log(\beta z) - 1] \\ &\quad + 384\pi z^2 \left[{}_1F_2\left(\frac{1}{2}; 1, 1; -z^2\right) - 2 {}_1F_2\left(\frac{1}{2}; 1, 2; -z^2\right) \right] \log(\beta z) + \log(\beta) \Big] \\ &\quad + \pi z^6 {}_4F_5\left(\frac{3}{2}, 2, 2, 2; 1, 3, 3, 3, 4; -z^2\right) \\ &\quad + 192\sqrt{\pi} z^2 \left[G_{2,4}^{2,1}\left(z^2 \middle| \frac{1}{2}, 1 \right) - 2G_{2,4}^{2,1}\left(z^2 \middle| \frac{1}{2}, 1 \right) \right] \\ &\quad - 48\pi \mu_n^2 z^4 J_2(z)^2 [4\beta^2 \delta\mu_n^2 + (2[\log(\beta z) - 1] \log(\beta z) + 1)] \\ &\quad \left. + 48\pi \beta^2 z^4 J_1(z) J_3(z) [4\beta^2 \delta\mu_n^2 + (2[\log(\beta z) - 1] \log(\beta z) + 1)] \right\} \end{aligned}$$

$$\begin{aligned}
 & + \frac{z^2 J_2(z)}{8\mu_n^2} \left\{ 2\sqrt{\pi} G_{3,5}^{2,2} \left(z, \frac{1}{2} \middle| \begin{matrix} 1, \frac{3}{2}, \frac{1}{2} \\ 1, 3, -1, 0, \frac{1}{2} \end{matrix} \right) [2\beta^2 \delta\mu_n^2 + \log^2(\beta z)] \right. \\
 & + 2\mu_n^2 \sqrt{\pi} \left[4G_{3,5}^{3,1} \left(z, \frac{1}{2} \middle| \begin{matrix} \frac{1}{2}, -\frac{1}{2}, 1 \\ 0, 0, 2, -1, -\frac{1}{2} \end{matrix} \right) - G_{4,6}^{2,3} \left(z, \frac{1}{2} \middle| \begin{matrix} 1, 1, \frac{3}{2}, \frac{1}{2} \\ 1, 3, -1, 0, 0, \frac{1}{2} \end{matrix} \right) \right] \log(\beta z) \\
 & + \mu_n^2 \sqrt{\pi} \left[4G_{4,6}^{4,1} \left(z, \frac{1}{2} \middle| \begin{matrix} \frac{1}{2}, -\frac{1}{2}, 1, 1 \\ 0, 0, 0, 2, -1, -\frac{1}{2} \end{matrix} \right) + G_{5,7}^{2,4} \left(z, \frac{1}{2} \middle| \begin{matrix} 1, 1, 1, \frac{3}{2}, \frac{1}{2} \\ 1, 3, -1, 0, 0, 0, \frac{1}{2} \end{matrix} \right) \right] \\
 & \left. + 8\mu_n^2 \log(z) [2 \log(\beta) + \log(z)] \right\} + \beta^2 z^2 [c_3 J_2(z) + c_4 Y_2(z)]. \tag{C28}
 \end{aligned}$$

Here $\beta \equiv \tilde{k}r_c/\mu_n$, $z \equiv (\mu_n/\tilde{k}r_c)e^{\tilde{k}r_c\varphi}$, $G_{i,j}^{k,l}$ are Meijer-G functions [37], ${}_iF_j$ are hypergeometric functions, c_1 is a normalization constant given in Eq. (C19), and c_3 and c_4 are constants that are determined from the Neumann boundary condition that the wave function must satisfy. Additionally, we provide these expressions as well as those valid also for arbitrary values of $\tilde{k}r_c$ in supplementary *Mathematica* files on GitHub [28].

Corrections to the mass to order ϵ^2 are calculated using Eq. (C12). The large kr_c limit is given in terms of Bessel functions and hypergeometric functions as follows:

$$\begin{aligned}
 \delta\mu_n^2 = & -\frac{c_1^2 \epsilon^2}{48\tilde{k}} \left\{ (j_{1,n})^2 \left[16J_0(j_{1,n})^2 (6\pi\tilde{k}r_c - 1) - 48J_2(j_{1,n})^2 (2\pi\tilde{k}r_c (\pi\tilde{k}r_c - 1) + 1) \right. \right. \\
 & + 96{}_3F_4 \left(1, 1, \frac{3}{2}; 2, 2, 2, 2; -(j_{1,n})^2 \right) - 96{}_3F_4 \left(1, 1, \frac{3}{2}; 2, 2, 2, 3; -(j_{1,n})^2 \right) \\
 & \left. \left. + (j_{1,n})^2 {}_4F_5 \left(\frac{3}{2}, 2, 2, 2; 1, 3, 3, 3, 4; -(j_{1,n})^2 \right) \right] - 32(3J_0(j_{1,n})^2 - 3) \right\}. \tag{C29}
 \end{aligned}$$

3. Spin-0 mass and wave functions in the DFGK model

To verify the sum rules to order ϵ^2 we need the radion mass squared and wave function corrections to order ϵ^2 . On the other hand, due to the normalization condition in Eq. (15) in the stiff-wall limit, the GW scalar wave functions do not have a ϵ^0 piece, but instead start at order ϵ . Therefore, it is only necessary to calculate their masses and wave functions to leading order in ϵ . To determine the wave functions to the required order in ϵ , we start with the Sturm-Liouville problem for the scalar modes defined in Eq. (12), and we perform an expansion of the same up to order ϵ^2 as follows:

$$\begin{aligned}
 & [\partial_\varphi^2 + 2\tilde{k}r_c \partial_\varphi + (\mu_{(n)}^{(0)})^2 e^{2\tilde{k}r_c\varphi}] \gamma_n^{(0)} + \epsilon \left[\frac{4\sqrt{6}}{\phi_1} \partial_\varphi \right] \gamma_n^{(0)} \\
 & + \epsilon^2 \{ [4\varphi \partial_\varphi - 4 + e^{2\tilde{k}r_c\varphi} (\delta\mu_{(n)}^2 + 2\varphi^2 (\mu_{(n)}^{(0)})^2)] \gamma_n^{(0)} + [\partial_\varphi^2 + 2\tilde{k}r_c \partial_\varphi + e^{2\tilde{k}r_c\varphi} (\mu_{(n)}^{(0)})^2] \gamma_n^{(2)} \} = 0. \tag{C30}
 \end{aligned}$$

a. Radion wave function and mass to order ϵ^2

We expand the wave function and mass in perturbation theory as described earlier. For the massless radion we start with the ansatz

$$\begin{aligned}
 \gamma_n & = \gamma_n^{(0)} + \gamma_n^{(2)} + \dots, \\
 \mu_{(n)}^2 & = (\mu_{(n)}^{(0)})^2 + \delta\mu_{(n)}^2 + \dots. \tag{C31}
 \end{aligned}$$

Note the absence of the order ϵ term in the expansion above although there is an explicit order ϵ term in the differential equation. It is easy to see that up to order ϵ , the radion wave

function is constant and only acquires nontrivial dependence at order ϵ^2 . We can substitute the above expansion into Eq. (C30) and solving for $\gamma_n^{(0)}$, $\gamma_n^{(2)}$, $(\mu_{(n)}^{(0)})^2$, and $\delta\mu_{(n)}^2$ order by order. This amounts to determining the unperturbed wave function and using Eq. (C13) to determine the perturbed wave function by solving the resulting nonhomogeneous differential equation

$$[\partial_\varphi^2 + 2\tilde{k}r_c \partial_\varphi + (\delta\mu_{(n)}^2 e^{2\tilde{k}r_c\varphi} - 4)] \gamma_0^{(0)} = 0. \tag{C32}$$

We find the normalized radion mass and wave function to order ϵ^2 to be

$$\mu_{(0)} = 2\epsilon \sqrt{\frac{2}{1 + e^{2\pi\tilde{k}r_c}}} \quad (\text{C33})$$

$$\begin{aligned} \gamma_0 = & \sqrt{\frac{\pi\tilde{k}r_c}{e^{2\pi\tilde{k}r_c} - 1}} + \epsilon^2 \frac{\sqrt{\pi(e^{2\pi\tilde{k}r_c} - 1)}}{6(\tilde{k}r_c)^{3/2}(e^{4\pi\tilde{k}r_c} - 1)^2} [-5 + 6e^{2\varphi\tilde{k}r_c} + 3e^{2\pi\tilde{k}r_c} - 6e^{(2\varphi+4\pi)\tilde{k}r_c} - 6\tilde{k}r_c(e^{2\pi\tilde{k}r_c} + 1)^2 \\ & \times (e^{2\pi\tilde{k}r_c}(\pi^2\tilde{k}r_c - 2\varphi + \pi) + 2\varphi) + 5e^{6\pi\tilde{k}r_c} - 6e^{2(\pi-\varphi)\tilde{k}r_c} + 6e^{2(3\pi-\varphi)\tilde{k}r_c} - 3e^{4\pi\tilde{k}r_c}] + \mathcal{O}(\epsilon^3). \end{aligned} \quad (\text{C34})$$

Note that the expression is valid for arbitrary values of $\tilde{k}r_c$. It is possible to simplify this expression further in the large $\tilde{k}r_c$ limit to

$$\gamma_0|_{\tilde{k}r_c \gg 1} = \gamma_0^{(0)} \left\{ 1 + \frac{\epsilon^2}{4\tilde{k}^2 r_c^2} \left(-\tilde{k}r_c(\pi^2\tilde{k}r_c - 2\varphi + \pi) + e^{-2\varphi\tilde{k}r_c} - e^{2(\varphi-\pi)\tilde{k}r_c} + \frac{5}{6} \right) \right\} + \mathcal{O}(\epsilon^3). \quad (\text{C35})$$

In the large $\tilde{k}r_c$ limit, the wave function for the unstabilized radion is $\gamma_0^{(0)} = e^{-\pi\tilde{k}r_c} \sqrt{\pi\tilde{k}r_c}$.

b. GW-scalar mass and wave function to leading order

As remarked upon earlier, to verify the sum rules to order ϵ^2 , we need the GW-scalar wave function to order ϵ , which, due to the normalization condition in Eq. (15), is in fact the leading order for the massive GW scalars. Hence we only need to solve the leading term in Eq. (C30). The normalized GW-scalar wave function to order ϵ is

$$\gamma_i = \frac{2\epsilon}{\mu_{(i)}} e^{-\tilde{k}r_c\varphi} \sqrt{\frac{\tilde{k}r_c\pi}{e^{2\pi\tilde{k}r_c} J_1\left(\frac{e^{\pi\tilde{k}r_c}\mu_{(i)}}{\tilde{k}r_c}\right)^2 - J_1\left(\frac{\mu_{(i)}}{\tilde{k}r_c}\right)^2}} J_1\left(\frac{e^{\tilde{k}r_c\varphi}\mu_{(i)}}{\tilde{k}r_c}\right). \quad (\text{C36})$$

Looking at the normalization condition of Eq. (15), we see that there is no ϵ^0 piece. Here we have used the large kr_c limit and omitted Bessel-Y functions. The masses of GW scalars are determined by solutions of the following transcendental equation:

$$J_2\left(\frac{e^{\tilde{k}r_c\varphi}\mu_{(i)}}{\tilde{k}r_c}\right) Y_2\left(\frac{\mu_{(i)}}{\tilde{k}r_c}\right) - J_2\left(\frac{\mu_{(i)}}{\tilde{k}r_c}\right) Y_2\left(\frac{e^{\pi\tilde{k}r_c}\mu_{(i)}}{\tilde{k}r_c}\right) = 0. \quad (\text{C37})$$

In the large kr_c limit, this reduces to the simple form

$$m_{(i)} = \frac{\mu_{(i)}}{r_c} = \tilde{k}e^{-\pi\tilde{k}r_c} j_{2,i}, \quad (\text{C38})$$

where $j_{2,i}$ are roots of J_2 .

-
- [1] T. Kaluza, Zum Unitätsproblem der physik, *Int. J. Mod. Phys. D* **27**, 1870001 (2018); *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1921**, 966 (1921).
 - [2] O. Klein, Quantum theory and five-dimensional theory of relativity. (In German and English), *Z. Phys.* **37**, 895 (1926).
 - [3] L. Randall and R. Sundrum, A Large Mass Hierarchy from a Small Extra Dimension, *Phys. Rev. Lett.* **83**, 3370 (1999).
 - [4] L. Randall and R. Sundrum, An Alternative to Compactification, *Phys. Rev. Lett.* **83**, 4690 (1999).
 - [5] M. Fierz and W. Pauli, On relativistic wave equations for particles of arbitrary spin in an electromagnetic field, *Proc. R. Soc. A* **173**, 211 (1939).
 - [6] N. Arkani-Hamed, H. Georgi, and M.D. Schwartz, Effective field theory for massive gravitons and gravity in theory space, *Ann. Phys. (Amsterdam)* **305**, 96 (2003).
 - [7] K. Hinterbichler, Theoretical aspects of massive gravity, *Rev. Mod. Phys.* **84**, 671 (2012).
 - [8] C. de Rham, Massive gravity, *Living Rev. Relativity* **17**, 7 (2014).
 - [9] R. Sekhar Chivukula, D. Foren, K. A. Mohan, D. Sengupta, and E. H. Simmons, Scattering amplitudes of massive spin-2 Kaluza-Klein states grow only as $\mathcal{O}(s)$, *Phys. Rev. D* **101**, 055013 (2020).
 - [10] R. Sekhar Chivukula, D. Foren, K. A. Mohan, D. Sengupta, and E. H. Simmons, Sum rules for massive spin-2 Kaluza-Klein elastic scattering amplitudes, *Phys. Rev. D* **100**, 115033 (2019).
 - [11] R. S. Chivukula, D. Foren, K. A. Mohan, D. Sengupta, and E. H. Simmons, Massive spin-2 scattering amplitudes

- in extra-dimensional theories, *Phys. Rev. D* **101**, 075013 (2020).
- [12] D. Foren, Scattering amplitudes in theories of compactified gravity, Ph.D. thesis, Michigan State University, 2020.
- [13] J. Bonifacio and K. Hinterbichler, Unitarization from geometry, *J. High Energy Phys.* **12** (2019) 165.
- [14] Y.-F. Hang and H.-J. He, Structure of Kaluza-Klein graviton scattering amplitudes from the gravitational equivalence theorem and double copy, *Phys. Rev. D* **105**, 084005 (2022).
- [15] R. Hofmann, P. Kanti, and M. Pospelov, (De)stabilization of an extra dimension due to a Casimir force, *Phys. Rev. D* **63**, 124020 (2001).
- [16] R. S. Chivukula, D. Foren, K. A. Mohan, D. Sengupta, and E. H. Simmons, Spin-2 Kaluza-Klein mode scattering in models with a massive radion, *Phys. Rev. D* **103**, 095024 (2021).
- [17] W. D. Goldberger and M. B. Wise, Modulus Stabilization with Bulk Fields, *Phys. Rev. Lett.* **83**, 4922 (1999).
- [18] W. D. Goldberger and M. B. Wise, Phenomenology of a stabilized modulus, *Phys. Lett. B* **475**, 275 (2000).
- [19] H. M. Lee, M. Park, and V. Sanz, Gravity-mediated (or Composite) dark matter, *Eur. Phys. J. C* **74**, 2715 (2014).
- [20] M. Garny, M. Sandora, and M. S. Sloth, Planckian Interacting Massive Particles as Dark Matter, *Phys. Rev. Lett.* **116**, 101302 (2016).
- [21] M. G. Folgado, A. Donini, and N. Rius, Gravity-mediated scalar dark matter in warped extra-dimensions, *J. High Energy Phys.* **01** (2020) 161; Erratum, *J. High Energy Phys.* **02** (2022) 129.
- [22] A. de Giorgi and S. Vogl, Dark matter interacting via a massive spin-2 mediator in warped extra-dimensions, *J. High Energy Phys.* **11** (2021) 036.
- [23] C. Csaki, M. L. Graesser, and G. D. Kribs, Radion dynamics and electroweak physics, *Phys. Rev. D* **63**, 065002 (2001).
- [24] L. Kofman, J. Martin, and M. Peloso, Exact identification of the radion and its coupling to the observable sector, *Phys. Rev. D* **70**, 085015 (2004).
- [25] E. E. Boos, Y. S. Mikhailov, M. N. Smolyakov, and I. P. Volobuev, Physical degrees of freedom in stabilized brane world models, *Mod. Phys. Lett. A* **21**, 1431 (2006).
- [26] E. E. Boos, V. E. Bunichev, I. P. Volobuev, and M. N. Smolyakov, Geometry, physics, and phenomenology of the Randall-Sundrum model, *Phys. Part. Nucl.* **43**, 42 (2012).
- [27] W. R. Inc., *Mathematica*, Version 13.0.0, [Champaign, IL], <https://www.wolfram.com/mathematica> (2021).
- [28] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevD.107.035015> for GitHub link to *Mathematica* [27] files giving expressions for all spin-2 and spin-0 perturbative wave functions.
- [29] J. W. York, Jr., Role of Conformal Three Geometry in the Dynamics of Gravitation, *Phys. Rev. Lett.* **28**, 1082 (1972).
- [30] G. W. Gibbons and S. W. Hawking, Action integrals and partition functions in quantum gravity, *Phys. Rev. D* **15**, 2752 (1977).
- [31] E. Dyer and K. Hinterbichler, Boundary terms, variational principles and higher derivative modified gravity, *Phys. Rev. D* **79**, 024028 (2009).
- [32] C. Charmousis, R. Gregory, and V. A. Rubakov, Wave function of the radion in a brane world, *Phys. Rev. D* **62**, 067505 (2000).
- [33] C. T. Fulton, Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions, *Proc. R. Soc. Edinb., Sect. A* **77**, 293 (1977).
- [34] P. Binding, P. J. Browne, and K. Seddighi, Sturm–Liouville problems with eigenparameter dependent boundary conditions, *Proc. Edinb. Math. Soc.* **37**, 57 (1994).
- [35] R. S. Chivukula, E. H. Simmons, and X. Wang, Supersymmetry and sum rules in the Goldberger-Wise model, *Phys. Rev. D* **106**, 035026 (2022).
- [36] O. DeWolfe, D. Z. Freedman, S. S. Gubser, and A. Karch, Modeling the fifth-dimension with scalars and gravity, *Phys. Rev. D* **62**, 046008 (2000).
- [37] I. Gradshteyn and I. Ryzhik, *Table of Integrals, Series, and Products* (Elsevier Science, New York, 2014).