

Investigation of Z -boson decay into Ξ_{bc} and Ξ_{bb} baryons within the NRQCD factorization approach*

Xuan Luo (罗炫) Hai-Bing Fu (付海冰)[†] Hai-Jiang Tian (田海江)

Department of Physics, Guizhou Minzu University, Guiyang 550025, China

Abstract: Z -boson decay provides a good opportunity to search for the Ξ_{bQ} baryon because a large number of its events can be collected at high energy colliders. In this paper, we perform a complete investigation of the indirect production of the Ξ_{bc} and Ξ_{bb} baryons via the Z -boson decay process $Z \rightarrow \Xi_{bQ} + \bar{b} + \bar{Q}$ with a $Q = (c, b)$ quark under the NRQCD factorization approach. After considering the contribution from the diquark states $\langle bc \rangle [^3S_1]_{3/6}$, $\langle bc \rangle [^1S_0]_{3/6}$, $\langle bb \rangle [^1S_0]_6$, and $\langle bb \rangle [^3S_1]_{3/6}$, the calculated branching fractions are $\mathcal{B}(Z \rightarrow \Xi_{bc} + X) = 3.595 \times 10^{-5}$ and $\mathcal{B}(Z \rightarrow \Xi_{bb} + X) = 1.213 \times 10^{-6}$. Furthermore, the Ξ_{bc} and Ξ_{bb} production events are predicted to be of the order of 10^4 and 10^3 at the LHC collider, and the order of 10^7 and 10^6 for the CEPC collider. We then estimate the production ratio for $Z \rightarrow c\bar{c}(b\bar{b}) \rightarrow \Xi_{bc}^{+,0} + X$ with $\Xi_{bc}^{+,0} \rightarrow \Lambda_c^+ + X$ in Z -boson decay, that is, $\mathcal{R}(Z \rightarrow c\bar{c} \rightarrow \Xi_{bc}^{+,0} + X) \sim \mathcal{O}(10^{-6})$ and $\mathcal{R}(Z \rightarrow b\bar{b} \rightarrow \Xi_{bb}^{+,0} + X) \sim \mathcal{O}(10^{-5})$, respectively. Finally, we present the differential decay widths of Ξ_{bc} and Ξ_{bb} with respect to s_{23} and z distributions and analyze the uncertainties.

Keywords: z -boson decay, nonrelativistic QCD, doubly heavy baryons

DOI: 10.1088/1674-1137/acbc0e

I. INTRODUCTION

Doubly heavy baryons composed by two heavy quarks and one light quark are expected by the quark model [1–4]. Investigating doubly heavy baryons is significant because it can provide a unique test for perturbative quantum chromodynamics (QCD) and nonrelativistic QCD (NRQCD). In the past few decades, research on doubly heavy baryons has developed rapidly from both the experimental and theoretical perspectives.

Experimentally, the doubly charmed baryon Ξ_{cc}^{++} was first observed by the LHCb collaboration based on the decay channel $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ (\rightarrow pK^-\pi^+) K^-\pi^+\pi^+$ [5]. It was subsequently identified via the measurement of $\Xi_{cc}^{++} \rightarrow \Xi_c^+ (\rightarrow pK^-\pi^+)\pi^+$ by the LHCb collaboration [6, 7]. Moreover, the first observation of the doubly charmed baryon Ξ_{cc}^+ was reported from $\Xi_{cc}^+ \rightarrow pD^+K^-$ by the SELEX collaboration. Over the past few years, the LHCb collaboration has published their observation of $\mathcal{R}(\Xi_{cc}^+)$, which is defined as $\mathcal{R}(\Xi_{cc}^+) = \sigma(\Xi_{cc}^+) \mathcal{B}(\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^-\pi^+) / \sigma(\Lambda_c^+)$, varying in the region $[0.9, 6.5] \times 10^{-3}$ for $\sqrt{s} = 8$ TeV, and $[0.12, 0.45] \times 10^{-3}$ for $\sqrt{s} = 13$ TeV [8]. These values are lower than the observations (9%) of the SELEX collabor-

ation.

Ξ_{bc} has also attracted the attention of researchers owing to its unique nature in the baryon family. In 2020, the LHCb collaboration searched for the doubly heavy baryon Ξ_{bc}^0 via its decay into the $D^0 p K^-$ final state, although no direct evidence was found [9]. Recently, Ξ_{bc}^0 and Ω_{bc}^0 were detected via the $\Lambda_c^+ \pi^-$ and $\Xi_c^+ \pi^-$ decay modes; however, evidence of the signal was not found [10]. Ξ_{bb} is yet to be detected. Overall, there is still no solid signal of the Ξ_{bQ} baryon with the heavy quark $Q = (c, b)$. To investigate the baryon production properties and further test the NRQCD method, considerable work has been conducted on both direct and indirect production [11–43].

Compared with direct production, such as hadroproduction, photoproduction, and e^+e^- annihilation, indirect production is also important because of the properties of baryons and their initial particles. Ξ_{bQ} baryons can be produced via different channels, such as top quark decay [44], W^+ -boson decay [45], and the $H \rightarrow \Xi_{bQ} + X$ process [46]. Other than the above channels, they can also be produced from Z -boson decays, for example, the process $Z \rightarrow \Xi_{cc} + X$ [47]. Ξ_{cc} events can reach 10^4 and 10^7 per year at the LHC and CEPC via Z -boson decays. Mean-

Received 4 October 2022; Accepted 14 February 2023; Published online 15 February 2023

* Supported in part by the Natural Science Foundation of China (12265010) and the Project of Guizhou Provincial Department of Education (KY[2021]030)

[†] E-mail: fuhb@cqu.edu.cn



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

while, the branching fraction $\mathcal{B}(Z \rightarrow \Xi_{cc} + X)$ is also comparable with $\mathcal{B}(Z \rightarrow J/\psi + X)$ [48, 49].

Thus, Z -boson decay can provide a good platform to study the Ξ_{bQ} baryon based on the large quantity of Z -boson events. Up to 10^9 and 10^{12} -order Z -boson events are produced per year at the LHC [50] and CEPC [51] colliders. Moreover, the decay channel $\Xi_{bc}^0 \rightarrow \Xi_{cc}^{++} + X$ has advantages over $\Xi_{bc}^0 \rightarrow \Lambda_c^+ \pi^-$, which offers a new experimental direction in the search for Ξ_{bc} [52]. In this paper, we first focus our attention on the indirect production of Ξ_{bQ} via Z -boson decay and then reveal whether a considerable amount of Ξ_{bQ} can be collected by Z -boson decay. In addition, we forecast $\mathcal{R}(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}^{+,0} + X)$ in Z -boson decay using the channel $\Xi_{bc}^{+,0} \rightarrow \Xi_{cc}^{++} + X$.

The rest of the paper is organized as follows: The detailed method is demonstrated in Sec. II, the phenomenological results and analyses are given in Sec. III, and a brief summary is given in Sec. IV.

II. CALCULATION TECHNOLOGY

Normally, production of the Ξ_{bQ} baryon can be treated in two steps [14, 37, 43, 53]. The first is by producing a bound state, which is also called a diquark $\langle bQ \rangle[n]$, with $[n]$ representing the color- and spin-combinations. Based on the decomposition $3 \otimes 3 = \bar{3} \oplus 6$ in the $SU_c(3)$ group and NRQCD, the quantum color number is only the color-antitriplet $\bar{3}$ and color-sextuplet 6 , and the quantum counts of the diquark $\langle bQ \rangle$ state are $[^3S_1]$ and $[^1S_0]$. The second step involves turning the diquark fragments into an observable baryon Ξ_{bQq} by hunting a light quark from the 'environment' with a fragmentation probability of almost one hundred percent. For convenience, we utilize the label Ξ_{bQ} instead of Ξ_{bQq} throughout this paper. Among this total "100%" fragmentation probability, the probability of both Ξ_{bQd} and Ξ_{bQu} is 43%, and the ratio for Ω_{bQs} is 14% [40, 54].

The diagrams for the process $Z(p_0) \rightarrow \langle bQ \rangle[n](p_1) + \bar{b}(p_2) + \bar{Q}(p_3)$ at tree level are shown in Fig. 1, where the heavy quark taken as $Q = c$ and b represents Ξ_{bc} and Ξ_{bb} , respectively. We can obtain the differential decay width of the process $Z(p_0) \rightarrow \langle bQ \rangle[n](p_1) + \bar{b}(p_2) + \bar{Q}(p_3)$ using the NRQCD factorization approach [55, 56],

$$d\Gamma = \sum_n d\hat{\Gamma}(Z \rightarrow \langle bQ \rangle[n] + \bar{b} + \bar{Q}) \langle O^H(n) \rangle. \quad (1)$$

Here, the long-distance matrix element $\langle O^H(n) \rangle$ describes the hadronization of the diquark state $\langle bQ \rangle[n]$ into the doubly heavy baryon Ξ_{bQ} . Generally, $\langle O^H(n) \rangle$ can be approximately obtained from the original value of the Schrödinger wave function or its derivative. In this paper, we take $\langle O^H(n) \rangle = (|\Psi_{bQ}(0)\rangle, |\Psi'_{bQ}(0)\rangle)$ for the S -wave and P -wave, which are derived from experimental data and non-perturbative theoretical methods, for example, the potential model, lattice QCD, and QCD sum rules [56–58].

The differential decay width $d\hat{\Gamma}(Z \rightarrow \langle bQ \rangle[n] + \bar{b} + \bar{Q})$ can be written as

$$d\hat{\Gamma}(Z \rightarrow \langle bQ \rangle[n] + \bar{b} + \bar{Q}) = \frac{1}{3} \frac{1}{2m_z} \sum |M[n]|^2 d\Phi_3, \quad (2)$$

where m_z and $|M[n]|$ are the Z -boson mass and hard amplitude, respectively. The constant $1/3$ arises from the spin average of the initial Z -boson, and the symbol " \sum " represents the sum of color and spin for all final particles. The three-body phase space $d\Phi_3$ with a massive quark or antiquark in the final state can be written as

$$d\Phi_3 = (2\pi)^4 \delta^4\left(p_0 - \sum_f p_f\right) \prod_f \frac{d^3 p_f}{(2\pi)^3 2p_f^0}. \quad (3)$$

The calculation for the three-body phase space has been discussed in Refs. [59, 60]. Then, Eq. (2) can be rewritten as

$$d\hat{\Gamma}(Z \rightarrow \langle bQ \rangle[n] + \bar{b} + \bar{Q}) = \frac{1}{2^8 \pi^3 m_z^3} \sum |M[n]|^2 ds_{12} ds_{23}, \quad (4)$$

with $s_{ij} = (p_i + p_j)^2$. After using charge parity $C = -i\gamma^2\gamma^0$, hard amplitude expressions $M[n]$ for baryon production are obtained, which can also be easily gained from the familiar meson production [26, 39]. Here, we give brief descriptions. $C = -i\gamma^2\gamma^0$ can be used to reverse




Fig. 1. (color online) Diagrams for $Z \rightarrow \langle bQ \rangle[n] + \bar{b} + \bar{Q}$ at the leading order, where the heavy quark $Q = c$ and b represents Ξ_{bc} and Ξ_{bb} , respectively.

one fermion line, which can be written as $L_1 = \bar{u}_{s_1}(k_1)\Gamma_{i+1}S_F(q_i, m_i)\cdots S_F(q_1, m_1)\Gamma_1 v_{s_2}(k_2)$, where Γ_i , $S_F(q_i, m_i)$, $s_{1,2}$, and $i = (0, 1, \dots)$ are the interaction vertex, fermion propagator, spin index, and quantity of interaction vertices in the fermion line, respectively. According to charge parity, we have

$$\begin{aligned} v_{s_2}^T(p)C &= -\bar{u}_{s_2}(p), & C^{-1}\Gamma_i^T C &= -\Gamma_i, \\ CC^{-1} &= I, & C^{-1}S_f^T(-q_i, m_i)C &= S_f(q_i, m_i), \\ C^{-1}\bar{u}_{s_1}^T(p_{12}) &= v_{s_1}(p_{12}), & C^{-1}(\gamma^\mu)^T C &= -\gamma^\mu, \\ C^{-1}(\gamma^\mu\gamma^5)^T C &= \gamma^\mu\gamma^5. \end{aligned} \quad (5)$$

If the fermion line does not include an axial vector vertex, we can readily obtain the expression

$$\begin{aligned} L_1 &= L_1^T = v_{s_2}^T(p_2)\Gamma_1^T S_F^T(q_1, m_1)\cdots S_F^T(q_1, m_1)\Gamma_{i+1}^T \bar{u}_{s_1}^T(p_{12}) \\ &= v_{s_2}^T(p_2)CC^{-1}\Gamma_1^T CC^{-1}S_F^T(q_1, m_1)CC^{-1} \\ &\quad \cdots CC^{-1}S_F^T(q_1, m_1)CC^{-1}\Gamma_{i+1}^T CC^{-1}\bar{u}_{s_1}^T(p_{12}) \end{aligned}$$

$$\begin{aligned} &= (-1)^{i+1}\bar{u}_{s_2}(p_2)\Gamma_1 S_F(-q_1, m_1) \\ &\quad \cdots S_F(-q_i, m_i)\Gamma_{i+1} v_{s_1}(p_{12}). \end{aligned} \quad (6)$$

Otherwise, the amplitudes of baryon production can be obtained from familiar meson production, except with an additional $(-1)^{(n+1)}$ coefficient for the pure vector case and a $(-1)^{(n+2)}$ factor when including an axial vector case. In other words, the amplitude of $Z \rightarrow \langle bQ \rangle[n] + \bar{b} + \bar{Q}$ can be written as

$$M_{\text{diquark}} = (M_1^a - M_1^v) + (M_2^a - M_2^v) + M_3 + M_4, \quad (7)$$

where M_i ($i = 1, 2, 3, 4$) is the hard amplitude of familiar meson production, and M_i^a and M_i^v are the components of the axial vector amplitudes and pure vector amplitudes of M_i , respectively.

Taking the traditional Feynman rules of Fig. 1 into consideration, we can obtain the amplitudes $M_l[n]$ with $l = (a, b, c, d)$, which have the following expressions:

$$\begin{aligned} M_a[n] &= -\kappa \frac{\bar{u}(p_{12})(-i\gamma^\nu)v(p_2)\bar{u}(p_{11})(-i\gamma^\nu)(m_Q + \not{p}_1 + \not{p}_2)\not{\epsilon}(p_0)(c_v^Q + c_a^Q\gamma^5)v(p_3)}{(p_{12} + p_2)^2[(p_1 + p_2)^2 - m_Q^2]}, \\ M_b[n] &= -\kappa \frac{\bar{u}(p_{12})(-i\gamma^\nu)(m_b + \not{p}_1 + \not{p}_3)\not{\epsilon}(p_0)(c_v^Q + c_a^Q\gamma^5)v(p_2)\bar{u}(p_{11})(-i\gamma^\nu)v(p_3)}{(p_{11} + p_3)^2[(p_1 + p_3)^2 - m_b^2]}, \\ M_c[n] &= -\kappa \frac{\bar{u}(p_{12})\not{\epsilon}(p_0)(c_v^Q + c_a^Q\gamma^5)(m_b - \not{p}_{11} - \not{p}_2 - \not{p}_3)(-i\gamma^\nu)v(p_2)\bar{u}(p_{11})(-i\gamma^\nu)v(p_3)}{(p_{11} + p_3)^2[(p_{11} + p_2 + p_3)^2 - m_b^2]}, \\ M_d[n] &= -\kappa \frac{\bar{u}(p_{12})(-i\gamma^\nu)v(p_2)\bar{u}(p_{11})\not{\epsilon}(p_0)(c_v^Q + c_a^Q\gamma^5)(m_Q - \not{p}_{12} - \not{p}_2 - \not{p}_3)(-i\gamma^\nu)v(p_3)}{(p_{12} + p_2)^2[(p_{12} + p_2 + p_3)^2 - m_Q^2]}. \end{aligned} \quad (8)$$

Here, $\kappa = -Cg_s^2$, with color factor C_{ijk} . p_{11} and p_{12} are the momenta of the bottom quark and another heavy quark $Q = (c, b)$ for Ξ_{bc} and Ξ_{bb} production. The vector and axial vector coupling constants of the $Z_{Q\bar{Q}}$ vertex, that is, c_v^Q and c_a^Q , have the following expressions:

$$\begin{aligned} c_v^c &= -\frac{e(8\sin^2\theta_w - 3)}{12\cos\theta_w\sin\theta_w}, & c_a^c &= -\frac{e}{4\cos\theta_w\sin\theta_w}, \\ c_v^b &= \frac{e(4\sin^2\theta_w - 3)}{12\cos\theta_w\sin\theta_w}, & c_a^b &= \frac{e}{4\cos\theta_w\sin\theta_w}. \end{aligned} \quad (9)$$

Here, θ_w is the Weinberg angle. With the help of Eq. (6) and inserting the spin projector $\Pi_{p_i}^{[n]}$, the amplitude can be re-written as

$$\begin{aligned}
M_a[n] &= -\kappa \frac{\bar{u}(p_2)(-i\gamma^\nu)\Pi_{p_1}^{[n]}(-i\gamma^\nu)(m_Q + \not{p}_1 + \not{p}_2)\not{e}(p_0)(c_v^Q + c_a^Q\gamma^5)v(p_3)}{(p_{12} + p_2)^2[(p_1 + p_2)^2 - m_Q^2]}, \\
M_b[n] &= -\kappa \frac{\bar{u}(p_2)\not{e}(p_0)(c_a^Q\gamma^5 - c_v^Q)(m_b - \not{p}_1 - \not{p}_3)(-i\gamma^\nu)\Pi_{p_1}^{[n]}(-i\gamma^\nu)v(p_3)}{(p_{11} + p_3)^2((p_1 + p_3)^2 - m_b^2)}, \\
M_c[n] &= -\kappa \frac{\bar{u}(p_2)(-i\gamma^\nu)(m_b + \not{p}_{11} + \not{p}_2 + \not{p}_3)\not{e}(p_0)(c_a^Q\gamma^5 - c_v^Q)\Pi_{p_1}^{[n]}(-i\gamma^\nu)v(p_3)}{(p_{11} + p_3)^2[(p_{11} + p_2 + p_3)^2 - m_b^2]}, \\
M_d[n] &= -\kappa \frac{\bar{u}(p_2)(-i\gamma^\nu)\Pi_{p_1}^{[n]}\not{e}(p_0)(c_v^Q + c_a^Q\gamma^5)(m_Q - \not{p}_{12} - \not{p}_2 - \not{p}_3)(-i\gamma^\nu)v(p_3)}{(p_{12} + p_2)^2[(p_{12} + p_2 + p_3)^2 - m_Q^2]}. \quad (10)
\end{aligned}$$

Here, the spin projector $\Pi_{p_1}^{[n]}$ has the following form [61]:

$$\begin{aligned}
\Pi_{p_1}^{[S_0]} &= \frac{1}{2\sqrt{M_{bQ}}}\gamma^5(\not{p}_1 + M_{bQ}), \\
\Pi_{p_1}^{[S_{\perp}]} &= \frac{1}{2\sqrt{M_{bQ}}}\not{e}(\not{p}_1 + M_{bQ}), \quad (11)
\end{aligned}$$

where $M_{bQ} \simeq m_b + m_Q$ is used to maintain gauge invariance.

Furthermore, the color factor $C_{i,j,k}$ can be easily obtained from Fig. 1, which has the following form:

$$C_{i,j,k} = N \times \sum_{a,m,n} (T^a)_{im}(T^a)_{jn} \times G_{mnk}. \quad (12)$$

Here, $k, a = (1, 2, \dots, 8)$, $N = \sqrt{1/2}$, and $i, j, m, n = (1, 2, 3)$ represent the diquark color indices, gluon color indices, normalization factor, and two outgoing antiquarks and two constituent quarks in the diquark color indices, respectively. For the $\bar{\mathbf{3}}(\mathbf{6})$ state, the function G_{mnk} is identical to the antisymmetric function ε_{mjk} and symmetric function f_{mjk} , which obey

$$\begin{aligned}
\varepsilon_{mjk}\varepsilon_{m'j'k} &= \delta_{mm'}\delta_{jj'} - \delta_{mj'}\delta_{jm'}, \\
f_{mjk}f_{m'j'k} &= \delta_{mm'}\delta_{jj'} + \delta_{mj'}\delta_{jm'}. \quad (13)
\end{aligned}$$

For color $\bar{\mathbf{3}}$ and $\mathbf{6}$ diquark state production, $C_{ijk}^2 = 4/3$ and $C_{ijk}^2 = 2/3$, respectively.

Meanwhile, diquark hadronization into a doubly heavy baryon is a non-perturbative procedure, which is factorized into a general coefficient $\langle O^H(n) \rangle$. This coefficient is connected to the wave function at the origin. In this paper, we take the usual assumption that the wave

function of the color $\bar{\mathbf{3}}$ state is equal to that of the color $\mathbf{6}$ state, as discussed in Refs. [26, 39, 44, 45, 53]¹⁾.

The transition probabilities of the color $\bar{\mathbf{3}}$ and $\mathbf{6}$ states are represented by $h_{\bar{\mathbf{3}}}$ and $h_{\mathbf{6}}$, respectively. Based on the NRQCD approach, a bound state of two heavy quarks with another light dynamical freedom of QCD Ξ_{bQ} can be described by a series of Fock states,

$$\begin{aligned}
|\Xi_{bQ}\rangle &= c_1(v)|(bQ)q\rangle + c_2(v)|(bQ)qg\rangle \\
&\quad + c_3(v)|(bQ)qgg\rangle + \dots, \quad (14)
\end{aligned}$$

where v denotes the small relative velocity between heavy quarks in the rest frame of the diquark. For the color $\bar{\mathbf{3}}$ state cases, one of the heavy quarks of the diquark can produce a gluon without altering its spin, which can divide into a light quark pair $q\bar{q}$. Then, the diquark can capture a light quark q to construct a baryon. Regarding the color $\mathbf{6}$ state, if the baryon is created by $|(bQ)q\rangle$, the emitted gluon would alter the spin of the heavy quark, causing suppression of $h_{\mathbf{6}}$. If the baryon is created from the $|(bQ)qg\rangle$ parts, one of the heavy quarks produces a gluon without altering the spin of the heavy quark. Then, the gluon separates into $q\bar{q}$. Additionally, a light quark q has the ability to produce gluons, which can be used to construct the component with qg . These contributions are at the same level because a light quark may produce gluons easily, that is, $c_1(v) \sim c_2(v) \sim c_3(v)$ [53]. We can then take the following approximation:

$$h_{\mathbf{6}} \sim h_{\bar{\mathbf{3}}} = \langle O^H(n) \rangle = |\Psi(0)|^2, \quad (15)$$

for the S -wave, and for P -wave,

$$h_{\mathbf{6}} \sim h_{\bar{\mathbf{3}}} = \langle O^H(n) \rangle = |\Psi'(0)|^2. \quad (16)$$

1) As reported in Ref. [39], the color $\mathbf{6}$ state is suppressed to the color $\bar{\mathbf{3}}$ state by order v^2 and its contribution can be disregarded, which can be ascribed to the one-gluon exchange interaction causes the interaction inside the diquark with the color $\mathbf{6}$ state to be repulsive rather than attractive. Meanwhile, as pointed out in Refs. [26, 44, 45, 53], the importance of the color $\mathbf{6}$ and the color $\bar{\mathbf{3}}$ states are equal.

III. NUMERICAL RESULTS

To perform the numerical calculation, the following input parameters are taken: The c , b -quark masses are $m_c = 1.8$ GeV and $m_b = 5.1$ GeV, respectively. The Z-boson mass $m_Z = 91.1876$ GeV and the decay width $\Gamma_Z = 2.4952$ GeV are from the PDG [62]. For the values of $|\Psi_{bc}(0)|^2$ and $|\Psi_{bb}(0)|^2$, we adopt 0.065 GeV^3 and 0.152 GeV^3 [15], respectively. The masses of the Ξ_{bc} and Ξ_{bb} baryons are taken as $m_{\Xi_{bc}} = 6.9$ GeV and $m_{\Xi_{bb}} = 10.2$ GeV, respectively. The remaining input parameters are [62] $G_F = 1.1663787 \times 10^{-5}$, which denotes the Fermi constant, and the Weinberg angle $\theta_w = \arcsin \sqrt{0.2312}$. The renormalization scale μ_r is taken as $2m_c(2m_b)$ for the indirect production of $\Xi_{bc}(\Xi_{bb})$.

First, the decay widths of two main Z-boson decay channels for Ξ_{bQ} production are given in Table 1. From Table 1, we can see that the $[^3S_1]_3$ state plays a leading role in the production of Ξ_{bb} . The contribution from the $[^3S_1]_3$ state can reach twice that from the $[^1S_0]_6$ state. As for Ξ_{bc} production, the situation is analogous to that of Ξ_{bb} . Moreover, in the case of Ξ_{bc} , the contribution from $Z \rightarrow c\bar{c}$ is significantly smaller than that from $Z \rightarrow b\bar{b}$, which is only a few percent.

To assess the doubly heavy baryon Ξ_{bQ} events generated at the LHC (CEPC), the corresponding branching ratio must be obtained from the Z-boson total decay width. At the LHC (CEPC), approximately $10^9(10^{12})$ Z-bosons can be produced per year [51, 63]. Based on the conditions mentioned above, the produced events of the double heavy baryon $\Xi_{bc}(\Xi_{bb})$ can be predicted at the LHC (CEPC). We list the total decay width, branching ratios, and events of the Ξ_{bc} and Ξ_{bb} baryons via Z-boson decay in Table 2, where contributions from each diquark state of the Z-boson decay channel are considered for the total

Table 1. Predicted decay widths $\Gamma(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}(\Xi_{bb}) + X)$ (unit: 10^{-6} GeV) with $Q = (c, b)$ for the Ξ_{bc} and Ξ_{bb} baryons from each Z-boson decay channel.

$\Gamma(Z \rightarrow Q\bar{Q})$	Ξ_{bc}				Ξ_{bb}	
	$[^3S_1]_3$	$[^3S_1]_6$	$[^1S_0]_3$	$[^1S_0]_6$	$[^3S_1]_3$	$[^1S_0]_6$
$Z \rightarrow c\bar{c}$	0.644	0.322	0.741	0.371	-	-
$Z \rightarrow b\bar{b}$	33.01	16.51	24.14	12.07	1.999	1.028

Table 2. Predicted total decay width (unit: GeV), branching fraction, and events of the Ξ_{bc} and Ξ_{bb} baryons in Z-boson decay.

	$Z \rightarrow \Xi_{bc}$	$Z \rightarrow \Xi_{bb}$
$\Gamma(Z \rightarrow \Xi_{bQ})$	89.71×10^{-6}	3.027×10^{-6}
$\mathcal{B}(Z \rightarrow \Xi_{bQ})$	35.95×10^{-6}	1.213×10^{-6}
LHC events	35.95×10^3	1.213×10^3
CEPC events	35.95×10^6	1.213×10^6

decay width. From Table 2, we can reach the following conclusions:

- For the production of Ξ_{bb} , the branching ratio $\mathcal{B}(Z \rightarrow \Xi_{bb} + X)$ is approximately 10^{-6} , which is comparable to the results given in Ref. [64].
- The branching ratio of $\mathcal{B}(Z \rightarrow \Xi_{bc} + X)$ reaches 10^{-5} for the production of Ξ_{bc} , which is also comparable to the predictions of $\mathcal{B}(Z \rightarrow B_c + X)$ [65].
- At the CEPC, there are approximately 10^7 -order Ξ_{bc} events and 10^6 -order Ξ_{bb} events obtained per year.
- In comparison, there are only approximately 10^4 order Ξ_{bc} events and 10^3 order Ξ_{bb} events produced at the LHC. However, the upgraded program of the HE(L)-LHC will significantly improve Z-boson yield events; thus, more Ξ_{bc} and Ξ_{bb} events will be produced.
- Considering the decay rate of the channels $\Xi_{bc}^+ \rightarrow \Xi_{cc}^{++} + X \simeq 7\%$ [52], $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+ \simeq 10\%$ [66], and $\Lambda_c^+ \rightarrow p K^+ \pi^+ \simeq 5\%$ [67], approximately 10^4 reconstructed Ξ_{bc}^+ events can be collected at the CEPC. These events are comparable to $\Xi_{cc}^{++(+)}$ [47], which proves the observability of Ξ_{bc}^+ via Z-boson decay.

Furthermore, the ratio of the $\Xi_{bc}^{+,0}$ production rate $\mathcal{R}(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}^{+,0} + X)$, which arises from Z-boson decay to Λ_c^+ accompanied by $K^- \pi^+ \pi^+$, has the following formula:

$$\mathcal{R}(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}^{+,0} + X) = \frac{\Gamma(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}^{+,0} + X)}{\Gamma(Z \rightarrow Q\bar{Q} \rightarrow \Lambda_c^+)} \times \mathcal{B}(\Xi_{bc}^{+,0} \rightarrow \Xi_{cc}^{++} + X) \mathcal{B}(\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+), \quad (17)$$

where X denotes all possible particles. First, we use the formula $\mathcal{B}(Z \rightarrow Q\bar{Q} \rightarrow \Lambda_c^+) = \mathcal{B}(Z \rightarrow Q\bar{Q}) \times f(Q \rightarrow \Lambda_c^+)$ to obtain $\Gamma(Z \rightarrow Q\bar{Q} \rightarrow \Lambda_c^+)$. This is based on the total decay width, which can be directly related to the branching fraction. The branching fractions of $Z \rightarrow Q\bar{Q}$ are taken from the PDG, that is, $\mathcal{B}(Z \rightarrow c\bar{c}) = 0.12$ and $\mathcal{B}(Z \rightarrow b\bar{b}) = 0.15$ [68]. The fragmentation fractions of a heavy quark to a particular charmed hadron are $f(c \rightarrow \Lambda_c^+) = 0.057$ and $f(b \rightarrow \Lambda_c^+) = 0.073$ [69]. Then, we have

$$\begin{aligned} \mathcal{B}(Z \rightarrow c\bar{c} \rightarrow \Lambda_c^+) &= 6.84 \times 10^{-3}, \\ \mathcal{B}(Z \rightarrow b\bar{b} \rightarrow \Lambda_c^+) &= 10.95 \times 10^{-3}. \end{aligned} \quad (18)$$

Second, according to the decay chains of $\Xi_{bc}^{+,0} \rightarrow \Xi_{cc}^{++} + X \simeq 7\%(1.5\%)$ [52], $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+ \simeq 10\%$ [66], and $\Lambda_c^+ \rightarrow p K^+ \pi^+ \simeq 5\%$ [67], we can get the final results




Fig. 2. (color online) Predictions for $\mathcal{R}(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}^{+,0} + X)$ with four different Z-boson decay channels, where $Q = (c, b)$. Here, the renormalization scale is set near $\mu_r = 2m_c$.

shown in Fig. 2. Here, the renormalization scale μ_r is set to $2m_c$. $\mathcal{R}(Z \rightarrow b\bar{b} \rightarrow \Xi_{bc}^{+,0} + X)$ is one magnitude larger than $\mathcal{R}(Z \rightarrow c\bar{c} \rightarrow \Xi_{bc}^{+,0} + X)$, which indicates that the decay channel $Z \rightarrow b\bar{b}$ provides key contributions compared with the $Z \rightarrow c\bar{c}$ channel for indirect $\Xi_{bc}^{+,0}$ production. Comparing the predictions of Ξ_{bc} in this study with Ξ_{cc} from our previous work [47] for Z-boson decay, there is a large gap between $\mathcal{R}(Z \rightarrow c\bar{c} \rightarrow \Xi_{cc}^{+,++} + X)$ and $\mathcal{R}(Z \rightarrow b\bar{b} \rightarrow \Xi_{bc}^{+,0} + X)$ of approximately one magnitude. This discrepancy indicates that it will be difficult to collect $\Xi_{bc}^{+,0}$ in experiment collaborations. Moreover, our predictions for $\mathcal{R}(Z \rightarrow Q\bar{Q} \rightarrow \Xi_{bc}^{+,0} + X)$ via the $\Xi_{cc}^{+,++}$ channel reach the order of 10^{-5} in Z-boson decay, which is larger than those for the $\Lambda_c^+\pi^-$ channel (of the 10^{-6} order) [70]. Thus, the observation of $\Xi_{bc}^{+,0}$ via the $\Xi_{cc}^{+,++} + X$ chan-

nel is more feasible than that via the $\Xi_{bc}^0 \rightarrow \Lambda_c^+\pi^-$ channel.

To further study the production of Ξ_{bcQ} via these considered decay channels and provide a reference for experimental research, we present the differential decay widths of Ξ_{bcQ} with respect to the invariant mass s_{23} and energy fraction z in Figs. 3 and 4, where $s_{ij} = (p_i + p_j)^2$, and $z = 2E_1/E_Z$, using the Ξ_{bcQ} energy E_1 and Z-boson energy E_Z .

- In Fig. 3, we find that the $[^3S_1]$ state plays the leading role in the cases of Ξ_{bc} and Ξ_{bb} production. The curves $d\Gamma/ds_{23}$ have a similar behavior, which increase initially and then decrease with s_{23} , with the peak located in the small region of s_{23} .

- As shown in Fig. 4, the behavior of the differential decay widths changes with the energy fraction z -distribution, that is, $d\Gamma/dz$ is similar to $d\Gamma/ds_{23}$, which increases initially and then decreases. In the case of Ξ_{bb} production, the peak of $d\Gamma/dz|_{(bb)[^3S_1]_{\bar{3}}}$ is approximately $z = 0.75$, and $d\Gamma/dz|_{(bb)[^1S_0]_{\bar{6}}}$ peaks near $z = 0.7$. As for Ξ_{bc} , the peak of $d\Gamma/dz|_{(bc)[^3S_1]_{\bar{3}(6)}}$ is approximately $z = 0.8$, and $d\Gamma/dz|_{(bc)[^1S_0]_{\bar{3}(6)}}$ peaks near $z = 0.85$. Owing to the dominant effect of the quark fragmentation mechanism, the peaks of the differential decay widths for $Z \rightarrow \Xi_{bc(bb)} + X$ with energy distribution are located in the larger z -region.

Finally, to discuss the theoretical uncertainties for the process $Z \rightarrow \Xi_{bcQ} + X$ precisely, c, b -quark masses of $m_c = 1.80 \pm 0.5$ GeV and $m_b = 5.1 \pm 0.5$ GeV, and the renormalization scale $\mu_r = 2m_c(m_z/2)$ for Ξ_{bc} and $\mu_r = 2m_b(m_z/2)$ for Ξ_{bb} can be considered. Here, the uncertainties from $|\Psi_{bQ}(0)|^2$ are not discussed; they are an overall coefficient in calculations and can be computed out easily. The total decay widths within uncertainties arising from the above input parameters are presented in Table 3, which shows that




Fig. 3. (color online) Invariant mass differential decay widths $d\Gamma/ds_{23}$ for the process $Z \rightarrow \Xi_{bc}(\Xi_{bb}) + X$ under the NRQCD factorization approach, where $\bar{3}(6)$ indicates that the color quantum number is $\bar{3}(6)$ of the diquark state, and "Total" denotes the total decay widths, which means that each diquark state has been summed.




Fig. 4. (color online) Differential decay widths $d\Gamma/dz$ for the process $Z \rightarrow \Xi_{bc}(\Xi_{bb}) + X$ under the NRQCD factorization approach, where $\bar{3}(6)$ indicates that the color quantum number is $\bar{3}(6)$ of the diquark state, and "Total" denotes the total decay widths, which means that each diquark state has been summed.

Table 3. Decay width Γ (unit: 10^{-6}) of the process $Z \rightarrow \Xi_{bc}(\Xi_{bb}) + X$ within theoretical uncertainties through changing the mass of the charm quark $m_c = 1.80 \pm 0.5$ GeV and the mass of the bottom quark $m_b = 5.10 \pm 0.5$ GeV.

μ_r	m_Q	Ξ_{bc}				Ξ_{bb}	
		$[^3S_1]_{\bar{3}}$	$[^3S_1]_6$	$[^1S_0]_{\bar{3}}$	$[^1S_0]_6$	$[^3S_1]_{\bar{3}}$	$[^1S_0]_6$
$2m_c$	$m_c = 1.30\text{GeV}$	100.4	50.19	67.07	33.53	1.999	1.028
	$m_c = 1.80\text{GeV}$	34.40	17.20	25.41	12.71	1.999	1.028
	$m_c = 2.30\text{GeV}$	15.64	7.820	12.46	6.231	1.999	1.028
$m_Z/2$	$m_c = 1.30\text{GeV}$	27.88	13.94	18.62	9.312	1.053	0.542
	$m_c = 1.80\text{GeV}$	9.552	4.776	7.057	3.528	1.053	0.542
	$m_c = 2.30\text{GeV}$	4.343	2.172	3.460	1.730	1.053	0.542
$2m_c$	$m_b = 4.60\text{GeV}$	33.89	16.95	25.91	12.96	3.044	1.385
	$m_b = 5.10\text{GeV}$	34.40	17.20	25.41	12.71	1.999	1.028
	$m_b = 5.60\text{GeV}$	34.92	17.46	25.00	12.50	1.345	0.697
$m_Z/2$	$m_b = 4.60\text{GeV}$	9.412	4.706	7.196	3.598	1.515	0.773
	$m_b = 5.10\text{GeV}$	9.552	4.776	7.057	3.528	1.053	0.542
	$m_b = 5.60\text{GeV}$	9.696	4.848	6.943	3.471	0.750	0.389

• For indirect Ξ_{bc} production in Z -boson decay, the decay width decreases as the c -quark mass increases, which is mainly ascribed to the suppression of phase space.

• Owing to the effect of the projector in Eq. (11), an abnormal phenomenon occurs in which the decay width increases as the b -quark mass increases for the $\langle bc \rangle [^3S_1]$ state in indirect production via Z -boson decay. The $Z \rightarrow \langle bc \rangle [^1S_0] + X$ decay width decreases as the b -quark mass increases. Moreover, the uncertainty caused by m_c is larger than that of m_b .

• For the process $Z \rightarrow \Xi_{bb} + X$ with the $[^3S_1]_3$ and $[^1S_0]_6$ cases, the decay width decreases as the b -quark mass increases.

IV. SUMMARY

In this study, we discuss in detail the indirect production of Ξ_{bc} and Ξ_{bb} via Z -boson decay based on the framework of NRQCD. After considering the contributions from the intermediate diquark states, that is, $\langle bc \rangle [^3S_1]_{3/6}$, $\langle bc \rangle [^1S_0]_{3/6}$, $\langle bb \rangle [^1S_0]_6$, and $\langle bb \rangle [^3S_1]_3$, the branching ratio $\mathcal{B}(Z \rightarrow \Xi_{bc} + X)$ is approximately of the order of 10^{-5} , and $\mathcal{B}(Z \rightarrow \Xi_{bb} + X)$ is of the 10^{-6} order. There will be $10^4(10^7)\Xi_{bc}$ events and $10^3(10^6)\Xi_{bb}$ events produced at the LHC (CEPC). Then, the change in the differential decay widths of Ξ_{bc} and Ξ_{bb} with s_{23} and z

distributions is presented. Moreover, we estimate the production ratio $\mathcal{R}(Z \rightarrow \Xi_{bc}^{+,0} + X)$ of $\Xi_{bc}^{+,0}$ to Λ_c^+ via the Z -boson decay channel $c\bar{c}$ and $b\bar{b}$ for the first time, resulting in values of up to 10^{-6} for the $c\bar{c}$ channel and 10^{-5} for the $b\bar{b}$ channel. Abundant Ξ_{bQ} baryon events and the considerable branching ratio $\mathcal{B}(Z \rightarrow \Xi_{bQ} + X)$ demonstrate the observability of the Ξ_{bQ} baryon in Z -boson decay during the experiment. Thus, we believe that it is worthwhile and feasible to search for the Ξ_{bQ} baryon through Z -boson decay at the LHC and CEPC.

In conclusion, at present, studies on the decay properties of doubly heavy baryons are discussed via their decay models. Especially in LHCb collaboration reports [71, 72] and theoretical results from Qin [52], decay models and their observational possibilities have been discussed. Inspired by the observation of the doubly-charmed baryon Ξ_{cc}^{++} , the Ξ_{bc} baryon may be detected by the decay channels $\Xi_{bc} \rightarrow \Xi_{cc}^{++}(\rightarrow pK^-\pi^+\pi^+) + X$, where X represents all possible particles. The advantage of this approach in detecting Ξ_{bc} is that the detection efficiency will be greatly improved because only Ξ_{cc}^{++} needs to be reconstructed, as discussed in Ref. [52]. Similar to the Ξ_{bc} baryon, Ξ_{bb} may also be observed via $\Xi_{bb} \rightarrow \Xi_{bc}(\rightarrow \Xi_{cc}) + X$.

ACKNOWLEDGMENTS

We are grateful for Professor Zhan Sun's valuable comments and suggestions.

References

- [1] M. Gell-Mann, *Phys. Lett.* **8**, 214 (1964)
- [2] D. Ebert, R. N. Faustov, V. O. Galkin *et al.*, *Z. Phys. C* **76**, 111 (1997), arXiv:hep-ph/9607314
- [3] S. M. Gerasyuta and D. V. Ivanov, *Nuovo Cim. A* **112**, 261 (1999), arXiv:hep-ph/0101310
- [4] C. Itoh, T. Minamikawa, K. Miura *et al.*, *Phys. Rev. D* **61**, 057502 (2000)
- [5] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **119**, 112001 (2017), arXiv:1707.01621
- [6] R. Aaij *et al.* (LHCb Collaboration), *Chin. Phys. C* **44**, 022001 (2020), arXiv:1910.11316
- [7] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **121**, 162002 (2018), arXiv:1807.01919
- [8] R. Aaij *et al.* (LHCb Collaboration), *Sci. China Phys. Mech. Astron.* **63**, 221062 (2020), arXiv:1909.12273
- [9] R. Aaij *et al.* (LHCb Collaboration), *JHEP* **11**, 095 (2020), arXiv:2009.02481
- [10] R. Aaij *et al.* (LHCb Collaboration), *Chin. Phys. C* **45**, 093002 (2021), arXiv:2104.04759
- [11] S. J. Brodsky, S. Groote, and S. Koshkarev, *Eur. Phys. J. C* **78**, 483 (2018), arXiv:1709.09903
- [12] V. V. Kiselev, A. K. Likhoded, and M. V. Shevlyagin, *Phys. Lett. B* **332**, 411 (1994), arXiv:hep-ph/9408407
- [13] A. F. Falk, M. E. Luke, M. J. Savage *et al.*, *Phys. Rev. D* **49**, 555 (1994), arXiv:hep-ph/9305315
- [14] C. H. Chang, J. P. Ma, C. F. Qiao *et al.*, *J. Phys. G* **34**, 845 (2007), arXiv:hep-ph/0610205
- [15] S. P. Baranov, *Phys. Rev. D* **54**, 3228 (1996)
- [16] G. T. Bodwin, E. Braaten, and G. P. Lepage, *Phys. Rev. D* **51**, 1125 (1995), arXiv:hep-ph/9407339
- [17] D. A. Gunter and V. A. Saleev, *Phys. Rev. D* **64**, 034006 (2001), arXiv:hep-ph/0104173
- [18] V. V. Kiselev, A. K. Likhoded, and M. V. Shevlyagin, *Phys. Atom. Nucl.* **58**, 1018 (1995)
- [19] A. V. Berezhnoy and A. K. Likhoded, *Phys. Atom. Nucl.* **70**, 478 (2007), arXiv:hep-ph/0602041
- [20] V. V. Braguta, V. V. Kiselev, and A. E. Chalov, *Phys. Atom. Nucl.* **65**, 1537 (2002)
- [21] E. Braaten, M. Kusunoki, Y. Jia *et al.*, *Phys. Rev. D* **70**, 054021 (2004), arXiv:hep-ph/0304280
- [22] S. Y. Li, Z. G. Si, and Z. J. Yang, *Phys. Lett. B* **648**, 284 (2007), arXiv:hep-ph/0701212
- [23] Z. J. Yang and T. Yao, *Chin. Phys. Lett.* **24**, 3378 (2007), arXiv:0710.0051
- [24] H. Y. Bi, R. Y. Zhang, X. G. Wu *et al.*, *Phys. Rev. D* **95**, 074020 (2017), arXiv:1702.07181
- [25] J. W. Zhang, X. G. Wu, T. Zhong *et al.*, *Phys. Rev. D* **83**, 034026 (2011), arXiv:1101.1130
- [26] J. Jiang, X. G. Wu, Q. L. Liao *et al.*, *Phys. Rev. D* **86**, 054021 (2012), arXiv:1208.3051
- [27] J. Jiang, X. G. Wu, S. M. Wang *et al.*, *Phys. Rev. D* **87**, 054027 (2013), arXiv:1302.0601

- [28] A. P. Martynenko and A. M. Trunin, *Phys. Rev. D* **89**, 014004 (2014), arXiv:1308.3998
- [29] Z. J. Yang and X. X. Zhao, *Chin. Phys. Lett.* **31**, 091301 (2014), arXiv:1408.5584
- [30] Z. J. Yang, P. F. Zhang, and Y. J. Zheng, *Chin. Phys. Lett.* **31**, 051301 (2014)
- [31] A. P. Martynenko and A. M. Trunin, *Eur. Phys. J. C* **75**, 138 (2015), arXiv:1405.0969
- [32] W. K. Lai and A. K. Leibovich, *Phys. Rev. D* **91**, 054022 (2015), arXiv:1410.2091
- [33] S. Koshkarev and V. Anikeev, *Phys. Lett. B* **765**, 171-174 (2017), arXiv:1605.03070
- [34] S. Koshkarev, *Acta Phys. Polon. B* **48**, 163 (2017), arXiv:1610.06125
- [35] S. Groote and S. Koshkarev, *Eur. Phys. J. C* **77**, 509 (2017), arXiv:1704.02850
- [36] X. Yao and B. Müller, *Phys. Rev. D* **97**, 074003 (2018), arXiv:1801.02652
- [37] C. H. Chang, C. F. Qiao, J. X. Wang *et al.*, *Phys. Rev. D* **73**, 094022 (2006), arXiv:hep-ph/0601032
- [38] G. Chen, X. G. Wu, J. W. Zhang *et al.*, *Phys. Rev. D* **89**, 074020 (2014), arXiv:1401.6269
- [39] X. C. Zheng, C. H. Chang, and Z. Pan, *Phys. Rev. D* **93**, 034019 (2016), arXiv:1510.06808
- [40] G. Chen, C. H. Chang, and X. G. Wu, *Eur. Phys. J. C* **78**, 801 (2018), arXiv:1808.03174
- [41] A. V. Berezhnoy, I. N. Belov and A. K. Likhoded, *Int. J. Mod. Phys. A* **34**, 1950038 (2019), arXiv:1811.07382
- [42] G. Chen, X. G. Wu and S. Xu, *Phys. Rev. D* **100**, 054022 (2019), arXiv:1903.00722
- [43] X. G. Wu, *Sci. China Phys. Mech. Astron.* **63**, 221063 (2020), arXiv:1912.01953
- [44] J. J. Niu, L. Guo, H. H. Ma *et al.*, *Phys. Rev. D* **98**, 094021 (2018), arXiv:1810.03834
- [45] P. H. Zhang, L. Guo, X. C. Zheng *et al.*, *Phys. Rev. D* **105**, 034016 (2022), arXiv:2202.01579
- [46] J. J. Niu, L. Guo, H. H. Ma *et al.*, *Eur. Phys. J. C* **79**, 339 (2019), arXiv:1904.02339
- [47] X. Luo, Y. Z. Jiang, G. Y. Zhang *et al.*, arXiv: 2206.05965
- [48] V. D. Barger, K. m. Cheung, and W. Y. Keung, *Phys. Rev. D* **41**, 1541 (1990)
- [49] E. Braaten, K. m. Cheung, and T. C. Yuan, *Phys. Rev. D* **48**, 4230 (1993), arXiv:hep-ph/9302307[hep-ph]
- [50] Q. L. Liao, Y. Yu, Y. Deng *et al.*, *Phys. Rev. D* **91**, 114030 (2015), arXiv:1505.03275
- [51] J. B. Guimarães da Costa *et al.* (CEPC Study Group), arXiv: 1811.10545
- [52] Q. Qin, Y. J. Shi, W. Wang *et al.*, *Phys. Rev. D* **105**(3), L031902 (2022), arXiv:2108.06716
- [53] J. P. Ma and Z. G. Si, *Phys. Lett. B* **568**, 135 (2003), arXiv:hep-ph/0305079
- [54] Z. Sun and X. G. Wu, *JHEP* **07**, 034 (2020), arXiv:2004.01012
- [55] A. Petrelli, M. Cacciari, M. Greco *et al.*, *Nucl. Phys. B* **514**, 245 (1998), arXiv:hep-ph/9707223
- [56] G. T. Bodwin, D. K. Sinclair, and S. Kim, *Phys. Rev. Lett.* **77**, 2376 (1996), arXiv:hep-ph/9605023
- [57] E. Bagan, H. G. Dosch, P. Gosdzinsky *et al.*, *Z. Phys. C* **64**, 57 (1994), arXiv:hep-ph/9403208
- [58] V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Nucl. Phys. B* **569**, 473 (2000), arXiv:hep-ph/9905359
- [59] C. H. Chang, J. X. Wang, and X. G. Wu, *Phys. Rev. D* **77**, 014022 (2008), arXiv:0711.1898
- [60] X. G. Wu, *Phys. Lett. B* **671**, 318 (2009), arXiv:0805.4511
- [61] G. T. Bodwin and A. Petrelli, *Phys. Rev. D* **66**, 094011 (2002), arXiv:hep-ph/0205210
- [62] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018)
- [63] G. Weiglein *et al.* (LHC/ILC Study Group), *Phys. Rept.* **426**, 47 (2006), arXiv:hep-ph/0410364
- [64] A. Ali, A. Y. Parkhomenko, Q. Qin *et al.*, *Phys. Lett. B* **782**, 412 (2018), arXiv:1805.02535
- [65] L. C. Deng, X. G. Wu, Z. Yang *et al.*, *Eur. Phys. J. C* **70**, 113 (2010), arXiv:1009.1453
- [66] F. S. Yu, H. Y. Jiang, R. H. Li *et al.*, *Chin. Phys. C* **42**, 051001 (2018), arXiv:1703.09086
- [67] R. Aaij *et al.* (LHCb Collaboration), *JHEP* **12**, 090 (2013), arXiv:1310.2538
- [68] P. A. Zyla *et al.* (Particle Data Group), *PTEP* **2020**, 083C01 (2020)
- [69] L. Gladilin, *Eur. Phys. J. C* **75**, 19 (2015), arXiv:1404.3888
- [70] W. Wang, F. S. Yu, and Z. X. Zhao, *Eur. Phys. J. C* **77**, 781 (2017), arXiv:1707.02834
- [71] R. Aaij *et al.* (LHCb Collaboration), arXiv: 1808.08865
- [72] R. Aaij *et al.* (LHCb), *Eur. Phys. J. C* **73**, 2373 (2013), arXiv:1208.3355