



# Oscillator approach to quantization of $AdS_5 \times S^5$ superparticle in twistor formulation

D.V. Uvarov

NSC Kharkov Institute of Physics and Technology, 61108 Kharkov, Ukraine

## ARTICLE INFO

### Article history:

Received 14 May 2020

Received in revised form 22 December 2020

Accepted 8 February 2021

Available online 15 February 2021

Editor: N. Lambert

## ABSTRACT

Using known relation between  $SU(2, 2|4)$  supertwistors and  $SU(2)$  bosonic and fermionic oscillators we identify the physical states of quantized massless  $AdS_5 \times S^5$  superparticle in supertwistor formulation and discuss how they fit into the spectrum of fluctuations of IIB supergravity on  $AdS_5 \times S^5$  superbackground.

© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

Significant progress attained over the last decade and a half in perturbative  $N = 4$  super-Yang-Mills theory (see, e.g. [1] for review) was substantially triggered by the seminal work [2] that, based on the previous findings [3], [4], showed how string and twistor theories can be coupled in a synergetic way. One of the developments was the construction of (ambi)twistor action functionals for  $N = 4$  super-Yang-Mills theory in [5], [6], [7] that results in efficient rearrangement of the perturbative series expansion.

Since  $N = 4$  super-Yang-Mills theory has a dual description as Type IIB string theory on  $AdS_5 \times S^5$  superbackground it is interesting to apply twistor methods also to the exploration of this duality [8], [9]. This necessitates working out supertwistor formulation on the  $AdS_5 \times S^5$  superbackground for the superstring [10], [11] and Type IIB supergravity [12], [13] that should start from identifying proper supertwistor variables. One of the feasible ways towards this goal is to consider supertwistor formulation on the  $AdS_5 \times S^5$  superbackground of the massless superparticle model that arises in the infinite-tension limit of the superstring.

Such approach was pursued in [14] (see also [15]) based on the generalization of the extended superparticle model [16] that has extra dynamical variables to linearly realize  $SU(2, 2|4)$  symmetry and extra gauge symmetries to get the same physical degrees of freedom as in the conventional superparticle models. In the model of Ref. [14] appropriately fixing part of the gauge symmetries allows to express the superparticle's Lagrangian in terms of the supertwistors arranged into two  $SU(2)$  doublets, one of which is a straight-forward generalization of the Ferber supertwistors [17], while components of another  $SU(2)$  doublet of supertwistors have different Grassmann parity, i.e.  $SU(2, 2)$  components are Grassmann odd, whereas  $SU(4)$  components are Grassmann even. Besides that supertwistors satisfy  $SU(2, 2|4)$ -invariant quadratic constraints that are generators of the  $su(2|2)$  gauge symmetry of the superparticle's action.

In [18], [19] we have found that this supertwistor Lagrangian can be obtained also from the supertwistor formulation of the conventional massless superparticle model on  $AdS_5 \times S^5$  superbackground [20], [21], [22], [23], [24]. One of the benefits of this approach is that it yields the incidence relations connecting supertwistor components and  $AdS_5 \times S^5$  superspace coordinates via the  $PSU(2, 2|4)/(SO(1, 4) \times SO(5))$  supercoset representative. Let us note that unlike the superspace formulation of the superparticle, in which the Lagrangian involves highly non-linear expressions for the  $AdS_5 \times S^5$  supervielbein bosonic components, kinetic term and the constraints in the supertwistor formulation are quadratic facilitating Dirac quantization. In [19] it was obtained the set of equations for the superparticle's wave function in the space of superambitwistors, however, open problem is to elaborate on the details of the Penrose transform in order to establish a mapping of the components of the wave function to the fields of Type IIB supergravity multiplet compactified on  $AdS_5 \times S^5$  superbackground.

In this note we use close relationship [25] of  $SU(2, 2|4)$  supertwistors and  $SU(2)$  bosonic and fermionic oscillators that were used to construct positive energy unitary irreducible representations (uir) of  $PSU(2, 2|4)$  [26], [12], to provide representation-theoretic characterization of the quantum states of  $AdS_5 \times S^5$  superparticle. Similar analysis of the massive bosonic particle model on  $AdS_5$  in the twistor

E-mail address: [d\\_uvarov@hotmail.com](mailto:d_uvarov@hotmail.com).

formulation [27]<sup>1</sup> was performed in [29]. After briefly recapitulating four-supertwistor formulation of the massless superparticle, we proceed to study oscillator realization of the  $u(2, 2|4)$  global symmetry and  $su(2|2)$  gauge symmetry generators, analyze gauge-invariant subspace of the quantum states of the superparticle and show how they fit the fluctuation spectrum of IIB supergravity over the  $AdS_5 \times S^5$  superbackground.

## 2. Four-supertwistor formulation of massless superparticle on $AdS_5 \times S^5$ superbackground

In [18], [19] it was derived four-supertwistor representation of the Lagrangian of the massless  $AdS_5 \times S^5$  superparticle starting from the first-order form of its superspace Lagrangian. Corresponding action functional reads

$$S = \int d\tau \mathcal{L}_{4\text{-stwistor}}$$

$$\begin{aligned} \mathcal{L}_{4\text{-stwistor}} = & \frac{i}{2} \left( \bar{Z}_{\mathcal{A}}^i \dot{Z}_i^{\mathcal{A}} - \dot{\bar{Z}}_{\mathcal{A}}^i Z_i^{\mathcal{A}} \right) + \frac{i}{2} \left( \bar{\Psi}_{\mathcal{A}}^q \dot{\Psi}_q^{\mathcal{A}} - \dot{\bar{\Psi}}_{\mathcal{A}}^q \Psi_q^{\mathcal{A}} \right) \\ & + \Lambda^j{}_i \left( \bar{Z}_{\mathcal{A}}^i Z_j^{\mathcal{A}} - \frac{1}{2} \delta_j^i \bar{Z}_{\mathcal{A}}^k Z_k^{\mathcal{A}} \right) + \Lambda^p{}_q \left( \bar{\Psi}_{\mathcal{A}}^q \Psi_p^{\mathcal{A}} - \frac{1}{2} \delta_p^q \bar{\Psi}_{\mathcal{A}}^r \Psi_r^{\mathcal{A}} \right) \\ & + \Lambda \left( \bar{Z}_{\mathcal{A}}^i Z_i^{\mathcal{A}} + \bar{\Psi}_{\mathcal{A}}^q \Psi_q^{\mathcal{A}} \right) + i \Lambda_q^i \bar{\Psi}_{\mathcal{A}}^q Z_i^{\mathcal{A}} + i \bar{\Lambda}_i^q \bar{Z}_{\mathcal{A}}^i \Psi_q^{\mathcal{A}}, \end{aligned} \quad (1)$$

where components of the two doublets of  $SU(2, 2|4)$  supertwistors have different Grassmann parity, that is Ferber or  $c$ -type supertwistors

$$Z_i^{\mathcal{A}} = \begin{pmatrix} Z_{i\alpha}^{\mathcal{A}} \\ \eta_i^{\mathcal{A}} \end{pmatrix}, \quad \bar{Z}_{\mathcal{A}}^i = (Z_i^{\mathcal{B}})^{\dagger} \mathcal{H}^{\mathcal{B}}{}_{\mathcal{A}} = (\bar{Z}_{\alpha}^i \bar{\eta}_{\mathcal{A}}^i), \quad \mathcal{H}^{\mathcal{B}}{}_{\mathcal{A}} = \begin{pmatrix} H^{\mathcal{B}}{}_{\alpha} & 0 \\ 0 & \delta_{\mathcal{A}}^{\mathcal{B}} \end{pmatrix} \quad (2)$$

have Grassmann-even  $SU(2, 2)$  components  $Z_i^{\mathcal{A}}$ ,  $\bar{Z}_{\mathcal{A}}^i$  and Grassmann-odd  $SU(4)$  components  $\eta_i^{\mathcal{A}}$ ,  $\bar{\eta}_{\mathcal{A}}^i$ , whereas  $a$ -type supertwistors

$$\Psi_q^{\mathcal{A}} = \begin{pmatrix} \xi_q^{\mathcal{A}} \\ L_q^{\mathcal{A}} \end{pmatrix}, \quad \bar{\Psi}_{\mathcal{A}}^q = (\Psi_q^{\mathcal{B}})^{\dagger} \mathcal{H}^{\mathcal{B}}{}_{\mathcal{A}} = (\bar{\xi}_{\alpha}^q \bar{L}_{\mathcal{A}}^q) \quad (3)$$

have Grassmann-odd  $SU(2, 2)$  components  $\xi_q^{\mathcal{A}}$ ,  $\bar{\xi}_{\alpha}^q$  and Grassmann-even  $SU(4)$  components  $L_q^{\mathcal{A}}$ ,  $\bar{L}_{\mathcal{A}}^q$ . The action (1) coincides with that obtained in Ref. [14] by partial gauge fixing  $2T$  superparticle model in  $2 + 10$  dimensions. The advantage of the approach pursued in [18], [19] is not only in establishing the direct connection with the superparticle's formulation in  $AdS_5 \times S^5$  superspace but also in obtaining the incidence relations for the components of supertwistors and superspace coordinates parametrizing  $PSU(2, 2|4)/(SO(1, 4) \times SO(5))$  supercoset representative  $\mathcal{G}$

$$Z_i^{\mathcal{A}} = G^{\mathcal{A}}{}_{\mathcal{B}} v_i^{\mathcal{B}}, \quad v_i^{\mathcal{B}} = \begin{pmatrix} v_i^{\mathcal{B}} \\ 0 \end{pmatrix}, \quad \bar{Z}_{\mathcal{A}}^i = v_i^{\mathcal{B}} G^{-1\mathcal{B}}{}_{\mathcal{A}} \quad (4)$$

and

$$\Psi_q^{\mathcal{A}} = G^{\mathcal{A}}{}_{\mathcal{B}} \ell_q^{\mathcal{B}}, \quad \ell_q^{\mathcal{B}} = \begin{pmatrix} 0 \\ \ell_q^{\mathcal{B}} \end{pmatrix}, \quad \bar{\Psi}_{\mathcal{A}}^q = \ell_q^{\mathcal{B}} G^{-1\mathcal{B}}{}_{\mathcal{A}}. \quad (5)$$

Unlike the eight-supertwistor formulation also considered in [19], in the four-supertwistor formulation there are only the first-class constraints introduced in (1) via the Lagrange multipliers – seven bosonic

$$\bar{Z}_{\mathcal{A}}^i Z_j^{\mathcal{A}} - \frac{1}{2} \delta_j^i \bar{Z}_{\mathcal{A}}^k Z_k^{\mathcal{A}} \approx 0, \quad (6a)$$

$$\bar{\Psi}_{\mathcal{A}}^q \Psi_p^{\mathcal{A}} - \frac{1}{2} \delta_p^q \bar{\Psi}_{\mathcal{A}}^r \Psi_r^{\mathcal{A}} \approx 0, \quad (6b)$$

$$T = \bar{Z}_{\mathcal{A}}^i Z_i^{\mathcal{A}} + \bar{\Psi}_{\mathcal{A}}^q \Psi_q^{\mathcal{A}} \approx 0 \quad (6c)$$

and eight fermionic

$$\bar{\Psi}_{\mathcal{A}}^q Z_i^{\mathcal{A}} \approx 0, \quad \bar{Z}_{\mathcal{A}}^i \Psi_q^{\mathcal{A}} \approx 0 \quad (7)$$

that are the generators of the classical  $su(2|2)$  gauge symmetry of the superparticle's action. It was shown in [18], [19] that, when this gauge symmetry is taken into account, the number of gauge-invariant physical degrees of freedom in the supertwistor formulation is the same as in the superspace formulation. The presence of only the first-class constraints greatly facilitates Dirac quantization of the model but requires insights into the Penrose transform of the functions of superambitwistors of  $c$ - and  $a$ -type as was discussed in [19], where the set of the first-order differential equations for the superparticle's wave function was derived.

<sup>1</sup> This twistor formulation of the bosonic particle on  $AdS_5$  was recently revisited and generalized to dimensions 4 and 7 in [28], for which one can benefit from using two-component spinors with complex, real and quaternionic entries respectively.

### 3. From $SU(2, 2|4)$ supertwistors to $SU(2)$ oscillators

Let us start this section with a few remarks regarding oscillator construction of the representations of  $D = 4$  conformal algebra and its  $N = 4$  supersymmetric extension. Bosonic oscillators have been utilized since 1960's in the field-theoretical models with underlying conformal symmetry to construct the generators of  $so(2, 4) \cong su(2, 2)$  algebra and describe their quantum states, as well as to realize certain  $su(2, 2)$  representations (see, e.g. [30], [31] and notably [32], where quantized twistors and oscillators were treated in the uniform way and used to describe ladder representations). In the mid 1980's the oscillator approach was advanced further in [33], [26] and applied in [12] to construct positive energy (lowest weight) uirs of  $su(2, 2)$  (of which ladder representations constitute a particular case) that arise upon the compactification of Type IIB supergravity on  $AdS_5 \times S^5$  superbackground. Subsequently the oscillator approach was also used to study the  $AdS_5/CFT_4$  correspondence [34], [35].

Infinite-dimensional positive energy uirs, that can be constructed with the aid of quantized bosonic oscillators, have (half-)integer labels and correspond to finite-component fields in  $AdS_5$  or on its conformal boundary given by the 4-dimensional Minkowski space-time. Their classification was finalized in [36] based on the method of induced representations from the maximal parabolic subalgebra  $(D \oplus sl(2, \mathbb{C})) \ltimes K \subset su(2, 2)$ , where  $D$  and  $K$  are generators of the conformal dilatations and boosts.<sup>2</sup> As was shown in Ref. [37] positive energy uirs with (half-)integer labels correspond to the holomorphic discrete series uirs of  $su(2, 2)$  or its limits. Recently there was proposed a generalization of the oscillator approach to describe uirs of  $su(p, q)$  and its superextensions with non-(half-)integer labels [38].

The construction of Ref. [33] makes use of the bosonic oscillators carrying fundamental representation(s) of the semisimple part of the maximal compact subalgebra of the non-compact Lie algebra in question. Upon generalization [26] to non-compact simple superalgebras they acquire superpartners – fermionic oscillators transforming in the fundamental representation of the subalgebra of  $R$ -symmetry and generating its finite-dimensional uirs. For  $psu(2, 2|4)$  superalgebra (recall that  $su(2, 2|4)$  is not simple) this construction implies introduction of the two species of bosonic oscillators associated with two  $su(2)$  subalgebras of the maximal compact subalgebra of  $su(2, 2)$  and two species of fermionic oscillators for two  $su(2)$  subalgebras of  $su(4)$ . Positive energy uirs of  $psu(2, 2|4)$  are grouped into supermultiplets – massive (long), massless (short) and doubleton supermultiplets – generated from respective lowest-weight vectors (lwv) that carry certain finite-dimensional uirs of the maximal compact subalgebra of  $su(2, 2)$  and  $su(2) \oplus su(2) \subset su(4)$ . Each member of the supermultiplet corresponds to a finite-component field in  $AdS_5$  or on its conformal boundary transforming according to some finite-dimensional representation of the  $su(4)$   $R$ -symmetry algebra. Classification of  $psu(2, 2|4)$  supermultiplets was done in [39] without employing oscillators (see also [38] for consideration relying on oscillators).

Let us now present our definition [40] of the bosonic  $SU(2)$  oscillators. Introduce  $SL(2, \mathbb{C})$  constituents of the  $SU(2, 2)$  components of  $c$ -type supertwistors

$$Z_i^\alpha = \begin{pmatrix} \mu_i^\alpha \\ \bar{\Lambda}_{\dot{\alpha}i} \end{pmatrix}, \quad \bar{Z}_\alpha^i = (\Lambda_\alpha^i \bar{\mu}^{\dot{\alpha}i}). \tag{8}$$

Then two species of the  $SU(2)$  bosonic oscillators are defined by the linear combinations of the twistor components (8)

$$\begin{pmatrix} a_i^\alpha \\ b_i^\alpha \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} -I & I \\ I & I \end{pmatrix} \begin{pmatrix} \mu_i^\alpha \\ \bar{\Lambda}_{\dot{\alpha}i} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} -\mu_i^\alpha + \bar{\Lambda}_{\dot{\alpha}i} \\ \mu_i^\alpha + \bar{\Lambda}_{\dot{\alpha}i} \end{pmatrix} \tag{9}$$

and

$$(a_\alpha^i \ b_\alpha^i) = \frac{1}{\sqrt{2}} (\bar{\mu}^{\dot{\alpha}i} \ \Lambda_\alpha^i) \begin{pmatrix} -I & I \\ I & I \end{pmatrix} = \frac{1}{\sqrt{2}} (-\bar{\mu}^{\dot{\alpha}i} + \Lambda_\alpha^i \ \bar{\mu}^{\dot{\alpha}i} + \Lambda_\alpha^i), \tag{10}$$

where  $I$  is the unit  $2 \times 2$  matrix. Unitary transformation (9)-(10) describes transition from the basis of  $D = 2 + 4$   $\gamma$ -matrices, in which the generator of conformal dilatations  $D$  and  $so(1, 3)$  generators are block-diagonal in the spinor representation, to the basis, in which generators of the maximal compact subalgebra  $\mathcal{K} = su(2) \oplus su(2) \oplus u(1) \subset so(2, 4)$  take block-diagonal form. The generators of the  $su(2)$  algebras in  $\mathcal{K}$  are given by the linear combinations of the  $su(2) \subset so(1, 3)$  generators and  $so(2, 4) - so(1, 3)$  generators that have no time indices (see [35] and Appendix B in [38] for details). The former basis corresponds to the realization of the  $su(2, 2)$  algebra as  $D = 4$  conformal algebra that is at the heart of the twistor theory, while the latter underlies the oscillator approach. Transformation (9)-(10) amounts to the diagonalization of the matrix  $H^\alpha_\beta$  that enters the definition of dual supertwistors in (2) and (3)

$$H^\alpha_\beta = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \rightarrow \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix}. \tag{11}$$

Note some abuse of the notation in (9)-(10) and below since, as in [38], small lightface Greek letters stand for the indices of both  $sl(2, \mathbb{C})$  two-component spinors and two-component spinors of the two  $su(2)$  algebras in  $\mathcal{K}$ . Moreover, following [33] we use the same letters to label indices of  $a$ - and  $b$ -oscillators although they carry spinor representations of different  $su(2)$  algebras and are not contracted. It suffices to take into account that  $sl(2, \mathbb{C})$  spinor indices are carried by the  $SU(2, 2)$  twistor components and space-time fields (discussed at the end of Section 5), whereas oscillators and  $psu(2, 2|4)$  supersymmetry generators constructed out of them carry  $su(2)$  spinor indices.

Dirac bracket (D.B.) relations for the bosonic components of  $c$ -type supertwistors that follow from the kinetic term of the Lagrangian (1)

$$\{Z_i^\alpha, \bar{Z}_\beta^j\}_{D.B.} = i\delta_i^j \delta_\beta^\alpha \tag{12}$$

<sup>2</sup> Equivalence of these approaches was discussed in [35] including a one-to-one mapping of the representation labels.

in the quantum theory become commutators<sup>3</sup>

$$[Z_i^\alpha, \bar{Z}_\beta^j] = \delta_i^j \delta_\beta^\alpha \quad (13)$$

that upon above transformation translate into commutators of  $a$ - and  $b$ -oscillators

$$[a_\alpha^i, a_j^\beta] = \delta_j^i \delta_\alpha^\beta, \quad [b_i^\alpha, b_\beta^j] = \delta_i^j \delta_\beta^\alpha. \quad (14)$$

Thus oscillators  $a_i^\alpha$  and  $b_i^\alpha$  are treated as raising operators, whereas  $a_\alpha^i$  and  $b_i^\alpha$  are lowering operators annihilating oscillator vacuum  $|0\rangle$ .

Similarly D.B. relations of the odd components of  $c$ -type supertwistors (2) read

$$\{\eta_i^A, \bar{\eta}_B^j\}_{D.B.} = i \delta_i^j \delta_B^A. \quad (15)$$

On the quantum level they become anticommutators

$$\{\eta_i^A, \bar{\eta}_B^j\} = \delta_i^j \delta_B^A \quad (16)$$

that can be viewed as defining relations of the two sets of  $SU(4)$  fermionic oscillators. Decomposing  $SU(4)$  (anti)fundamental representation on the fundamental representations of its two  $SU(2)$  subgroups introduces two kinds of  $SU(2)$  fermionic oscillators [12], [34]<sup>4</sup>

$$\eta_i^A = \begin{pmatrix} \alpha_i^a \\ \beta_i^{\dot{a}} \end{pmatrix}, \quad \bar{\eta}_A^i = \begin{pmatrix} \alpha_a^i & \beta_{\dot{a}}^i \end{pmatrix} \quad (17)$$

that satisfy

$$\{\alpha_i^a, \alpha_b^j\} = \delta_b^j \delta_i^a, \quad \{\beta_i^{\dot{a}}, \beta_b^{\dot{j}}\} = \delta_b^{\dot{j}} \delta_i^{\dot{a}}. \quad (18)$$

Admissible choice is to treat  $\alpha_i^a$  and  $\beta_i^{\dot{a}}$  as raising oscillators, then  $\alpha_a^i$  and  $\beta_{\dot{a}}^i$  are lowering ones annihilating oscillator vacuum  $|0\rangle$ .

Note that in such a way we get two copies of bosonic and fermionic oscillators of each kind transforming as the doublet of one of the  $su(2)$  subalgebras of  $su(2|2)$  gauge superalgebra that is an instance of the color algebras in the oscillator approach. Note also that to realize  $su(2, 2)$  and  $psu(2, 2|4)$  generators it suffices to take just one copy of the oscillators of each kind. Using these minimal realization of  $su(2, 2)$  it is possible to construct only ladder uirs (named doubletons in the oscillator approach) and their supermultiplets, among which the  $N = 4$  super-Yang-Mills multiplet is the only self-conjugate one [12], [34]. Other positive energy  $su(2, 2)$  (and  $psu(2, 2|4)$ ) uirs can be obtained by decomposing tensor products of finitely many (super)doubletons.

One can analogously introduce oscillators associated with the  $a$ -type supertwistors (3). Namely, taking linear combinations of the  $SL(2, \mathbb{C})$  constituents of the odd components of  $a$ -type supertwistors

$$\xi_q^\alpha = \begin{pmatrix} m_q^\alpha \\ \bar{n}_{\dot{\alpha}q} \end{pmatrix}, \quad \bar{\xi}_\alpha^q = (n_\alpha^q \bar{m}^{\dot{\alpha}q}) \quad (19)$$

defines another two species of  $SU(2)$  fermionic oscillators

$$\begin{aligned} \omega_q^\alpha &= \frac{1}{\sqrt{2}}(-m_q^\alpha + \bar{n}_{\dot{\alpha}q}), & \omega_\alpha^q &= \frac{1}{\sqrt{2}}(-\bar{m}^{\dot{\alpha}q} + n_\alpha^q), \\ \varphi_q^\alpha &= \frac{1}{\sqrt{2}}(m_q^\alpha + \bar{n}_{\dot{\alpha}q}), & \varphi_\alpha^q &= \frac{1}{\sqrt{2}}(\bar{m}^{\dot{\alpha}q} + n_\alpha^q) \end{aligned} \quad (20)$$

in addition to (17). The fact that Grassmann parity of the  $SU(2, 2)$  components of  $a$ -type supertwistors differs from that of the  $c$ -type supertwistors has far-reaching consequences for the future analysis. In particular, from the anticommutators of  $\xi_q^\alpha$  and  $\bar{\xi}_\beta^p$

$$\{\xi_q^\alpha, \bar{\xi}_\beta^p\} = \delta_q^p \delta_\beta^\alpha \quad (21)$$

it follows that

$$\{\omega_q^\alpha, \omega_\beta^p\} = -\delta_q^p \delta_\beta^\alpha, \quad \{\varphi_q^\alpha, \varphi_\beta^p\} = \delta_q^p \delta_\beta^\alpha, \quad (22)$$

i.e.  $\omega$ -oscillators satisfy wrong-sign anticommutation relations.

Grassmann-even components of  $a$ -type supertwistors that in the quantum theory satisfy commutation relations

$$[L_q^A, \bar{L}_B^p] = \delta_q^p \delta_B^A \quad (23)$$

admit decomposition on the  $SU(2)$  constituents in analogy with (17)

$$L_q^A = \begin{pmatrix} c_q^a \\ d_q^{\dot{a}} \end{pmatrix}, \quad \bar{L}_A^q = (c_a^q \bar{d}_{\dot{a}}^q). \quad (24)$$

This yields two extra species of  $SU(2)$  bosonic oscillators. Commutation relations for them can be read off from (23)

$$[c_q^a, c_b^p] = \delta_q^p \delta_b^a, \quad [d_q^{\dot{a}}, d_b^{\dot{p}}] = \delta_q^p \delta_b^{\dot{a}}. \quad (25)$$

As will be shown in the next section for one type of the introduced  $SU(2)$  bosonic oscillators the commutation relations are necessarily the wrong-sign ones.

<sup>3</sup> We do not place hats over the quantum operators not to overburden the notation. Hopefully this will not cause a confusion.

<sup>4</sup> Another option – to use  $SU(4)$  oscillators as they stand to construct  $psu(2, 2|4)$  uirs was considered in [41], [42], [38].

#### 4. Oscillator realization of global $u(2, 2|4)$ symmetry

As was mentioned above superparticle's action (1) is invariant under the  $su(2|2)$  gauge symmetry. The constraints (6) and (7) span the  $su(2|2)$  superalgebra named color superalgebra in [15], [14] since it generalizes bosonic color symmetries that accompany oscillator realizations of  $su(2, 2)$  and  $psu(2, 2|4)$  generators [12], [34], [35], [41]. Superparticle's action (1) is also invariant under  $U(2, 2|4)$  global symmetry generated by

$$z_i^A \bar{z}_B^i - \Psi_q^A \bar{\Psi}_B^q = u(2, 2|4). \quad (26)$$

Important property of these  $u(2, 2|4)$  generators is that they (anti)commute with the generators of the  $su(2|2)$  gauge symmetry due to the color symmetry invariance.<sup>5</sup> When expressed in terms of oscillators introduced in (9), (10), (17), (20) and (24), these global symmetry generators are used to build the space of states of quantized superparticle model as the certain representation space of  $u(2, 2|4)$ . These generators are of three types: quadratic in creation oscillators, quadratic in annihilation oscillators and composed of the products of creation and annihilation oscillators in accordance with the three-grading structure of the  $(ps)u(2, 2|4)$  superalgebra [34]. This three-grading structure allows to choose which of the oscillators associated with the  $a$ -type supertwistors represent creation and annihilation operators once such choice is made for the oscillators related to the  $c$ -type supertwistors.

Consider first the  $su(2, 2) \subset u(2, 2|4)$  algebra. Three-grading decomposition of its generators is as follows

$$\mathfrak{g}_{(0)} = \{E, L^\alpha_\beta, R^\alpha_\beta\}, \quad \mathfrak{g}_{(-1)} = \{T^{(-)\alpha}_\beta\}, \quad \mathfrak{g}_{(+1)} = \{T^{(+)\alpha}_\beta\}: \quad (27)$$

$$2E = a_i^\alpha a_\alpha^i + b_\alpha^i b_i^\alpha - \omega_q^\alpha \omega_\alpha^q + \varphi_q^\alpha \varphi_\alpha^q, \quad (28a)$$

$$L^\alpha_\beta = a_i^\alpha a_\beta^i - \omega_q^\alpha \omega_\beta^q - \frac{1}{2} \delta_\beta^\alpha (a_i^\gamma a_\gamma^i - \omega_q^\gamma \omega_\gamma^q), \quad (28b)$$

$$R^\alpha_\beta = b_i^\alpha b_\beta^i - \varphi_q^\alpha \varphi_\beta^q - \frac{1}{2} \delta_\beta^\alpha (b_i^\gamma b_\gamma^i - \varphi_q^\gamma \varphi_\gamma^q), \quad (28c)$$

$$T^{(-)\alpha}_\beta = b_i^\alpha a_\beta^i - \varphi_q^\alpha \omega_\beta^q, \quad (28d)$$

$$T^{(+)\alpha}_\beta = a_i^\alpha b_\beta^i - \omega_q^\alpha \varphi_\beta^q. \quad (28e)$$

Above expressions deserve a number of comments. From (28d) and (28e) it follows that  $\omega_\beta^q$  and  $\varphi_q^\alpha$  are annihilation operators, whereas  $\omega_q^\alpha$  and  $\varphi_\beta^q$  are creation operators. In (27)  $\mathfrak{g}_{(0)}$  is spanned by the generators of the maximal compact subalgebra  $u(1)_E \oplus su(2) \oplus su(2)$  of  $su(2, 2)$ , where  $E$  is the  $AdS_5$  energy (conformal dimension) operator that defines grading of  $su(2, 2)$  generators:

$$[E, \mathfrak{g}_{(\sigma)}] = \sigma \mathfrak{g}_{(\sigma)} \quad \sigma = 0, \pm 1. \quad (29)$$

As a result  $su(2, 2)$  commutation relations can be written in concise schematic form

$$[\mathfrak{g}_{(\sigma)}, \mathfrak{g}_{(\sigma')}] = \mathfrak{g}_{(\sigma+\sigma')}. \quad (30)$$

This three grading decomposition can be viewed as a modification of the Cartan decomposition. Expression (28a) corresponds to creation-annihilation ordering of oscillators and was obtained from the manifestly Hermitian one by moving raising oscillators to the left. Introducing oscillator number operators  $N_{(a)} = a_i^\alpha a_\alpha^i$ ,  $N_{(b)} = b_\alpha^i b_i^\alpha$ ,  $N_{(\omega)} = -\omega_q^\alpha \omega_\alpha^q$  and  $N_{(\varphi)} = \varphi_q^\alpha \varphi_\alpha^q$  it can be brought to the form

$$E = \frac{1}{2}(N_{(a)} + N_{(b)} + N_{(\omega)} + N_{(\varphi)}). \quad (31)$$

Note that  $N_{(\omega)}$  includes extra minus sign that correlates with the wrong sign in the anticommutator of the  $\omega$ -oscillators in (22). Though  $E$  is defined to be non-negative, in (31) there is no positive  $c$ -number term in distinction to the case when  $su(2, 2)$  generators are realized solely in terms of the bosonic oscillators [12], [34]. Such positive  $c$ -number contribution ensures that obtained irreducible representations (irreps) lie above or saturate respective unitarity bounds. There is no ordering ambiguity in the expressions (28b) and (28c) for  $L^\alpha_\beta$  and  $R^\alpha_\beta$  that generate two  $su(2)$  subalgebras of  $su(2, 2)$ . Lwv's carry spin  $s_1$  and  $s_2$  irreps of these  $su(2)$  subalgebras and are eigenvectors of  $E$  so that two spins  $s_1, s_2$  and  $AdS_5$  energy are, as is common, used to label the  $su(2, 2)$  irreps. Whatever oscillator realization of the  $su(2, 2)$  generators is, positive energy representations are constructed by repeated action of  $T^{(+)\alpha}_\beta$  generators on the lww's annihilated by  $T^{(-)\alpha}_\beta$ . In the model under consideration lww corresponding to the superparticle's ground state is also annihilated by the  $su(2|2)$  gauge symmetry generators.

Generators of the  $su(4) \subset u(2, 2|4)$  algebra also admit three-grading decomposition

$$h_{(0)} = \{C, A^a_b, B^{\dot{a}}_{\dot{b}}\}, \quad h_{(-1)} = \{V^{(-)\dot{a}}_{\dot{b}}\}, \quad h_{(+1)} = \{V^{(+)\dot{a}}_{\dot{b}}\}: \quad (32)$$

$$2C = -\alpha_i^a \alpha_a^i - \beta_{\dot{a}}^i \beta_i^{\dot{a}} + c_q^a c_a^q - d_q^{\dot{a}} d_{\dot{a}}^q, \quad (33a)$$

$$A^a_b = \alpha_i^a \alpha_b^i - c_q^a c_b^q - \frac{1}{2} \delta_b^a (\alpha_i^c \alpha_c^i - c_q^c c_c^q), \quad (33b)$$

$$B^{\dot{a}}_{\dot{b}} = \beta_i^{\dot{a}} \beta_{\dot{b}}^i - d_q^{\dot{a}} d_{\dot{b}}^q - \frac{1}{2} \delta_{\dot{b}}^{\dot{a}} (\beta_i^c \beta_c^i - d_q^c d_c^q), \quad (33c)$$

$$V^{(-)\dot{a}}_{\dot{b}} = \beta_i^{\dot{a}} \alpha_b^i - d_q^{\dot{a}} c_b^q, \quad (33d)$$

$$V^{(+)\dot{a}}_{\dot{b}} = \alpha_i^a \beta_b^i - c_q^a d_b^q \quad (33e)$$

<sup>5</sup> This is a manifestation of the Howe duality that has been recently discussed in [43] with the emphasis on the oscillator approach.

so that commutation relations of  $su(4)$  algebra can be brought to the form respecting this three-grading structure

$$[h_{(\tau)}, h_{(\tau')}] = h_{(\tau+\tau')}, \quad (34)$$

where grading of a generator is read off from its commutator with  $C$

$$[C, h_{(\tau)}] = -\tau h_{(\tau)} \quad \tau = 0, \pm 1. \quad (35)$$

The form of the diannihilation and dcreation operators (33d) and (33e) suggests that bosonic oscillators  $c_b^q$  and  $d_b^q$  are lowering ones, while  $c_a^q$  and  $d_b^q$  are raising. In view of (25) this choice implies that  $c$ -oscillators satisfy wrong-sign commutation relations. Then the operator  $C$  can be written in the form similar to that of the  $AdS_5$  energy operator (31)

$$C = -\frac{1}{2}(N_{(\alpha)} + N_{(\beta)} + N_{(c)} + N_{(d)}), \quad (36)$$

where  $N_{(\alpha)} = \alpha_i^a \alpha_a^i$ ,  $N_{(\beta)} = \beta_a^i \beta_i^a$ ,  $N_{(c)} = -c_a^q c_a^q$  and  $N_{(d)} = d_a^q d_a^q$  are oscillator number operators and the overall minus sign has been introduced to conform with the definition of  $C$  in [34]. It is readily seen to be non-positive definite, whereas for the  $su(4)$  uirs constructed in [12] its eigenvalues are non-negative due to the positive  $c$ -number contribution equal the number of copies of the oscillators.  $su(4)$  irreps are labeled by two spins  $j_1$  and  $j_2$  of the irreps of  $su(2)$  algebras spanned by  $A^a_b$  and  $B^a_b$  that are carried by lww's as well as eigenvalues of  $C$ . Like  $su(2, 2)$  lowest-weight representations,  $su(4)$  lowest-weight representations are constructed via repeated action of  $V^{(+)}_b^a$  on respective lww's annihilated by  $V^{(-)}_a^b$ . If  $su(4)$  generators are constructed using finite number of fermionic oscillators as in [12], [34] resulting lowest-weight representations are finite-dimensional.

Two additional bosonic generators of  $u(2, 2|4)$  are the constraint (6c) that in oscillator form equals

$$T = -N_{(a)} + N_{(b)} - N_{(\alpha)} + N_{(\beta)} - N_{(\omega)} + N_{(\varphi)} - N_{(c)} + N_{(d)} \approx 0 \quad (37)$$

and

$$U = -N_{(a)} + N_{(b)} + N_{(\alpha)} - N_{(\beta)} - N_{(\omega)} + N_{(\varphi)} + N_{(c)} - N_{(d)}. \quad (38)$$

Since  $T$  is the constraint, for all physical states of the quantized superparticle its eigenvalue is zero but eigenvalues of  $Y = -N_{(a)} + N_{(b)} - N_{(\omega)} + N_{(\varphi)}$  can be used as an additional label together with the  $su(2, 2)$  and  $su(4)$  labels.

Among 32 odd generators of  $u(2, 2|4)$  eight are constructed out of the raising oscillators

$$Q^{(+)\alpha}_a = a_i^\alpha \beta_a^i - \omega_q^\alpha d_a^q, \quad Q^{(+)\alpha}_\alpha = \alpha_i^a b_\alpha^i - c_q^\alpha \varphi_\alpha^q \quad (39)$$

and belong to the  $+1$  eigenstate w.r.t. the grading defined by  $E - C$ . Their repeated action on the  $su(2, 2) \oplus su(4)$  lww of the ground state of the superparticle, that will be identified in the next Section, yields all the lww's of the supermultiplet of the physical states.

## 5. Physical states of quantized superparticle

As discussed above  $u(2, 2|4)$  generators (26) (anti)commute with  $su(2|2)$  gauge symmetry generators (6) and (7), so the latter act directly on the lww's and one can examine the constraints they impose on the ground state of the quantized superparticle. Bosonic constraints (6) imply that sought for lww is a singlet of the two  $su(2)$  color subalgebras of  $su(2|2)$  and is annihilated by  $T$ . So the ground state can be constructed by acting on the vacuum  $|0\rangle$  with the products of the color-singlet combinations of raising oscillators, namely, the determinants of bosonic oscillators  $\Delta_{(a)} = \frac{1}{2} \varepsilon^{ij} \varepsilon_{\alpha\beta} a_i^\alpha a_j^\beta$ ,  $\Delta_{(b)} = \frac{1}{2} \varepsilon_{ij} \varepsilon^{\alpha\beta} b_\alpha^i b_\beta^j$ ,  $\Delta_{(c)} = \frac{1}{2} \varepsilon^{pq} \varepsilon_{ab} c_p^a c_q^b$  and  $\Delta_{(d)} = \frac{1}{2} \varepsilon_{pq} \varepsilon^{ab} d_p^a d_q^b$ , and antisymmetrized in color indices products of the fermionic oscillators  $\alpha^{ab} = \alpha_i^a \varepsilon^{ij} \alpha_j^b$ ,  $\beta_{ab} = \beta_a^i \varepsilon_{ij} \beta_b^j$ ,  $\omega^{\alpha\beta} = \omega_p^\alpha \varepsilon^{pq} \omega_q^\beta$  and  $\varphi_{\alpha\beta} = \varphi_p^\alpha \varepsilon_{pq} \varphi_q^\beta$  raised to the powers balanced in such a way as to be annihilated by  $T^{(-)\alpha}_\beta$ ,  $V^{(-)}_a^b$  and  $T$ . Among these lww's only vacuum itself is annihilated by all the fermionic constraints (7) and hence is the ground state.

Other lww's of the supermultiplet of physical states are obtained by repeated action of the supersymmetry generators (39) on the ground state. Consider first lww's that can be obtained by acting solely with  $Q^{(+)\alpha}_a$  generators. For example, the product of two such supersymmetry generators can be decomposed on three types of contributions depending on the oscillators involved

$$Q^{(+)\alpha_1}_{a_1} Q^{(+)\alpha_2}_{a_2} = a_i^{\alpha_1} a_j^{\alpha_2} \beta_{a_1}^i \beta_{a_2}^j + \omega_q^{\alpha_1} \omega_p^{\alpha_2} d_{a_1}^q d_{a_2}^p + a_i^{\alpha_1} \omega_q^{\alpha_2} \beta_{a_1}^i d_{a_2}^q - a_i^{\alpha_2} \omega_q^{\alpha_1} \beta_{a_2}^i d_{a_1}^q. \quad (40)$$

Further decomposing the products of oscillators on irreducible tensors gives two lww's:

$$-\frac{1}{4} \varepsilon_{a_1 a_2} \left( a_i^{\alpha_1} a_j^{\alpha_2} \beta^{ij} - 2\omega^{\alpha_1 \alpha_2} \Delta_{(d)} + 2a_i^{\alpha_1} \omega_q^{\alpha_2} \beta_b^i \varepsilon^{bc} d_c^q \right) |0\rangle, \quad \beta^{ij} = \beta_b^i \varepsilon^{bc} \beta_c^j \quad (41)$$

with spins  $s_1 = 1$ ,  $s_2 = 0$  with respect to  $su(2)$  subalgebras of  $su(2, 2)$  and  $j_1 = j_2 = 0$  with respect to  $su(2)$  subalgebras of  $su(4)$ , and

$$\frac{1}{4} \varepsilon^{\alpha_1 \alpha_2} \left( 2\Delta_{(a)} \beta_{a_1 a_2} - \omega_{pq} d_{(a_1)}^p d_{(a_2)}^q - 2a_i^\beta \varepsilon_{\beta\gamma} \omega_p^\gamma \beta_{(a_1)}^i d_{(a_2)}^p \right) |0\rangle, \quad \omega_{pq} = \omega_p^\beta \varepsilon_{\beta\gamma} \omega_q^\gamma \quad (42)$$

with spins  $s_1 = s_2 = 0$  and  $j_1 = 0$ ,  $j_2 = 1$  presented in the third and fourth rows of the Table 1. Since each of the other products of  $Q^{(+)\alpha}_a$ 's gives rise to single lww and the products of more than four supercharges vanish, one finds six independent lww's presented in the Table 1. Spin content of these lww's coincides with the spin content of the lww's in the first column of the Table 1 in [12]<sup>6</sup> that have

<sup>6</sup> We prefer to label lww's by four spins of four  $su(2)$  subalgebras of  $psu(2, 2|4)$  rather than by Young tableau as in [12].



**Table 1**

$su(2, 2)$  and  $su(4)$  labels of lww's obtained by acting with the powers of  $Q^{(+)\alpha}_a$  on  $|0\rangle$  and respective fields on  $AdS_5 \times S^5$ .

$su(2, 2)$ labels ( $E, s_1, s_2$ )	$su(4)$ labels ( $C, j_1, j_2$ )	$Y$	fields on $AdS_5 \times S^5$
(0, 0, 0)	(0, 0, 0)	0	$\varphi^{(1)}(y, z)$
(1/2, 1/2, 0)	(-1/2, 0, 1/2)	-1	$\lambda_{\alpha\dot{a}}^{(1)}(y, z)$
(1, 1, 0)	(-1, 0, 0)	-2	$A_{\alpha(2)}^{(1)}(y, z)$
(1, 0, 0)	(-1, 0, 1)	-2	$\varphi_{\dot{a}(2)}^{(2)}(y, z)$
(3/2, 1/2, 0)	(-3/2, 0, 1/2)	-3	$\lambda_{\alpha\dot{a}}^{(2)}(y, z)$
(2, 0, 0)	(-2, 0, 0)	-4	$\varphi^{(3)}(y, z)$

**Table 2**

$su(2, 2)$  and  $su(4)$  labels of lww's obtained by acting with the powers of  $Q^{(+)\alpha}_\alpha$  on  $|0\rangle$  and respective fields on  $AdS_5 \times S^5$ .

$su(2, 2)$ labels ( $E, s_1, s_2$ )	$su(4)$ labels ( $C, j_1, j_2$ )	$Y$	fields on $AdS_5 \times S^5$
(1/2, 0, 1/2)	(-1/2, 1/2, 0)	1	$\lambda_{\alpha\dot{a}}^{(1)}(y, z)$
(1, 0, 1)	(-1, 0, 0)	2	$A_{\alpha(2)}^{(1)}(y, z)$
(1, 0, 0)	(-1, 1, 0)	2	$\varphi_{\dot{a}(2)}^{(2)}(y, z)$
(3/2, 0, 1/2)	(-3/2, 1/2, 0)	3	$\lambda_{\alpha\dot{a}}^{(2)}(y, z)$
(2, 0, 0)	(-2, 0, 0)	4	$\bar{\varphi}^{(3)}(y, z)$

units in the second and third digits. We normalize  $AdS_5$  energy  $E$  in such a way that its eigenvalues are (half-)integer for lww's with (half-)integer  $s_1 + s_2$  and eigenvalues of  $Y$  take integer values.

Similarly repeated application of  $Q^{(+)\alpha}_\alpha$  produces five other lww's with interchanged spins  $s_1 \leftrightarrow s_2$  and  $j_1 \leftrightarrow j_2$  presented in the Table 2. Their spin content is the same as that of the lww's given in the first column of the Table 1 in [12] with units in the first and fourth digits. Multiple application of the bosonic generators (28e) and (33e) to obtained lww's produces basis vectors of the infinite-dimensional  $su(2, 2) \oplus su(4)$  irreps. Respective fields on  $AdS_5 \times S^5$ , coordinatized by  $y$  and  $z$ , are listed in the utmost right column of the Tables 1 and 2. To make contact with the consideration of Ref. [12] we use similar notation for these fields. In particular, fields with the same spins of  $su(2) \oplus su(2) \subset su(2, 2)$  are denoted by the same letters and superscripts with numbers in round brackets are used to label them in order of increasing  $AdS_5$  energy. In the subscripts numbers in round brackets following  $sl(2, \mathbb{C})$  or  $su(2)$  indices denote a group of symmetrized indices equal to that number.

There remains to consider simultaneous application of  $Q^{(+)\alpha}_a$  and  $Q^{(+)\alpha}_\alpha$  to the ground state. To find respective lww's one has to decompose the products of oscillators into irreps of four  $su(2)$  subalgebras of  $psu(2, 2|4)$  and leave, according to the criterion formulated in [41], only traceless tensors of the two  $su(2)$  color subalgebras. The role of the latter requirement can be explained on the simple example of the product  $Q^{(+)\alpha}_a Q^{(+)\beta}_b$ . There arise four contributions

$$Q^{(+)\alpha}_a Q^{(+)\beta}_b = a_i^\alpha b_j^\beta \rho_a^i \alpha_j^b + \omega_p^\alpha \varphi_p^q d_a^p c_q^b + a_i^\alpha \varphi_p^q \rho_a^i c_q^b - \omega_p^\alpha b_j^\beta d_a^p \alpha_j^b. \tag{43}$$

It is easy to see that neither  $a_i^\alpha b_j^\beta |0\rangle$  and  $\omega_p^\alpha \varphi_p^q |0\rangle$  nor their traces in the color indices form  $su(2, 2)$  lww's since the r.h.s. of the commutators of  $a_i^\alpha b_j^\beta$  and  $\omega_p^\alpha \varphi_p^q$  with the lowering generators (28d) do not annihilate the oscillator vacuum. However, color-traceless combinations  $(a_i^\alpha b_j^\beta - 1/2\delta_i^j a_k^\alpha b_k^\beta) |0\rangle$  and  $(\omega_p^\alpha \varphi_p^q - 1/2\delta_p^q \omega_r^\alpha \varphi_r^q) |0\rangle$  indeed are the lww's, as well as  $a_i^\alpha \varphi_p^q |0\rangle$  and  $\omega_p^\alpha b_j^\beta |0\rangle$  that ab initio transform irreducibly under both color subalgebras. Note that due to the color symmetry invariance of the  $su(2, 2) \oplus su(4)$  lww's, taking color-traceless combinations of the products of oscillators associated with  $su(2, 2)$  selects color-traceless combinations of the products of oscillators associated with  $su(4)$ . In fact there is a relationship between the  $su(2) \oplus su(2) \subset su(2, 2)$  representations,  $su(2) \oplus su(2) \subset su(4)$  representations and those of the color symmetry carried by a lww.<sup>7</sup> So  $(\beta_a^i \alpha_j^b - 1/2\delta_j^i \beta_a^k \alpha_k^b) |0\rangle$  and  $(d_a^p c_q^b - 1/2\delta_q^p d_a^r c_r^b) |0\rangle$  form  $su(4)$  lww's together with  $\beta_a^i c_q^b |0\rangle$  and  $d_a^p \alpha_j^b |0\rangle$ . Respective spin values  $s_1 = s_2 = 1/2$  and  $j_1 = j_2 = 1/2$  are given in the first row of the Table 3. Examining other products of the supercharges yields extra 24 lww's, whose labels can formally be obtained by summing those of all pairs of the lww's from Tables 1 and 2, except for the vacuum.

We conclude this section with a few comments regarding comparison of the results obtained in this note with known descriptions of the compactification spectrum of IIB supergravity on  $AdS_5 \times S^5$  superbackground. In distinction to the approach of Ref. [12] superparticle model in the supertwistor formulation has finitely many physical states. This can be explained by the fact that in [12] the spectrum of IIB supergravity on  $AdS_5 \times S^5$  is described in terms of the  $S^5$  Fourier modes of the on-shell  $10d$  fields, that is the fields on  $AdS_5$  transforming according to finite-dimensional  $su(4)$  uirs. Fourier modes presented in each line of the Table 1 in [12] have the same  $su(2) \oplus su(2) \subset su(2, 2)$  spins and are parametrized by integer  $p$  with  $AdS_5$  energies and  $su(4)$  labels proportional to  $p$ . The states of the superparticle model collected in Tables 1-3 (apart from the leading mode of  $\varphi^{(1)}$  that corresponds to the scalar field carrying the  $su(2, 2)$  representation) are naturally identified with the  $AdS_5 \times S^5$  fields obtained by summing Fourier modes in each line of the Table 1 in [12]. On the other hand, Fourier modes with the same  $p$  in different lines make up supermultiplets of the  $psu(2, 2|4)$  supersymmetry.

<sup>7</sup> In the non-supersymmetric case color symmetry is not essential in constructing  $su(2, 2)$  lww's (see, e.g. sections 5-6 in Ref. [34]).

**Table 3**  
 $su(2, 2)$  and  $su(4)$  labels of lmv's obtained by acting with the powers of  $Q^{(+)\alpha_a}$  and  $Q^{(+)\dot{a}_\alpha}$  on  $|0\rangle$  and respective fields on  $AdS_5 \times S^5$ .

$su(2, 2)$ labels $(E, s_1, s_2)$	$su(4)$ labels $(C, j_1, j_2)$	$Y$	fields on $AdS_5 \times S^5$
(1, 1/2, 1/2)	(-1, 1/2, 1/2)	0	$A_{\alpha\dot{\alpha}\dot{a}a}^{(1)}(y, z)$
(3/2, 1, 1/2)	(-3/2, 1/2, 0)	-1	$\Psi_{\alpha(2)\dot{a}a}^{(1)}(y, z)$
(3/2, 0, 1/2)	(-3/2, 1/2, 1)	-1	$\lambda_{\dot{\alpha}\dot{a}a(2)}^{(3)}(y, z)$
(3/2, 1/2, 1)	(-3/2, 0, 1/2)	1	$\Psi_{\alpha\dot{a}(2)\dot{a}}^{(1)}(y, z)$
(3/2, 1/2, 0)	(-3/2, 1, 1/2)	1	$\lambda_{\alpha\dot{a}(2)\dot{a}}^{(3)}(y, z)$
(2, 1, 0)	(-2, 1, 0)	0	$A_{\alpha(2)a(2)}^{(2)}(y, z)$
(2, 0, 1)	(-2, 0, 1)	0	$A_{\dot{\alpha}(2)\dot{a}(2)}^{(2)}(y, z)$
(2, 1, 1)	(-2, 0, 0)	0	$h_{\alpha(2)\dot{a}(2)}(y, z)$
(2, 0, 0)	(-2, 1, 1)	0	$\varphi_{\dot{a}(2)\dot{a}(2)}^{(4)}(y, z)$
(2, 1/2, 1/2)	(-2, 1/2, 1/2)	-2	$A_{\alpha\dot{\alpha}\dot{a}a}^{(2)}(y, z)$
(2, 1/2, 1/2)	(-2, 1/2, 1/2)	2	$\tilde{A}_{\alpha\dot{\alpha}\dot{a}a}^{(2)}(y, z)$
(5/2, 1/2, 1)	(-5/2, 0, 1/2)	-1	$\Psi_{\alpha\dot{a}(2)\dot{a}}^{(2)}(y, z)$
(5/2, 1/2, 0)	(-5/2, 1, 1/2)	-1	$\lambda_{\alpha\dot{a}(2)\dot{a}}^{(5)}(y, z)$
(5/2, 1, 1/2)	(-5/2, 1/2, 0)	1	$\Psi_{\alpha(2)\dot{a}a}^{(2)}(y, z)$
(5/2, 0, 1/2)	(-5/2, 1/2, 1)	1	$\lambda_{\dot{\alpha}\dot{a}a(2)}^{(5)}(y, z)$
(5/2, 0, 1/2)	(-5/2, 1/2, 0)	-3	$\lambda_{\dot{\alpha}\dot{a}}^{(4)}(y, z)$
(5/2, 1/2, 0)	(-5/2, 0, 1/2)	3	$\lambda_{\alpha\dot{a}}^{(4)}(y, z)$
(3, 1/2, 1/2)	(-3, 1/2, 1/2)	0	$A_{\alpha\dot{\alpha}\dot{a}a}^{(3)}(y, z)$
(3, 0, 1)	(-3, 0, 0)	-2	$A_{\dot{\alpha}(2)}^{(3)}(y, z)$
(3, 0, 0)	(-3, 1, 0)	-2	$\varphi_{\dot{a}(2)}^{(5)}(y, z)$
(3, 1, 0)	(-3, 0, 0)	2	$A_{\alpha(2)}^{(3)}(y, z)$
(3, 0, 0)	(-3, 0, 1)	2	$\varphi_{\dot{a}(2)}^{(5)}(y, z)$
(7/2, 0, 1/2)	(-7/2, 1/2, 0)	-1	$\lambda_{\dot{\alpha}\dot{a}}^{(6)}(y, z)$
(7/2, 1/2, 0)	(-7/2, 0, 1/2)	1	$\lambda_{\alpha\dot{a}}^{(6)}(y, z)$
(4, 0, 0)	(-4, 0, 0)	0	$\varphi^{(6)}(y, z)$

In particular, leading modes in the  $S^5$  harmonic expansion of  $\lambda_{\alpha\dot{a}}^{(1)}, \lambda_{\dot{\alpha}a}^{(1)}, A_{\alpha(2)}^{(1)}, A_{\dot{\alpha}(2)}^{(1)}$  and subleading mode of  $\varphi^{(1)}$  make up  $N = 4$  super-Yang-Mills multiplet that corresponds to modes with  $p = 1$  in Table 1 of [12]. This multiplet is known to decouple from other modes given by the  $D = 5$   $N = 8$  gauged supergravity multiplet ( $p = 2$ ) and massive  $psu(2, 2|4)$  supermultiplets with spins ranging from 0 to 2 ( $p > 2$ ).

$AdS_5$  energies of some of the superparticle's states in Tables 1-3 with spins  $s_1$  and  $s_2$  lie below the unitarity bound for massless fields on  $AdS_5$   $E \geq s_1 + s_2 + 2$  ( $s_1 s_2 \neq 0$ ) and the bound for massless fields on the Minkowski boundary of  $AdS_5$   $E \geq s_1 + s_2 + 1$  ( $s_1 s_2 = 0$ )<sup>8</sup> and hence correspond to non-unitary representations with non-negative energy. To explain the possible origin of such irreps take, for instance, fields  $\varphi^{(1)}$  in the first row of the Table 1 in [12]. Respective  $su(2, 2)$  uirs have energies  $E_p^{\varphi^{(1)}} = p$  (in our normalization) and their sum  $\sum_{p=1}^{\infty} E_p^{\varphi^{(1)}}$  diverges but can be  $\zeta$ -regularized to take finite value of  $-1/12$ . Also the description of the  $su(4)$  irreps by means of ever increasing number of fermionic oscillators in the limit results in the infinite-dimensional non-unitary irreps that can be realized in terms of finitely many bosonic oscillators. Making these arguments more rigorous requires further study.

Finally let us confront the results of the present work with those of Ref. [13], where the spectrum of IIB supergravity on  $AdS_5 \times S^5$  superbackground had been obtained starting from the field equations in  $D = 1 + 9$  dimensions expanded over the background values of the metric and Ramond-Ramond (RR) self-dual five-form and linearized in small excitations. This allowed not only to find the set of fields on  $AdS_5$  but also the equations they satisfy and to trace their  $10d$  origin. So Table 4 below serves twofold purpose – to match the fields that arise in the spectrum of physical states of the considered superparticle model with those found in [13] and to establish their relation to the fields from the  $D = 1 + 9$  supermultiplet of IIB supergravity. The data presented require some explanatory comments. In distinction to our notation, in [13] small Greek letters are used for curved-space indices of  $AdS_5$  ( $\mu, \nu = 0, \dots, 4$ ) and  $S^5$  ( $\alpha, \beta = 1, \dots, 5$ ), whereas indices of four-components spinors of  $Spin(1, 4)$  and  $Spin(5)$  are not shown explicitly. Bosonic fields that appear twice in the third column, such as  $\varphi^{(1)}$  and  $\varphi^{(6)}$ , stand for eigenvectors of the  $2 \times 2$  matrices that enter the systems of coupled equations for the fields in the second column, like  $S^5$  Fourier modes of  $h_{\alpha}^{\alpha}$  and  $a_{\alpha\beta\gamma\delta}$ . Homogeneous equations for  $A_{\mu\nu}$  factorize so that there are two solutions and  $S^5$  expansion of  $A_{\alpha\beta}$  includes harmonics of two types that are eigenfunctions of the 'square root' of the  $S^5$  Laplacian with different eigenvalues also resulting in two independent solutions. Expansions of the fermionic fields include spinor (labeled by L) and vector-spinor

<sup>8</sup> Unitarity conditions found by G. Mack [36] for the  $SU(2, 2)$  positive energy irreps are the special cases of the unitarity bounds for the positive energy irreps of  $so(2, d)$  [44] and for the positive energy irreps of  $su(p, q)$  [38].



**Table 4**  
Correspondence between the fields of IIB supergravity and the spectrum of its compactification on  $AdS_5 \times S^5$  superbackground.

Bosonic fields		Fermionic fields			
10d fields	$AdS_5 \times S^5$ fields		10d fields	$AdS_5 \times S^5$ fields	
	Ref. [13]	ours		Ref. [13]	ours
graviton	$h'_{\mu\nu}$	$h_{\alpha(2)\dot{\alpha}(2)}$	$\Psi_{\alpha(2)\dot{\alpha}a}^{(1)}$	$\Psi_{\alpha(2)\dot{\alpha}a}^{(1)}$	
	$h_{\mu\alpha}$	$A_{\alpha\dot{\alpha}a\dot{a}}^{(1)}$ $A_{\alpha\dot{\alpha}a\dot{a}}^{(3)}$			$\Psi_{\alpha\dot{\alpha}(2)\dot{a}}^{(1)}$
	$h_{(\alpha\beta)}$	$\varphi_{a(2)\dot{a}(2)}^{(4)}$	$\Psi_{\alpha(2)\dot{\alpha}a}^{(2)}$	$\Psi_{\alpha\dot{\alpha}(2)\dot{a}}^{(2)}$	
complex scalar	$h_{\alpha}^{\alpha}$	$\varphi^{(1)}$ $\varphi^{(6)}$	complex gravitino	$\lambda_{\alpha\dot{a}}^{(1)}$	
	$B$	$\varphi^{(3)}, \bar{\varphi}^{(3)}$		$\Psi^{L-}$	$\lambda_{\dot{a}a}^{(1)}$
complex 2-form	$A_{\mu\nu}$	$A_{\alpha(2)}^{(1)}, A_{\dot{\alpha}(2)}^{(1)}$ $A_{\alpha(2)}^{(3)}, A_{\dot{\alpha}(2)}^{(3)}$	$\Psi^{L+}$	$\lambda_{\alpha\dot{a}}^{(6)}$ $\lambda_{\dot{a}a}^{(6)}$	
	$A_{\mu\alpha}$	$A_{\alpha\dot{\alpha}a\dot{a}}^{(2)}, \bar{A}_{\alpha\dot{\alpha}a\dot{a}}^{(2)}$	$\Psi^{T-}$	$\lambda_{\alpha a(2)\dot{a}}^{(3)}$ $\lambda_{\dot{a}a\dot{a}(2)}^{(3)}$	
	$A_{\alpha\beta}$	$\varphi_{a(2)}^{(2)}, \varphi_{\dot{a}(2)}^{(2)}$ $\varphi_{a(2)}^{(5)}, \varphi_{\dot{a}(2)}^{(5)}$	$\Psi^{T+}$	$\lambda_{\alpha a(2)\dot{a}}^{(5)}$ $\lambda_{\dot{a}a\dot{a}(2)}^{(5)}$	
RR 4-form	$a_{\mu\nu\alpha\beta}$	$A_{\alpha(2)a(2)}^{(2)}, A_{\dot{\alpha}(2)\dot{a}(2)}^{(2)}$	complex spinor	$\lambda^{L-}$	$\lambda_{\alpha\dot{a}}^{(2)}$ $\lambda_{\dot{a}a}^{(2)}$
	$a_{\mu\alpha\beta\gamma}$	$A_{\alpha\dot{\alpha}a\dot{a}}^{(1)}$ $A_{\alpha\dot{\alpha}a\dot{a}}^{(3)}$		$\lambda^{L+}$	$\lambda_{\alpha\dot{a}}^{(4)}$ $\lambda_{\dot{a}a}^{(4)}$
	$a_{\alpha\beta\gamma\delta}$	$\varphi^{(1)}$ $\varphi^{(6)}$			

(labeled by T) harmonics that are eigenspinors of the Dirac operator on  $S^5$  with opposite eigenvalues so expansion modes are endowed + and – superscripts. The presence of spinor and vector-spinor harmonics with different eigenvalues results in two independent solutions with different ‘masses’ ( $AdS_5$  energies) of the equations for fermionic fields of each type.

### 6. Discussion

In this note we have shown how physical states of the quantized  $AdS_5 \times S^5$  massless superparticle in the twistor formulation can be described in terms of the  $su(2)$  bosonic and fermionic oscillators some of which were used previously to construct unitary supermultiplets of  $psu(2, 2|4)$  [12], [34], [35]. These physical states of the superparticle have been mapped to those in the spectrum of fluctuations of IIB supergravity on  $AdS_5 \times S^5$  superbackground [12], [13] and  $N = 4$  super-Yang-Mills multiplet on  $D = 4$  Minkowski boundary of  $AdS_5$ . We observed that the description of IIB supergravity spectrum on  $AdS_5 \times S^5$  superbackground in the context of  $D = 10$  superparticle model differs from the original one in Ref. [12], where it was presented in the form of an infinite sum of unitary  $psu(2, 2|4)$  supermultiplets,  $l_{wv}$ 's of which were obtained by acting on the oscillator vacuum with raising supersymmetry generators of the  $psu(2, 2|4)$  superalgebra constructed out of  $P \geq 1$  copies of the four sets of  $su(2)$  oscillators coming from the linear combinations of the components of  $SU(2, 2|4)$  Ferber supertwistors. In these supermultiplets fields corresponding to the irreps that have the same  $su(2) \oplus su(2) \subset su(2, 2)$  spins and  $Y$  eigenvalues but transform according to different representations of  $su(4)$  represent expansion coefficients of the 10d fields over  $S^5$  harmonics [13]. On the contrary, dynamical variables of the  $AdS_5 \times S^5$  superparticle model in the four-supertwistor formulation are two Ferber or  $c$ -type supertwistors and two  $a$ -type supertwistors, whose components have another Grassmann parity. The latter give rise to  $su(2)$  oscillators some of which satisfy wrong-sign (anti)commutation relations. As a result for some states their  $su(2, 2)$   $l_{wv}$ 's have  $AdS_5$  energies, though non-negative but lying below the unitarity bounds for respective values of spins and hence corresponding to non-unitary irreps. We conjectured that such irreps correspond to infinite sums of the unitary ones. There also arise infinite-dimensional non-unitary  $su(4)$  representations that can be interpreted as resulting from summing up the  $S^5$  Fourier modes.

It would be interesting to study quantization of the superparticle model directly in terms of the supertwistors. In [19] there were derived the first-order differential equations for the superparticle's wave function, whose arguments are  $c$ - and  $a$ -type superambitwistors. To provide the supertwistor description of the IIB supergravity fluctuations over  $AdS_5 \times S^5$  superbackground as solution of these equations it is necessary to develop ambitwistor transform of respective fields on  $AdS_5$ . This would be an important step towards the twistor formulation of the  $AdS_5/CFT_4$  duality.

The approach reported here can also be extended to describe fluctuation spectra over other highly supersymmetric backgrounds of known supergravity theories in the framework of appropriate supertwistor formulations of the models of superparticles and extended objects such as superstrings and branes.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The author is grateful to A.A. Zheltukhin for interesting discussions. This work was partially supported by the Ministry of Education and Science of Ukraine (research grant No. 0120U102252).

## References

- [1] H. Elvang, Yu-tin Huang, *Scattering Amplitudes in Gauge Theory and Gravity*, Cambridge University Press, 2015, arXiv:1308.1697 [hep-th].
- [2] E. Witten, Perturbative gauge theory as a string theory in twistor space, *Commun. Math. Phys.* 252 (2004) 189, arXiv:hep-th/0312171.
- [3] S.J. Parke, T.R. Taylor, Amplitude for  $n$ -gluon scattering, *Phys. Rev. Lett.* 56 (1986) 2459.
- [4] V.P. Nair, A current algebra for some gauge theory amplitudes, *Phys. Lett. B* 214 (1988) 215.
- [5] L.J. Mason, Twistor actions for non-self-dual fields: a new foundation for twistor-string theory, *J. High Energy Phys.* 0510 (2005) 009, arXiv:hep-th/0507269.
- [6] L.J. Mason, D. Skinner, An ambitwistor Yang-Mills Lagrangian, *Phys. Lett. B* 636 (2006) 60, arXiv:hep-th/0510262.
- [7] R. Boels, L. Mason, D. Skinner, Supersymmetric gauge theories in twistor space, *J. High Energy Phys.* 0702 (2007) 014, arXiv:hep-th/0604040.
- [8] W. Siegel, AdS/CFT in superspace, arXiv:1005.2317 [hep-th];  
W. Siegel, Embedding versus 6D twistors, arXiv:1204.5679 [hep-th].
- [9] T. Adamo, D. Skinner, J. Williams, Twistor methods for  $AdS_5$ , *J. High Energy Phys.* 1608 (2016) 167, arXiv:1607.03763 [hep-th].
- [10] R.R. Metsaev, A.A. Tseytlin, Type IIB superstring action in  $AdS_5 \times S^5$  background, *Nucl. Phys. B* 533 (1998) 109, arXiv:hep-th/9805028.
- [11] R. Kallosh, J. Rahmfeld, A. Rajaraman, Near horizon superspace, *J. High Energy Phys.* 9809 (1998) 002, arXiv:hep-th/9805217.
- [12] M. Gunaydin, N. Marcus, The spectrum of the  $S^5$  compactification of the chiral  $N=2$ ,  $D=10$  supergravity and the unitary supermultiplets of  $U(2, 2|4)$ , *Class. Quantum Gravity* 2 (1985) L11.
- [13] H.J. Kim, L.J. Romans, P. van Nieuwenhuizen, Mass spectrum of chiral ten-dimensional  $N=2$  supergravity on  $S^5$ , *Phys. Rev. D* 32 (1985) 389.
- [14] I. Bars, Twistors and 2T-physics, *AIP Conf. Proc.* 767 (2005) 3, arXiv:hep-th/0502065;  
I. Bars, Lectures on twistors, *Bulg. J. Phys.* 33 (S1) (2006) 487, arXiv:hep-th/0601091.
- [15] I. Bars, Twistor superstring in 2T-physics, *Phys. Rev. D* 70 (2004) 104022, arXiv:hep-th/0407239.
- [16] I. Bars, 2T physics formulation of superconformal dynamics relating to twistors and supertwistors, *Phys. Lett. B* 483 (2000) 248, arXiv:hep-th/0004090.
- [17] A. Ferber, Supertwistors and conformal supersymmetry, *Nucl. Phys. B* 132 (1978) 55.
- [18] D.V. Uvarov, Supertwistor formulation for massless superparticle in  $AdS_5 \times S^5$  superspace, *Nucl. Phys. B* 936 (2019) 690, arXiv:1807.08318 [hep-th].
- [19] D.V. Uvarov, Multitwistor mechanics of massless superparticle on  $AdS_5 \times S^5$  superbackground, *Nucl. Phys. B* 950 (2020) 114830, arXiv:1907.13613 [hep-th].
- [20] R.R. Metsaev, C.B. Thorn, A.A. Tseytlin, Light cone superstring in AdS space-time, *Nucl. Phys. B* 596 (2001) 151, arXiv:hep-th/0009171.
- [21] I. Bars, Hidden twelve-dimensional structures in  $AdS_5 \times S^5$  and  $M^4 \times R^6$  supergravities, *Phys. Rev. D* 66 (2002) 105024, arXiv:hep-th/0208012.
- [22] T. Horigane, Y. Kazama, Exact quantization of a superparticle in  $AdS_5 \times S^5$ , *Phys. Rev. D* 81 (2010) 045004, arXiv:0912.1166 [hep-th].
- [23] W. Siegel, Spacecone quantization of AdS superparticle, arXiv:1005.5049 [hep-th].
- [24] M. Heinze, Spectrum and quantum symmetries of the  $AdS_5 \times S^5$  superstring, arXiv:1507.03005 [hep-th].
- [25] P. Claus, M. Gunaydin, R. Kallosh, J. Rahmfeld, Y. Zunger, Supertwistors as quarks of  $SU(2, 2|4)$ , *J. High Energy Phys.* 9905 (1999) 019, arXiv:hep-th/9905112.
- [26] I. Bars, M. Gunaydin, Unitary representations of non-compact supergroups, *Commun. Math. Phys.* 91 (1983) 31.
- [27] P. Claus, J. Rahmfeld, Y. Zunger, A simple particle action from a twistor parametrization of  $AdS_5$ , *Phys. Lett. B* 466 (1999) 181, arXiv:hep-th/9906118.
- [28] A.S. Arvanitakis, A.E. Barns-Graham, P.K. Townsend, Twistor variables for anti-de Sitter (super)particles, *Phys. Rev. Lett.* 118 (2017) 141601, arXiv:1608.04380 [hep-th];  
A.S. Arvanitakis, A.E. Barns-Graham, P.K. Townsend, Twistor description of spinning particles in AdS, *J. High Energy Phys.* 1801 (2018) 059, arXiv:1710.09557 [hep-th].
- [29] P. Claus, R. Kallosh, J. Rahmfeld, BRST quantization of a particle in  $AdS_5$ , *Phys. Lett. B* 462 (1999) 285, arXiv:hep-th/9906195.
- [30] Y. Nambu, Relativistic wave equations for particles with internal structure and mass spectrum, *Prog. Theor. Phys. Suppl.* 37–38 (1966) 368.
- [31] A.O. Barut, H. Kleinert, Current operators and Majorana equation for the hydrogen atom from dynamical groups, *Phys. Rev.* 157 (1967) 1180.
- [32] G. Mack, I. Todorov, Irreducibility of the ladder representations of  $U(2, 2)$  when restricted to the Poincare subgroup, *J. Math. Phys.* 10 (1969) 2078.
- [33] M. Gunaydin, C. Sacliglu, Oscillator-like unitary representations of non-compact groups with a Jordan structure and the non-compact groups of supergravity, *Commun. Math. Phys.* 87 (1982) 159.
- [34] M. Gunaydin, D. Minic, M. Zagermann, 4d doubleton conformal theories, CPT and IIB string on  $AdS_5 \times S^5$ , *Nucl. Phys. B* 534 (1998) 96, arXiv:hep-th/9806042.
- [35] M. Gunaydin, D. Minic, M. Zagermann, Novel supermultiplets of  $SU(2, 2|4)$  and the  $AdS_5/CFT_4$  duality, *Nucl. Phys. B* 544 (1999) 737, arXiv:hep-th/9810226.
- [36] G. Mack, All unitary ray representations of the conformal group  $SU(2, 2)$  with positive energy, *Commun. Math. Phys.* 55 (1977) 1.
- [37] V.K. Dobrev, Positive energy representations, holomorphic discrete series and finite-dimensional irreps, *J. Phys. A, Math. Theor.* 41 (2008) 425206, arXiv:0712.4375 [hep-th].
- [38] M. Gunaydin, D. Volin, The complete unitary dual of non-compact Lie superalgebra  $su(p, q|m)$  via the generalized oscillator formalism, and non-compact Young diagrams, *Commun. Math. Phys.* 367 (2019) 873, arXiv:1712.01811 [hep-th].
- [39] V.K. Dobrev, V.B. Petkova, All positive energy unitary irreducible representations of extended conformal supersymmetry, *Phys. Lett. B* 162 (1985) 127.
- [40] D.V. Uvarov, Ambitwistors, oscillators and massless fields on  $AdS_5$ , *Phys. Lett. B* 762 (2016) 415, arXiv:1607.05233 [hep-th].
- [41] I. Bars, A mysterious zero in  $AdS_5 \times S^5$  supergravity, *Phys. Rev. D* 66 (2002) 105023, arXiv:hep-th/0205194.
- [42] C. Marboe, D. Volin, The full spectrum of  $AdS_5/CFT_4$  I: representation theory and one-loop  $Q$ -system, *J. Phys. A* 51 (2018) 165401, arXiv:1701.03704 [hep-th].
- [43] T. Basile, Euihun Joung, K. Mkrtchyan, M. Mojaza, Dual pair correspondence in physics: oscillator realizations and representations, *J. High Energy Phys.* 2009 (2020) 020, arXiv:2006.07102 [hep-th].
- [44] R. Metsaev, Lowest eigenvalues of the energy operator for totally (anti)symmetric massless fields of the  $n$ -dimensional anti-de Sitter group, *Class. Quantum Gravity* 11 (1994) L141;  
R. Metsaev, Massless mixed symmetry bosonic free fields in  $d$ -dimensional anti-de Sitter space-time, *Phys. Lett. B* 354 (1995) 78.