



Determination of γ and $-2\beta_s$ from charmless two-body decays of beauty mesons



LHCb Collaboration

ARTICLE INFO

Article history:

Received 19 August 2014
 Received in revised form 27 November 2014
 Accepted 5 December 2014
 Available online 9 December 2014
 Editor: L. Rolandi

ABSTRACT

Using the latest LHCb measurements of time-dependent CP violation in the $B_s^0 \rightarrow K^+K^-$ decay, a U-spin relation between the decay amplitudes of $B_s^0 \rightarrow K^+K^-$ and $B^0 \rightarrow \pi^+\pi^-$ decay processes allows constraints to be placed on the angle γ of the unitarity triangle and on the B_s^0 mixing phase $-2\beta_s$. Results from an extended approach, which uses additional inputs on $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays from other experiments and exploits isospin symmetry, are also presented. The dependence of the results on the maximum allowed amount of U-spin breaking is studied. At 68% probability, the value $\gamma = (63.5_{-6.7}^{+7.2})^\circ$ modulo 180° is determined. In an alternative analysis, the value $-2\beta_s = -0.12_{-0.16}^{+0.14}$ rad is found. In both measurements, the uncertainties due to U-spin breaking effects up to 50% are included.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Introduction

The understanding of flavour dynamics is one of the most important aims of particle physics. Charge–parity (CP) violation and rare decay processes involving weak decays of B mesons provide tests of the Cabibbo–Kobayashi–Maskawa (CKM) mechanism [1, 2] in the Standard Model (SM). The CKM matrix describes all flavour changing transitions of quarks in the SM. These include tree-level decays, which are expected to be largely unaffected by non-SM contributions, and flavour changing neutral current transitions characterized by the presence of loops in the relevant diagrams, which are sensitive to the presence of non-SM physics. Tests of the CKM matrix structure, commonly represented by the unitarity triangle (UT), are of fundamental importance.

Although significant hadronic uncertainties usually complicate the experimental determination of the CKM matrix elements V_{ij} , there are certain cases where the V_{ij} can be derived with reduced or even negligible hadronic uncertainty. One of these cases involves the determination of the UT angle γ . The angle γ , defined as $\arg[-(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*)]$, can be measured using decays that involve tree diagrams only, with almost vanishing theoretical uncertainty [3]. However, γ is experimentally the least known of the UT angles. World averages of the measurements performed by BaBar, Belle and LHCb [4–7], provided by the UTfit Collaboration and CKMfitter group, are $\gamma = (70.1 \pm 7.1)^\circ$ and $\gamma = (68.0_{-8.5}^{+8.0})^\circ$, respectively¹ [8,9].

An alternative strategy to determine γ using two-body charmless B decays, namely $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$, has also been proposed [10–12]. Knowledge of the B^0 mixing phase 2β , where $\beta = \arg[-(V_{cd}V_{cb}^*)/(V_{td}V_{tb}^*)]$, is needed as an input. Due to the presence of penguin diagrams in the decay amplitudes, in addition to tree diagrams, the interpretation of the observables requires knowledge of hadronic factors that cannot at present be calculated accurately from quantum chromodynamics (QCD). However, the hadronic parameters entering the $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays are related by the U-spin symmetry of strong interactions. This symmetry, related to the exchange of d and s quarks in the decay diagrams, can be exploited to determine the unknown hadronic factors. A more sophisticated analysis has also been proposed [13], where it is suggested to combine the U-spin analysis of $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays with the isospin analysis of $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays [14], in order to achieve a more robust determination of γ with respect to U-spin breaking effects. The B_s^0 mixing phase $-2\beta_s$, where $\beta_s = \arg[-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)]$, can also be determined with either analysis approach.

An analysis based on Bayesian statistics, aimed at determining probability density functions (PDFs) for γ and $-2\beta_s$, is presented in this Letter. This uses the latest LHCb measurements of time-dependent CP violation in the $B_s^0 \rightarrow K^+K^-$ decay, exploiting U-spin symmetry with the $B^0 \rightarrow \pi^+\pi^-$ decay. An extended analysis, including measurements on $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays from other experiments, is also performed. The Letter is organized as follows. First, the theoretical formalism needed to describe CP violation is introduced in Section 2, including the SM parameterization of the decay amplitudes of the various decays.

¹ The measurements of γ are given modulo 180° throughout this Letter.

The experimental status is given in Section 3. In Section 4 we present the determination of γ and $-2\beta_s$ using $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays, and in Section 5 we also add information from $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays. The dependence of the measurements of γ and $-2\beta_s$ on the amount of U-spin breaking is studied in detail in both cases. Finally, conclusions are drawn in Section 6.

2. Theoretical formalism

Assuming *CPT* invariance, the *CP* asymmetry as a function of decay time for a neutral B^0 or B_s^0 meson decaying to a self-conjugate final state f , with $f = \pi^+\pi^-$, $\pi^0\pi^0$ or K^+K^- , is given by

$$\begin{aligned} \mathcal{A}(t) &\equiv \frac{\Gamma_{\bar{B}_{(s)}^0 \rightarrow f}(t) - \Gamma_{B_{(s)}^0 \rightarrow f}(t)}{\Gamma_{\bar{B}_{(s)}^0 \rightarrow f}(t) + \Gamma_{B_{(s)}^0 \rightarrow f}(t)} \\ &= \frac{-C_f \cos(\Delta m_{d(s)}t) + S_f \sin(\Delta m_{d(s)}t)}{\cosh\left(\frac{\Delta\Gamma_{d(s)}}{2}t\right) + A_f^{\Delta\Gamma} \sinh\left(\frac{\Delta\Gamma_{d(s)}}{2}t\right)}, \end{aligned} \quad (1)$$

where $\Delta m_{d(s)} \equiv m_{d(s),H} - m_{d(s),L}$ and $\Delta\Gamma_{d(s)} \equiv \Gamma_{d(s),L} - \Gamma_{d(s),H}$ are the mass and width differences of the $B_{(s)}^0 - \bar{B}_{(s)}^0$ system mass eigenstates. The subscripts H and L denote the heavy and light eigenstates. With this convention, the value of $\Delta m_{d(s)}$ is positive by definition, and that of $\Delta\Gamma_s$ is measured to be positive [15], $\Delta\Gamma_s = 0.106 \pm 0.011(\text{stat}) \pm 0.007(\text{syst}) \text{ ps}^{-1}$ [16]. The value of $\Delta\Gamma_d$ is also positive in the SM and is expected to be much smaller than that of $\Delta\Gamma_s$, $\Delta\Gamma_d \simeq 3 \times 10^{-3} \text{ ps}^{-1}$ [8]. The quantities C_f , S_f and $A_f^{\Delta\Gamma}$ are

$$\begin{aligned} C_f &\equiv \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}, \\ S_f &\equiv \frac{2\text{Im}\lambda_f}{1 + |\lambda_f|^2} \quad \text{and} \quad A_f^{\Delta\Gamma} \equiv -\frac{2\text{Re}\lambda_f}{1 + |\lambda_f|^2}, \end{aligned} \quad (2)$$

where λ_f is given by

$$\lambda_f \equiv \frac{q}{p} \frac{\bar{A}_f}{A_f}. \quad (3)$$

The two mass eigenstates of the effective Hamiltonian in the $B_{(s)}^0 - \bar{B}_{(s)}^0$ system are $p|B_{(s)}^0\rangle \pm q|\bar{B}_{(s)}^0\rangle$, where p and q are complex parameters satisfying the relation $|p|^2 + |q|^2 = 1$. The parameter λ_f is thus related to $B_{(s)}^0 - \bar{B}_{(s)}^0$ mixing (via q/p) and to the decay amplitudes of the $B_{(s)}^0 \rightarrow f$ decay (A_f) and of the $\bar{B}_{(s)}^0 \rightarrow f$ decay (\bar{A}_f). Assuming negligible *CP* violation in mixing ($|q/p| = 1$), as expected in the SM and supported by current experimental determinations [17,18], the terms C_f and S_f parameterize *CP* violation in the decay and in the interference between mixing and decay, respectively. From the definitions given in Eq. (2), it follows that

$$(C_f)^2 + (S_f)^2 + (A_f^{\Delta\Gamma})^2 = 1. \quad (4)$$

It is then possible to express the magnitude (but not the sign) of $A_f^{\Delta\Gamma}$ as a function of C_f and S_f . There are therefore two independent parameters, which can be chosen, for example, to be $\text{Re}\lambda_f$ and $\text{Im}\lambda_f$, or C_f and S_f . In the latter case, the sign of $A_f^{\Delta\Gamma}$ carries additional information.

The *CP*-averaged branching fraction is given by

$$\mathcal{B}_f = \frac{1}{2} F(B_{(s)}^0 \rightarrow f) (|\bar{A}_f|^2 + |A_f|^2), \quad (5)$$

where

$$F(B^0 \rightarrow \pi^+\pi^-) = \frac{\sqrt{m_{B^0}^2 - 4m_{\pi^+}^2}}{m_{B^0}^2} \tau_{B^0}, \quad (6)$$

$$F(B^0 \rightarrow \pi^0\pi^0) = \frac{\sqrt{m_{B^0}^2 - 4m_{\pi^0}^2}}{m_{B^0}^2} \tau_{B^0}, \quad (7)$$

$$\begin{aligned} F(B_s^0 \rightarrow K^+K^-) \\ = \frac{\sqrt{m_{B_s^0}^2 - 4m_{K^+}^2}}{m_{B_s^0}^2} [2\tau_{B_s^0} - (1 - y_s^2)\tau(B_s^0 \rightarrow K^+K^-)], \end{aligned} \quad (8)$$

with $\tau_{B^0} \equiv 1/\Gamma_d$, $\tau_{B_s^0} \equiv 1/\Gamma_s$ and $y_s \equiv \Delta\Gamma_s/(2\Gamma_s)$. The term m_x is the mass of the meson x , $\Gamma_{d(s)} \equiv (\Gamma_{d(s),L} + \Gamma_{d(s),H})/2$ is the average decay width of the $B_{(s)}^0$ meson, and $\tau(B_s^0 \rightarrow K^+K^-)$ is the effective lifetime measured using $B_s^0 \rightarrow K^+K^-$ decays. The extra term in Eq. (8) follows from the fact that the $\bar{B}_s^0 - B_s^0$ meson system is characterized by a sizeable decay width difference. This leads to a difference between the measured (*i.e.* decay-time-integrated) branching fraction and the theoretical branching fraction, and a correction is applied using the corresponding effective lifetime measurement [19].

In the case of a B^+ meson decaying to a final state f , the *CP* asymmetry is given by

$$\mathcal{A}_f = \frac{|\bar{A}_f|^2 - |A_f|^2}{|\bar{A}_f|^2 + |A_f|^2}, \quad (9)$$

and the *CP*-averaged branching fraction is

$$\mathcal{B}_f = \frac{1}{2} F(B^+ \rightarrow f) (|\bar{A}_f|^2 + |A_f|^2), \quad (10)$$

where

$$F(B^+ \rightarrow \pi^+\pi^0) = \frac{\sqrt{m_{B^+}^2 - (m_{\pi^+} + m_{\pi^0})^2}}{m_{B^+}^2} \tau_{B^+}, \quad (11)$$

with τ_{B^+} the lifetime and m_{B^+} the mass of the B^+ meson.

Adopting the parameterization from Ref. [10] and its extension from Ref. [13], assuming isospin symmetry and neglecting electroweak penguin contributions, the following expressions for the various *CP* asymmetry terms and branching fractions are obtained in the framework of the SM

$$C_{\pi^+\pi^-} = -\frac{2d \sin(\vartheta) \sin(\gamma)}{1 - 2d \cos(\vartheta) \cos(\gamma) + d^2}, \quad (12)$$

$$S_{\pi^+\pi^-} = -\frac{\sin(2\beta + 2\gamma) - 2d \cos(\vartheta) \sin(2\beta + \gamma) + d^2 \sin(2\beta)}{1 - 2d \cos(\vartheta) \cos(\gamma) + d^2}, \quad (13)$$

$$C_{\pi^0\pi^0} = -\frac{2dq \sin(\vartheta_q - \vartheta) \sin(\gamma)}{q^2 + 2dq \cos(\vartheta_q - \vartheta) \cos(\gamma) + d^2}, \quad (14)$$

$$\mathcal{A}_{\pi^+\pi^0} = 0, \quad (15)$$

$$C_{K^+K^-} = \frac{2\tilde{d}' \sin(\vartheta') \sin(\gamma)}{1 + 2\tilde{d}' \cos(\vartheta') \cos(\gamma) + \tilde{d}'^2}, \quad (16)$$

$$\begin{aligned} S_{K^+K^-} = & -\left(\frac{\sin(-2\beta_s + 2\gamma) + 2\tilde{d}' \cos(\vartheta') \sin(-2\beta_s + \gamma)}{1 + 2\tilde{d}' \cos(\vartheta') \cos(\gamma) + \tilde{d}'^2} \right. \\ & \left. + \frac{\tilde{d}'^2 \sin(-2\beta_s)}{1 + 2\tilde{d}' \cos(\vartheta') \cos(\gamma) + \tilde{d}'^2} \right), \end{aligned} \quad (17)$$

$$\mathcal{B}_{\pi^+\pi^-} = F(B^0 \rightarrow \pi^+\pi^-) |D|^2 \times (1 - 2d \cos(\vartheta) \cos(\gamma) + d^2), \quad (18)$$

$$\mathcal{B}_{\pi^0\pi^0} = F(B^0 \rightarrow \pi^0\pi^0) \frac{|D|^2}{2} \times (q^2 + 2dq \cos(\vartheta_q - \vartheta) \cos(\gamma) + d^2), \quad (19)$$

$$\mathcal{B}_{\pi^+\pi^0} = F(B^+ \rightarrow \pi^+\pi^0) \frac{|D|^2}{2} (1 + q^2 + 2q \cos(\vartheta_q)), \quad (20)$$

$$\mathcal{B}_{K^+K^-} = F(B_s^0 \rightarrow K^+K^-) \frac{\lambda^2}{(1 - \lambda^2/2)^2} |D'|^2 \times (1 + 2\tilde{d}' \cos(\vartheta') \cos(\gamma) + \tilde{d}'^2), \quad (21)$$

where $\tilde{d}' \equiv d'(1 - \lambda^2)/\lambda^2$ and $\lambda \equiv |V_{us}|/\sqrt{|V_{ud}|^2 + |V_{us}|^2}$. In addition, $A_{K^+K^-}^{\Delta\Gamma}$ can be expressed as

$$A_{K^+K^-}^{\Delta\Gamma} = - \left(\frac{\cos(-2\beta_s + 2\gamma) + 2\tilde{d}' \cos(\vartheta') \cos(-2\beta_s + \gamma)}{1 + 2\tilde{d}' \cos(\vartheta') \cos(\gamma) + \tilde{d}'^2} + \frac{\tilde{d}'^2 \cos(-2\beta_s)}{1 + 2\tilde{d}' \cos(\vartheta') \cos(\gamma) + \tilde{d}'^2} \right). \quad (22)$$

The quantities $|D|$, d , ϑ , q and ϑ_q are real-valued hadronic parameters related to the decay amplitudes of $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays, whereas $|D'|$, d' and ϑ' are the analogues of $|D|$, d and ϑ for the $B_s^0 \rightarrow K^+K^-$ decay. They are defined as

$$D^{(\prime)} \equiv A\lambda^3 R_u (-T^{(\prime)} - P^{(\prime)u} + P^{(\prime)t}), \quad (23)$$

$$d^{(\prime)} e^{i\vartheta^{(\prime)}} \equiv \frac{1}{R_u} \frac{P^{(\prime)c} - P^{(\prime)t}}{T^{(\prime)} + P^{(\prime)u} - P^{(\prime)t}}, \quad (24)$$

$$qe^{i\vartheta_q} \equiv \frac{C - P^u + P^t}{T + P^u - P^t}, \quad (25)$$

where T and C represent the contributions from $\bar{b} \rightarrow \bar{u}W^+ (\rightarrow \bar{u}\bar{d})$ tree and colour-suppressed tree transitions, P^q represents the contributions from $\bar{b} \rightarrow \bar{d}g (\rightarrow \bar{u}u)$ or $\bar{b} \rightarrow \bar{d}g (\rightarrow \bar{d}d)$ penguin transitions (the index $q \in \{u, c, t\}$ indicates the flavour of the internal quark in the penguin loop), R_u is one of the sides of the UT

$$R_u = \frac{1}{\lambda} \left(1 - \frac{\lambda^2}{2} \right) \left| \frac{V_{ub}}{V_{cb}} \right|, \quad (26)$$

and $A \equiv 1/\lambda |V_{cb}/V_{us}|$. Analogously, T' represents the contribution from $\bar{b} \rightarrow \bar{u}W^+ (\rightarrow \bar{u}\bar{s})$ tree transitions, and P'^q represents the contributions from $\bar{b} \rightarrow \bar{s}g (\rightarrow \bar{u}u)$ penguin transitions.

3. Experimental status

CP violation both in decay amplitudes and in their interference with the B^0 - \bar{B}^0 mixing amplitude has been seen in $B^0 \rightarrow \pi^+\pi^-$ decays by the BaBar [20] and Belle [21] experiments, which also provided measurements of CP violation in the $B^+ \rightarrow \pi^+\pi^0$ [22,23] and $B^0 \rightarrow \pi^0\pi^0$ [20,24] decays. LHCb has recently published measurements of CP violation in $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays [25]. Measurements of branching fractions for $B_s^0 \rightarrow \pi^+\pi^-$, $B^+ \rightarrow \pi^+\pi^0$ and $B^0 \rightarrow \pi^0\pi^0$ decays have been made by BaBar [20,22,26] and Belle [23,24]. CDF and LHCb have also measured the $B^0 \rightarrow \pi^+\pi^-$ branching fraction, as well as that of the $B_s^0 \rightarrow K^+K^-$ decay [27,28], using the world average of the $B^0 \rightarrow K^+\pi^-$ branching fraction for normalization [17]. The current experimental knowledge is summarized in Table 1.

The LHCb measurement of $C_{K^+K^-}$ and $S_{K^+K^-}$ in Ref. [25] was obtained using the constraint

$$A_{K^+K^-}^{\Delta\Gamma} = -\sqrt{1 - (C_{K^+K^-})^2 - (S_{K^+K^-})^2} \quad (27)$$

in the maximum likelihood fit. In the same analysis, the sign of $A_{K^+K^-}^{\Delta\Gamma}$ was verified to be negative, as expected in the SM. A measurement of $A_{K^+K^-}^{\Delta\Gamma}$ has also been made by LHCb via an effective lifetime measurement of the $B_s^0 \rightarrow K^+K^-$ decay, using the same data sample as in Ref. [25], but with different event selection. The result is $A_{K^+K^-}^{\Delta\Gamma} = -0.87 \pm 0.17(\text{stat}) \pm 0.13(\text{syst})$ [29]. In the analysis presented in this Letter, $A_{K^+K^-}^{\Delta\Gamma}$ is constrained to have a negative value.

4. Determination of γ and $-2\beta_s$ from $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays

A method to determine γ and $-2\beta_s$ using CP asymmetries and branching fractions of $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays, exploiting the approximate U-spin symmetry of strong interactions, was proposed in Refs. [10–12]. Typical U-spin breaking corrections are expected to be around the 30% level [30,31]. In the limit of strict U-spin symmetry, one has $d = d'$, $\vartheta = \vartheta'$ and $|D| = |D'|$. As pointed out in Ref. [10], the equalities $d = d'$ and $\vartheta = \vartheta'$ do not receive U-spin breaking corrections within the factorization approximation, in contrast with the equality $|D| = |D'|$,

$$\left| \frac{D'}{D} \right|_{\text{fact}} = \frac{f_K f_{B_s^0 K}^+ (m_K^2) m_{B_s^0}^2 - m_K^2}{f_\pi f_{B^0 \pi}^+ (m_\pi^2) m_{B^0}^2 - m_\pi^2}, \quad (28)$$

where f_K and f_π are the kaon and pion decay constants, and $f_{B_s^0 K}^+(m_K^2)$ and $f_{B^0 \pi}^+(m_\pi^2)$ parameterize hadronic matrix elements. These quantities have been determined using QCD sum rules [32], yielding

$$\left| \frac{D'}{D} \right|_{\text{fact}} = 1.41_{-0.11}^{+0.20}.$$

To take into account non-factorizable U-spin breaking corrections, we parameterize the effect of the breaking as

$$|D'| = \left| \frac{D'}{D} \right|_{\text{fact}} |D| |1 + r_D e^{i\vartheta_{rD}}|, \quad (29)$$

$$d' e^{i\vartheta'} = d e^{i\vartheta} \frac{1 + r_G e^{i\vartheta_{rG}}}{1 + r_D e^{i\vartheta_{rD}}}, \quad (30)$$

where r_D and r_G are relative magnitudes, and ϑ_{rD} and ϑ_{rG} are phase shifts caused by the breaking. In the absence of non-factorizable U-spin breaking, one has $r_D = 0$ and $r_G = 0$.

We perform two distinct analyses, to determine either γ or $-2\beta_s$. They are referred to as analyses A and B, respectively. To improve the precision on the determination of γ , in analysis A the value of $-2\beta_s$ is constrained as

$$-2\beta_s = -2\lambda^2 \bar{\eta} [1 + \lambda^2 (1 - \bar{\rho})], \quad (31)$$

which is valid in the SM up to terms of order λ^4 . The parameters $\bar{\rho}$ and $\bar{\eta}$ determine the apex of the UT, and are defined as $\bar{\rho} + i\bar{\eta} \equiv -(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*)$. Since $\bar{\rho}$ and $\bar{\eta}$ can be written as functions of β and γ as

$$\bar{\rho} = \frac{\sin \beta \cos \gamma}{\sin(\beta + \gamma)}, \quad \bar{\eta} = \frac{\sin \beta \sin \gamma}{\sin(\beta + \gamma)}, \quad (32)$$

we can express $-2\beta_s$ in terms of β and γ . To determine $-2\beta_s$ in analysis B, the world average value of γ from tree-level decays,

Table 1
Current knowledge of CP violation parameters and CP-averaged branching fractions of $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$, $B^+ \rightarrow \pi^+\pi^0$ and $B_s^0 \rightarrow K^+K^-$ decays from BaBar, Belle, CDF and LHCb. The parameter $\rho(X, Y)$ is the statistical correlation between X and Y . The first uncertainties are statistical and the second systematic.

Quantity	BaBar	Belle	CDF	LHCb
$C_{\pi^+\pi^-}$	$-0.25 \pm 0.08 \pm 0.02$	$-0.33 \pm 0.06 \pm 0.03$	–	$-0.38 \pm 0.15 \pm 0.02$
$S_{\pi^+\pi^-}$	$-0.68 \pm 0.10 \pm 0.03$	$-0.64 \pm 0.08 \pm 0.03$	–	$-0.71 \pm 0.13 \pm 0.02$
$\rho(C_{\pi^+\pi^-}, S_{\pi^+\pi^-})$	–0.06	–0.10	–	0.38
$\mathcal{B}_{\pi^+\pi^-} \times 10^6$	$5.5 \pm 0.4 \pm 0.3$	$5.04 \pm 0.21 \pm 0.18$	$5.02 \pm 0.33 \pm 0.35$	$5.08 \pm 0.17 \pm 0.37$
$C_{K^+K^-}$	–	–	–	$0.14 \pm 0.11 \pm 0.03$
$S_{K^+K^-}$	–	–	–	$0.30 \pm 0.12 \pm 0.04$
$\rho(C_{K^+K^-}, S_{K^+K^-})$	–	–	–	0.02
$\mathcal{B}_{K^+K^-} \times 10^6$	–	$38_{-9}^{+10} \pm 7$	$25.8 \pm 2.2 \pm 1.7$	$23.0 \pm 0.7 \pm 2.3$
$\mathcal{A}_{\pi^+\pi^0}$	$-0.03 \pm 0.08 \pm 0.01$	$-0.025 \pm 0.043 \pm 0.007$	–	–
$\mathcal{B}_{\pi^+\pi^0} \times 10^6$	$5.02 \pm 0.46 \pm 0.29$	$5.86 \pm 0.26 \pm 0.38$	–	–
$C_{\pi^0\pi^0}$	$-0.43 \pm 0.26 \pm 0.05$	$-0.44_{-0.52}^{+0.53} \pm 0.17$	–	–
$\mathcal{B}_{\pi^0\pi^0} \times 10^6$	$1.83 \pm 0.21 \pm 0.13$	$2.3_{-0.5-0.3}^{+0.4+0.2}$	–	–

Table 2
Experimental inputs used for the determination of γ and $-2\beta_s$ from $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays using U-spin symmetry. The parameter $\rho(X, Y)$ is the statistical correlation between X and Y . For $C_{\pi^+\pi^-}$ and $S_{\pi^+\pi^-}$ we perform our own weighted average of BaBar, Belle and LHCb results, accounting for correlations.

Quantity	Value	Source
$C_{\pi^+\pi^-}$	-0.30 ± 0.05	This Letter
$S_{\pi^+\pi^-}$	-0.66 ± 0.06	This Letter
$\rho(C_{\pi^+\pi^-}, S_{\pi^+\pi^-})$	–0.007	This Letter
$C_{K^+K^-}$	0.14 ± 0.11	LHCb [25]
$S_{K^+K^-}$	0.30 ± 0.13	LHCb [25]
$\rho(C_{K^+K^-}, S_{K^+K^-})$	0.02	LHCb [25]
$\mathcal{B}_{\pi^+\pi^-} \times 10^6$	5.10 ± 0.19	HFAG [17]
$\mathcal{B}_{K^+K^-} \times 10^6$	24.5 ± 1.8	HFAG [17]
$\sin 2\beta$	0.682 ± 0.019	HFAG [17]
γ (analysis B only)	$(70.1 \pm 7.1)^\circ$	UTfit [8]
λ	0.2253 ± 0.0007	PDG [33]
m_{B^0} [MeV/ c^2]	5279.55 ± 0.26	PDG [33]
$m_{B_s^0}$ [MeV/ c^2]	5366.7 ± 0.4	PDG [33]
m_{π^+} [MeV/ c^2]	139.57018 ± 0.00035	PDG [33]
m_{K^+} [MeV/ c^2]	493.677 ± 0.013	PDG [33]
τ_{B^0} [ps]	1.519 ± 0.007	HFAG [17]
$\tau_{B_s^0}$ [ps]	1.516 ± 0.011	HFAG [17]
$\Delta\Gamma_s/\Gamma_s$	0.160 ± 0.020	LHCb [16]
$\tau(B_s^0 \rightarrow K^+K^-)$ [ps]	1.452 ± 0.042	LHCb [17,34,35]

Table 3
Ranges of flat priors used for the determination of γ and $-2\beta_s$ from $B^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow K^+K^-$ decays using U-spin symmetry.

Quantity	Prior range
d	[0, 20]
ϑ	$[-180^\circ, 180^\circ]$
r_D	[0, κ]
ϑ_{r_D}	$[-180^\circ, 180^\circ]$
r_G	[0, κ]
ϑ_{r_G}	$[-180^\circ, 180^\circ]$
γ (analysis A only)	$[-180^\circ, 180^\circ]$
$-2\beta_s$ [rad] (analysis B only)	$[-\pi, \pi]$

$\gamma = (70.1 \pm 7.1)^\circ$ [8], is used as an input, and $-2\beta_s$ is left as a free parameter.

The inputs to the analyses are the measured values of $C_{\pi^+\pi^-}$, $S_{\pi^+\pi^-}$, $C_{K^+K^-}$, $S_{K^+K^-}$, $\mathcal{B}_{\pi^+\pi^-}$ and $\mathcal{B}_{K^+K^-}$. The corresponding constraints are given in Eqs. (12), (13), (16), (17), (18) and (21). In addition, the value of $A_{K^+K^-}^{\Delta\Gamma}$ is fixed to be negative. A summary of the experimental inputs is given in Table 2.

In both analyses, flat prior probability distributions, hereinafter referred to as priors, on d , ϑ , r_D , ϑ_{r_D} , r_G , ϑ_{r_G} and, where appropriate, on γ and $-2\beta_s$ are used. In particular, we allow the U-spin breaking phases ϑ_{r_D} and ϑ_{r_G} to be completely undeter-

mined, using flat priors between -180° and 180° . Concerning the parameters r_D and r_G , we adopt uniform priors between 0 and κ , where κ represents the maximum magnitude of non-factorizable U-spin breaking allowed. The ranges of the flat priors are summarized in Table 3. We study the sensitivity on γ and $-2\beta_s$ as a function of κ , ranging from 0 to 1, meaning from 0% up to 100% non-factorizable U-spin breaking. For all experimental inputs we use Gaussian PDFs. The values of $|D'|$, d' and ϑ' are determined using Eqs. (29) and (30).

The dependences on κ of the 68% and 95% posterior probability intervals for γ and $-2\beta_s$ are shown in Fig. 1. When the allowed amount of U-spin breaking becomes large enough, the PDF for γ is poorly constrained. In particular, it can be noted that for values of κ exceeding 0.6 the sensitivity on γ reduces significantly as a function of increasing κ . This fast transition is related to the non-linearity of the constraint equations. For $-2\beta_s$ the dependence of the sensitivity on κ is mild, but for values of κ exceeding 0.6 a slight shift of the distribution towards more negative values is observed.

In Fig. 2 we show the PDFs for γ obtained from analysis A and for $-2\beta_s$ obtained from analysis B, corresponding to $\kappa = 0.5$. The numerical results from both analyses are reported in Table 4. The 68% probability interval for γ is $[56^\circ, 70^\circ]$, and that for $-2\beta_s$ is $[-0.28, 0.02]$ rad.

5. Inclusion of physics observables from $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays

A method to determine the angle α of the UT using CP asymmetries and branching fractions of $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays was proposed in Ref. [14]. This method relies on the isospin symmetry of strong interactions and on the assumption of negligible contributions from electroweak penguin amplitudes. Isospin breaking and electroweak penguin contributions are known to be small, and their impact on the determination of the weak phase is at the level of 1° [36–39]. In Ref. [13] it was suggested to combine the isospin-based technique of Ref. [14] with that of Ref. [10] based on U-spin. Here we extend the study presented in Section 4 by including the experimental information on $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays, i.e. using also the observables $C_{\pi^0\pi^0}$, $\mathcal{B}_{\pi^0\pi^0}$ and $\mathcal{B}_{\pi^+\pi^0}$. The corresponding constraints are given in Eqs. (14), (19) and (20).

In complete analogy with the study presented in Section 4, we perform two distinct analyses, to determine either γ or $-2\beta_s$. They are referred to as analyses C and D, respectively. In analysis C, the value of $-2\beta_s$ is constrained as a function of β and γ , and γ is determined, whereas in analysis D, the world average




Fig. 1. Dependences of the 68% (hatched areas) and 95% (filled areas) probability intervals on the allowed amount of non-factorizable U-spin breaking, for (a) γ from analysis A and (b) $-2\beta_s$ from analysis B.




Fig. 2. Distributions of (a) γ from analysis A and (b) $-2\beta_s$ from analysis B, corresponding to $\kappa = 0.5$. The hatched areas correspond to 68% probability intervals, whereas the filled areas correspond to 95% probability intervals.

Table 4

Results obtained from analyses A and B with $\kappa = 0.5$. The results are given modulo 180° for ϑ , ϑ' and γ .

Quantity	Analysis A		Analysis B	
	68% prob.	95% prob.	68% prob.	95% prob.
d	[0.32, 0.53]	[0.25, 0.78]	[0.36, 0.58]	[0.29, 0.75]
ϑ	[136°, 157°]	[119°, 165°]	[141°, 157°]	[129°, 163°]
d'	[0.33, 0.50]	[0.28, 0.65]	[0.34, 0.52]	[0.28, 0.69]
ϑ'	[132°, 160°]	[114°, 176°]	[132°, 160°]	[117°, 175°]
$ D $ [MeV $^{\frac{1}{2}}$ ps $^{-\frac{1}{2}}$]	[0.102, 0.114]	[0.094, 0.121]	[0.101, 0.112]	[0.095, 0.117]
$ D' $ [MeV $^{\frac{1}{2}}$ ps $^{-\frac{1}{2}}$]	[0.130, 0.195]	[0.097, 0.231]	[0.122, 0.188]	[0.090, 0.224]
γ	[56°, 70°]	[49°, 82°]	–	–
$-2\beta_s$ [rad]	–	–	[–0.28, 0.02]	[–0.44, 0.17]

value of γ from tree-level decays is used as an input and $-2\beta_s$ is determined. A summary of the experimental inputs is given in Table 5.

In both analyses, flat priors on d , ϑ , q , ϑ_q , r_D , ϑ_{r_D} , r_G , ϑ_{r_G} and, where appropriate, on γ and $-2\beta_s$ are used. The ranges of the flat priors are summarized in Table 6. For all experimental inputs we use Gaussian PDFs. The values of $|D|$, d' and ϑ' are again determined using Eqs. (29) and (30).

The dependences on κ of the 68% and 95% probability intervals for γ and $-2\beta_s$ are shown in Fig. 3. Again, when the amount of U-spin breaking exceeds 60%, additional maxima appear in the posterior PDF for γ . By contrast, for $-2\beta_s$, the dependence of the sensitivity on κ is very weak. In Fig. 4 we show the PDFs for γ obtained from analysis C and for $-2\beta_s$ obtained from analysis D, corresponding to $\kappa = 0.5$. The numerical results from both analyses are reported in Table 7. The 68% probability interval for γ is [57°, 71°], and that for $-2\beta_s$ is [–0.28, 0.02] rad.

It is worth emphasizing that, although this study is similar to that presented in Ref. [13], there are two relevant differences, in

addition to the use of updated experimental inputs. First, the upper limits of the priors on d and q are chosen to be much larger, to include all nonzero likelihood regions and to remove any sizable dependence of the results on the choice of the priors. In particular, this leads to a bigger impact of U-spin breaking effects at very large κ values. Second, the adopted parameterization of non-factorizable U-spin breaking is slightly different, in order to propagate equally the effects of the breaking on every topology contributing to the total decay amplitudes.

6. Results and conclusions

Using the latest LHCb measurements of time-dependent CP violation in the $B_s^0 \rightarrow K^+K^-$ decay, and following the approaches outlined in Refs. [10,13], the angle γ of the unitarity triangle and the B_s^0 mixing phase $-2\beta_s$ have been determined. The approach of Ref. [10] relies on the use of the U-spin symmetry of strong interactions relating $B_s^0 \rightarrow K^+K^-$ with $B^0 \rightarrow \pi^+\pi^-$ decay amplitudes, whereas that of Ref. [13] relies on both isospin and U-spin

symmetries by combining the methods proposed in Refs. [10] and [14], i.e. considering also the information from $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ decays. To follow the latter approach, measure-

Table 5

Experimental inputs used for the determination of γ and $-2\beta_s$ from $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$, $B^+ \rightarrow \pi^+\pi^0$ and $B_s^0 \rightarrow K^+K^-$ decays, using isospin and U-spin symmetries. The parameter $\rho(X, Y)$ is the statistical correlation between X and Y . For $C_{\pi^+\pi^-}$ and $S_{\pi^+\pi^-}$ we perform our own weighted average of BaBar, Belle and LHCb results, accounting for correlations.

Quantity	Value	Source
$C_{\pi^+\pi^-}$	-0.30 ± 0.05	This Letter
$S_{\pi^+\pi^-}$	-0.66 ± 0.06	This Letter
$\rho(C_{\pi^+\pi^-}, S_{\pi^+\pi^-})$	-0.007	This Letter
$C_{\pi^0\pi^0}$	-0.43 ± 0.24	HFAG [17]
$C_{K^+K^-}$	0.14 ± 0.11	LHCb [25]
$S_{K^+K^-}$	0.30 ± 0.13	LHCb [25]
$\rho(C_{K^+K^-}, S_{K^+K^-})$	0.02	LHCb [25]
$\mathcal{B}_{\pi^+\pi^-} \times 10^6$	5.10 ± 0.19	HFAG [17]
$\mathcal{B}_{\pi^+\pi^0} \times 10^6$	5.48 ± 0.35	HFAG [17]
$\mathcal{B}_{\pi^0\pi^0} \times 10^6$	1.91 ± 0.23	HFAG [17]
$\mathcal{B}_{K^+K^-} \times 10^6$	24.5 ± 1.8	HFAG [17]
$\sin 2\beta$	0.682 ± 0.019	HFAG [17]
γ (analysis D only)	$(70.1 \pm 7.1)^\circ$	UTfit [8]
λ	0.2253 ± 0.0007	PDG [33]
m_{B^0} [MeV/ c^2]	5279.55 ± 0.26	PDG [33]
m_{B^+} [MeV/ c^2]	5279.25 ± 0.26	PDG [33]
$m_{B_s^0}$ [MeV/ c^2]	5366.7 ± 0.4	PDG [33]
m_{π^+} [MeV/ c^2]	139.57018 ± 0.00035	PDG [33]
m_{π^0} [MeV/ c^2]	134.9766 ± 0.0006	PDG [33]
m_{K^+} [MeV/ c^2]	493.677 ± 0.013	PDG [33]
τ_{B^0} [ps]	1.519 ± 0.007	HFAG [17]
τ_{B^+} [ps]	1.641 ± 0.008	HFAG [17]
$\tau_{B_s^0}$ [ps]	1.516 ± 0.011	HFAG [17]
$\Delta\Gamma_s/\Gamma_s$	0.160 ± 0.020	LHCb [16]
$\tau(B_s^0 \rightarrow K^+K^-)$ [ps]	1.452 ± 0.042	LHCb [17,34,35]

ments solely coming from other experiments have been included in the analysis.

We have studied the impact of large non-factorizable U-spin breaking corrections on the determination of γ and $-2\beta_s$. The relevant results in terms of 68% and 95% probability intervals, which include uncertainties due to non-factorizable U-spin breaking effects up to 50%, are summarized in Fig. 5. Typical U-spin breaking effects, including factorizable contributions, are expected to be much smaller, around the 30% level [30,31].

With up to 50% non-factorizable U-spin breaking, the approach of Ref. [13] gives marginal improvements in precision with respect to that of Ref. [10]. The former approach gives considerably more robust results for larger U-spin breaking values. Following the approach of Ref. [13] and taking the most probable value as central value, at 68% probability we obtain

$$\gamma = (63.5_{-6.7}^{+7.2})^\circ,$$

Table 6

Ranges of flat priors used for the determination of γ and $-2\beta_s$ from $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$, $B^+ \rightarrow \pi^+\pi^0$ and $B_s^0 \rightarrow K^+K^-$ decays, using isospin and U-spin symmetries.

Quantity	Prior range
d	[0, 20]
ϑ	$[-180^\circ, 180^\circ]$
q	[0, 20]
ϑ_q	$[-180^\circ, 180^\circ]$
r_D	[0, κ]
ϑ_{r_D}	$[-180^\circ, 180^\circ]$
r_G	[0, κ]
ϑ_{r_G}	$[-180^\circ, 180^\circ]$
γ (analysis C only)	$[-180^\circ, 180^\circ]$
$-2\beta_s$ [rad] (analysis D only)	$[-\pi, \pi]$




Fig. 3. Dependences of the 68% (hatched areas) and 95% (filled areas) probability intervals on the allowed amount of non-factorizable U-spin breaking, for (a) γ from analysis C and (b) $-2\beta_s$ from analysis D.





Fig. 4. Distributions of (a) γ from analysis C and (b) $-2\beta_s$ from analysis D, corresponding to $\kappa = 0.5$. The hatched areas correspond to 68% probability intervals, whereas the filled areas correspond to 95% probability intervals.

Table 7Results obtained from analyses C and D with $\kappa = 0.5$. The results are given modulo 180° for ϑ , ϑ' and γ .

Quantity	Analysis C		Analysis D	
	68% prob.	95% prob.	68% prob.	95% prob.
d	[0.33, 0.57]	[0.28, 0.79]	[0.37, 0.59]	[0.31, 0.77]
ϑ	[139°, 157°]	[125°, 164°]	[142°, 157°]	[132°, 163°]
d'	[0.34, 0.50]	[0.28, 0.65]	[0.34, 0.52]	[0.29, 0.70]
ϑ'	[132°, 160°]	[119°, 176°]	[133°, 160°]	[119°, 176°]
q	[1.04, 1.21]	[0.94, 1.30]	[1.04, 1.21]	[0.95, 1.30]
ϑ_q	[−82°, −58°]	[−88°, −35°]	[−78°, −57°]	[−85°, 38°]
$ D $ [MeV $^{\frac{1}{2}}$ ps $^{-\frac{1}{2}}$]	[0.101, 0.113]	[0.094, 0.118]	[0.100, 0.111]	[0.094, 0.116]
$ D' $ [MeV $^{\frac{1}{2}}$ ps $^{-\frac{1}{2}}$]	[0.129, 0.193]	[0.097, 0.228]	[0.122, 0.187]	[0.089, 0.221]
γ	[57°, 71°]	[52°, 82°]	–	–
$-2\beta_s$ [rad]	–	–	[−0.28, 0.02]	[−0.44, 0.17]

**Fig. 5.** Results for (top) γ and (bottom) $-2\beta_s$ with 50% ($\kappa = 0.5$) non-factorizable U-spin breaking. As a comparison, other reference values are also reported. The most likely values are indicated by the vertical lines inside the boxes. The boxes and the error bars delimit the 68% and 95% probability intervals, respectively.

and, in an alternative analysis,

$$-2\beta_s = -0.12^{+0.14}_{-0.16} \text{ rad.}$$

These results have been verified to be robust with respect to the choice of the priors and of the parameterization of non-factorizable U-spin breaking contributions. The value of γ shows no significant deviation from the averages of γ from tree-level decays provided by the UTfit Collaboration and the CKMfitter group that quote $\gamma = (70.1 \pm 7.1)^\circ$ and $\gamma = (68.0^{+8.0}_{-8.5})^\circ$, respectively [8,9]. Analogously, the value of $-2\beta_s$ is compatible with the LHCb result from $b \rightarrow c\bar{c}s$ transitions, $\phi_s = 0.01 \pm 0.07$ (stat) ± 0.01 (syst) rad [16], obtained using a data sample of pp collisions corresponding to an integrated luminosity of 1.0 fb^{-1} .

In summary, the value of γ from charmless two-body decays of beauty mesons is found to be compatible and competitive with that from tree-level decays. However, since the impact of U-spin breaking corrections is significant, further improvements in the measurement of γ are primarily limited by theoretical understanding of U-spin breaking. By contrast, the impact of U-spin breaking

effects on the value of $-2\beta_s$ is small, and significant improvements are anticipated with the advent of larger samples of data. It is worth emphasizing that the information on $-2\beta_s$ comes solely from the measurement of CP violation in the $B_s^0 \rightarrow K^+ K^-$ decay [25], also based on a data sample of pp collisions corresponding to an integrated luminosity of 1.0 fb^{-1} . At present, the overall uncertainty on $-2\beta_s$, which also includes theoretical uncertainties, is only two times larger than that obtained using $b \rightarrow c\bar{c}s$ transitions, as reported above.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MINECO (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are indebted to the communities behind the multiple open source software packages on which we depend. We are also thankful for the computing resources and the access to software R&D tools provided by Yandex LLC (Russia). Individual groups or members have received support from EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR (Russia), XuntaGal and GENCAT (Spain), Royal Society and Royal Commission for the Exhibition of 1851 (United Kingdom).

References

- [1] N. Cabibbo, Unitary symmetry and leptonic decays, *Phys. Rev. Lett.* 10 (1963) 531.
- [2] M. Kobayashi, T. Maskawa, CP violation in the renormalizable theory of weak interaction, *Prog. Theor. Phys.* 49 (1973) 652.
- [3] J. Brod, J. Zupan, The ultimate theoretical error on γ from $B \rightarrow DK$ decays, *J. High Energy Phys.* 1401 (2014) 051, arXiv:1308.5663.
- [4] BaBar Collaboration, J.P. Lees, et al., Observation of direct CP violation in the measurement of the Cabibbo–Kobayashi–Maskawa angle γ with $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ decays, *Phys. Rev. D* 87 (2013) 052015, arXiv:1301.1029.
- [5] K. Trabelsi, Study of direct CP in charmed B decays and measurement of the CKM angle γ at Belle, arXiv:1301.2033.
- [6] LHCb Collaboration, R. Aaij, et al., Measurement of the CKM angle γ from a combination of $B^\pm \rightarrow Dh^\pm$ analyses, *Phys. Lett. B* 726 (2013) 151, arXiv:1305.2050.
- [7] LHCb Collaboration, Improved constraints on γ from $B^\pm \rightarrow DK^\pm$ decays including first results on 2012 data, LHCb-CONF-2013-006.

- [8] UTfit Collaboration, M. Bona, et al., The unitarity triangle fit in the Standard Model and hadronic parameters from lattice QCD: a reappraisal after the measurements of Δm_s and $\text{BR}(B \rightarrow \tau \nu_\tau)$, *J. High Energy Phys.* 0610 (2006) 081, arXiv:hep-ph/0606167, updated results and plots available at <http://www.utfit.org/>.
- [9] CKMfitter group, J. Charles, et al., Predictions of selected flavour observables within the Standard Model, *Phys. Rev. D* 84 (2011) 033005, arXiv:1106.4041, updated results and plots available at <http://ckmfitter.in2p3.fr/>.
- [10] R. Fleischer, New strategies to extract β and γ from $B_d \rightarrow \pi^+ \pi^-$ and $B_s \rightarrow K^+ K^-$, *Phys. Lett. B* 459 (1999) 306, arXiv:hep-ph/9903456.
- [11] R. Fleischer, $B_{s,d} \rightarrow \pi \pi$, πK , KK : status and prospects, *Eur. Phys. J. C* 52 (2007) 267, arXiv:0705.1121.
- [12] R. Fleischer, R. Kneegjens, In pursuit of new physics with $B_s^0 \rightarrow K^+ K^-$, *Eur. Phys. J. C* 71 (2011) 1532, arXiv:1011.1096.
- [13] M. Ciuchini, E. Franco, S. Mishima, L. Silvestrini, Testing the Standard Model and searching for new physics with $B_d \rightarrow \pi \pi$ and $B_s \rightarrow KK$ decays, *J. High Energy Phys.* 1210 (2012) 029, arXiv:1205.4948.
- [14] M. Gronau, D. London, Isospin analysis of CP asymmetries in B decays, *Phys. Rev. Lett.* 65 (1990) 3381.
- [15] LHCb Collaboration, R. Aaij, et al., Determination of the sign of the decay width difference in the B_s^0 system, *Phys. Rev. Lett.* 108 (2012) 241801, arXiv:1202.4717.
- [16] LHCb Collaboration, R. Aaij, et al., Measurement of CP violation and the B_s^0 meson decay width difference with $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays, *Phys. Rev. D* 87 (2013) 112010, arXiv:1304.2600.
- [17] Heavy Flavor Averaging Group, Y. Amhis, et al., Averages of b -hadron, c -hadron, and τ -lepton properties as of early 2012, arXiv:1207.1158, updated results and plots available at <http://www.slac.stanford.edu/xorg/hfag/>.
- [18] LHCb Collaboration, R. Aaij, et al., Measurement of the flavour-specific CP-violating asymmetry $a_{\text{CP}}^{\text{eff}}$ in B_s^0 decays, *Phys. Lett. B* 728 (2014) 607, arXiv:1308.1048.
- [19] K. De Bruyn, et al., Branching ratio measurements of B_s decays, *Phys. Rev. D* 86 (2012) 014027, arXiv:1204.1735.
- [20] BaBar Collaboration, J.P. Lees, et al., Measurement of CP asymmetries and branching fractions in charmless two-body B-meson decays to pions and kaons, *Phys. Rev. D* 87 (2013) 052009, arXiv:1206.3525.
- [21] Belle Collaboration, I. Adachi, et al., Measurement of the CP violation parameters in $B^0 \rightarrow \pi^+ \pi^-$ decays, *Phys. Rev. D* 88 (2013) 092003, arXiv:1302.0551.
- [22] BaBar Collaboration, B. Aubert, et al., Study of $B^0 \rightarrow \pi^0 \pi^0$, $B^\pm \rightarrow \pi^\pm \pi^0$, and $B^\pm \rightarrow K^\pm \pi^0$ decays, and isospin analysis of $B \rightarrow \pi \pi$ decays, *Phys. Rev. D* 76 (2007) 091102, arXiv:0707.2798.
- [23] Belle Collaboration, Y.-T. Duh, et al., Measurements of branching fractions and direct CP asymmetries for $B \rightarrow K\pi$, $B \rightarrow \pi\pi$ and $B \rightarrow KK$ decays, *Phys. Rev. D* 87 (2013) 031103, arXiv:1210.1348.
- [24] Belle Collaboration, K. Abe, et al., Observation of $B^0 \rightarrow \pi^0 \pi^0$, *Phys. Rev. Lett.* 94 (2005) 181803, arXiv:hep-ex/0408101.
- [25] LHCb Collaboration, R. Aaij, et al., First measurement of time-dependent CP violation in $B_s^0 \rightarrow K^+ K^-$ decays, *J. High Energy Phys.* 1310 (2013) 183, arXiv:1308.1428.
- [26] BaBar Collaboration, B. Aubert, et al., Improved measurements of the branching fractions for $B^0 \rightarrow \pi^+ \pi^-$ and $B^0 \rightarrow K^+ \pi^-$, and a search for $B^0 \rightarrow K^+ K^-$, *Phys. Rev. D* 75 (2007) 012008, arXiv:hep-ex/0608003.
- [27] CDF Collaboration, T. Aaltonen, et al., Measurements of direct CP violating asymmetries in charmless decays of strange bottom mesons and bottom baryons, *Phys. Rev. Lett.* 106 (2011) 181802, arXiv:1103.5762.
- [28] LHCb Collaboration, R. Aaij, et al., Measurement of b -hadron branching fractions for two-body decays into charmless charged hadrons, *J. High Energy Phys.* 1210 (2012) 037, arXiv:1206.2794.
- [29] LHCb Collaboration, R. Aaij, et al., Effective lifetime measurements in the $B_s^0 \rightarrow K^+ K^-$, $B^0 \rightarrow K^+ \pi^-$ and $B_s^0 \rightarrow \pi^+ K^-$ decays, *Phys. Lett. B* (2014), in press, arXiv:1406.7204.
- [30] M. Gronau, U-spin breaking in CP asymmetries in B decays, *Phys. Lett. B* 727 (2013) 136, arXiv:1308.3448.
- [31] M. Nagashima, A. Szytnkman, D. London, U-spin tests of the standard model and new physics, *Mod. Phys. Lett. A* 23 (2008) 1175, arXiv:hep-ph/0701199.
- [32] G. Duplancić, B. Melić, B. $B_s \rightarrow K$ form factors: an update of light-cone sum rule results, *Phys. Rev. D* 78 (2008) 054015, arXiv:0805.4170.
- [33] Particle Data Group, J. Beringer, et al., Review of particle physics, *Phys. Rev. D* 86 (2012) 010001, and 2013 partial update for the 2014 edition.
- [34] LHCb Collaboration, R. Aaij, et al., Measurement of the effective $B_s^0 \rightarrow K^+ K^-$ lifetime, *Phys. Lett. B* 707 (2012) 349, arXiv:1111.0521.
- [35] LHCb Collaboration, R. Aaij, et al., Measurement of the effective $B_s^0 \rightarrow K^+ K^-$ lifetime, *Phys. Lett. B* 716 (2012) 393, arXiv:1207.5993.
- [36] A.J. Buras, R. Fleischer, A general analysis of γ determinations from $B \rightarrow \pi K$ decays, *Eur. Phys. J. C* 11 (1999) 93, arXiv:hep-ph/9810260.
- [37] M. Gronau, D. Pirjol, T.-M. Yan, Model independent electroweak penguins in B decays to two pseudoscalars, *Phys. Rev. D* 60 (1999) 034021, arXiv:hep-ph/9810482.
- [38] J. Zupan, Penguin pollution estimates relevant for the extraction of α/ϕ_2 , *Nucl. Phys. B, Proc. Suppl.* 170 (2007) 33, arXiv:hep-ph/0701004.
- [39] F. Botella, D. London, J.P. Silva, Looking for $\Delta I = 5/2$ amplitude components in $B \rightarrow \pi\pi$ and $B \rightarrow \rho\rho$ experiments, *Phys. Rev. D* 73 (2006) 071501, arXiv:hep-ph/0602060.

LHCb Collaboration

R. Aaij⁴¹, C. Abellán Beteta⁴⁰, B. Adeva³⁷, M. Adinolfi⁴⁶, A. Affolder⁵², Z. Ajaltouni⁵, S. Akar⁶, J. Albrecht⁹, F. Alessio³⁸, M. Alexander⁵¹, S. Ali⁴¹, G. Alkhazov³⁰, P. Alvarez Cartelle³⁷, A.A. Alves Jr^{25,38}, S. Amato², S. Amerio²², Y. Amhis⁷, L. An³, L. Anderlini^{17,g}, J. Anderson⁴⁰, R. Andreassen⁵⁷, M. Andreotti^{16,f}, J.E. Andrews⁵⁸, R.B. Appleby⁵⁴, O. Aquines Gutierrez¹⁰, F. Archilli³⁸, A. Artamonov³⁵, M. Artuso⁵⁹, E. Aslanides⁶, G. Auriemma^{25,n}, M. Baalouch⁵, S. Bachmann¹¹, J.J. Back⁴⁸, A. Badalov³⁶, C. Baesso⁶⁰, W. Baldini¹⁶, R.J. Barlow⁵⁴, C. Barschel³⁸, S. Barsuk⁷, W. Barter⁴⁷, V. Batozskaya²⁸, V. Battista³⁹, A. Bay³⁹, L. Beaucourt⁴, J. Beddow⁵¹, F. Bedeschi²³, I. Bediaga¹, S. Belogurov³¹, K. Belous³⁵, I. Belyaev³¹, E. Ben-Haim⁸, G. Bencivenni¹⁸, S. Benson³⁸, J. Benton⁴⁶, A. Berezhnoy³², R. Bernet⁴⁰, M.-O. Bettler⁴⁷, M. van Beuzekom⁴¹, A. Bien¹¹, S. Bifani⁴⁵, T. Bird⁵⁴, A. Bizzeti^{17,i}, P.M. Bjørnstad⁵⁴, T. Blake⁴⁸, F. Blanc³⁹, J. Blouw¹⁰, S. Blusk⁵⁹, V. Bocci²⁵, A. Bondar³⁴, N. Bondar^{30,38}, W. Bonivento^{15,38}, S. Borghi⁵⁴, A. Borgia⁵⁹, M. Borsato⁷, T.J.V. Bowcock⁵², E. Bowen⁴⁰, C. Bozzi¹⁶, T. Brambach⁹, J. Bressieux³⁹, D. Brett⁵⁴, M. Britsch¹⁰, T. Britton⁵⁹, J. Brodzicka⁵⁴, N.H. Brook⁴⁶, H. Brown⁵², A. Bursche⁴⁰, G. Busetto^{22,r}, J. Buytaert³⁸, S. Cadeddu¹⁵, R. Calabrese^{16,f}, M. Calvi^{20,k}, M. Calvo Gomez^{36,p}, P. Campana^{18,38}, D. Campora Perez³⁸, A. Carbone^{14,d}, G. Carboni^{24,l}, R. Cardinale^{19,38,j}, A. Cardini¹⁵, L. Carson⁵⁰, K. Carvalho Akiba², G. Casse⁵², L. Cassina²⁰, L. Castillo Garcia³⁸, M. Cattaneo³⁸, Ch. Cauet⁹, R. Cenci⁵⁸, M. Charles⁸, Ph. Charpentier³⁸, M. Chefdeville⁴, S. Chen⁵⁴, S.-F. Cheung⁵⁵, N. Chiapolini⁴⁰, M. Chrzaszcz^{40,26}, K. Ciba³⁸, X. Cid Vidal³⁸, G. Ciezarek⁵³, P.E.L. Clarke⁵⁰, M. Clemencic³⁸, H.V. Cliff⁴⁷, J. Closier³⁸, V. Coco³⁸, J. Cogan⁶, E. Cogneras⁵, L. Cojocariu²⁹, P. Collins³⁸, A. Comerma-Montells¹¹, A. Contu^{15,38}, A. Cook⁴⁶, M. Coombes⁴⁶, S. Coquereau⁸, G. Corti³⁸, M. Corvo^{16,f}, I. Counts⁵⁶, B. Couturier³⁸, G.A. Cowan⁵⁰, D.C. Craik⁴⁸, M. Cruz Torres⁶⁰, S. Cunliffe⁵³, R. Currie⁵⁰, C. D'Ambrosio³⁸, J. Dalseno⁴⁶, P. David⁸,

P.N.Y. David⁴¹, A. Davis⁵⁷, K. De Bruyn⁴¹, S. De Capua⁵⁴, M. De Cian¹¹, J.M. De Miranda¹, L. De Paula², W. De Silva⁵⁷, P. De Simone¹⁸, D. Decamp⁴, M. Deckenhoff⁹, L. Del Buono⁸, N. Déleage⁴, D. Derkach⁵⁵, O. Deschamps⁵, F. Dettori³⁸, A. Di Canto³⁸, H. Dijkstra³⁸, S. Donleavy⁵², F. Dordei¹¹, M. Dorigo³⁹, A. Dosil Suárez³⁷, D. Dossett⁴⁸, A. Dovbnya⁴³, K. Dreimanis⁵², G. Dujany⁵⁴, F. Dupertuis³⁹, P. Durante³⁸, R. Dzhelyadin³⁵, A. Dziurda²⁶, A. Dzyuba³⁰, S. Easo^{49,38}, V. Egorychev³¹, S. Eidelman³⁴, S. Eisenhardt⁵⁰, U. Eitschberger⁹, R. Ekelhof⁹, L. Eklund⁵¹, I. El Rifai⁵, Ch. Elsasser⁴⁰, S. Ely⁵⁹, S. Esen¹¹, H.-M. Evans⁴⁷, T. Evans⁵⁵, A. Falabella¹⁴, C. Färber¹¹, C. Farinelli⁴¹, N. Farley⁴⁵, S. Farry⁵², R.F. Fay⁵², D. Ferguson⁵⁰, V. Fernandez Albor³⁷, F. Ferreira Rodrigues¹, M. Ferro-Luzzi³⁸, S. Filippov³³, M. Fiore^{16,f}, M. Fiorini^{16,f}, M. Firlej²⁷, C. Fitzpatrick³⁹, T. Fiutowski²⁷, P. Fol⁵³, M. Fontana¹⁰, F. Fontanelli^{19,j}, R. Forty³⁸, O. Francisco², M. Frank³⁸, C. Frei³⁸, M. Frosini^{17,g}, J. Fu^{21,38}, E. Furfaro^{24,l}, A. Gallas Torreira³⁷, D. Galli^{14,d}, S. Gallorini^{22,38}, S. Gambetta^{19,j}, M. Gandelman², P. Gandini⁵⁹, Y. Gao³, J. García Pardiñas³⁷, J. Garofoli⁵⁹, J. Garra Tico⁴⁷, L. Garrido³⁶, C. Gaspar³⁸, R. Gauld⁵⁵, L. Gavardi⁹, G. Gavrilo³⁰, A. Geraci^{21,v}, E. Gersabeck¹¹, M. Gersabeck⁵⁴, T. Gershon⁴⁸, Ph. Ghez⁴, A. Gianelle²², S. Gianì³⁹, V. Gibson⁴⁷, L. Giubega²⁹, V.V. Gligorov³⁸, C. Göbel⁶⁰, D. Golubkov³¹, A. Golutvin^{53,31,38}, A. Gomes^{1,a}, C. Gotti²⁰, M. Grabalosa Gándara⁵, R. Graciani Diaz³⁶, L.A. Granado Cardoso³⁸, E. Graugés³⁶, G. Graziani¹⁷, A. Grecu²⁹, E. Greening⁵⁵, S. Gregson⁴⁷, P. Griffith⁴⁵, L. Grillo¹¹, O. Grünberg⁶², B. Gui⁵⁹, E. Gushchin³³, Yu. Guz^{35,38}, T. Gys³⁸, C. Hadjivasiliou⁵⁹, G. Haefeli³⁹, C. Haen³⁸, S.C. Haines⁴⁷, S. Hall⁵³, B. Hamilton⁵⁸, T. Hampson⁴⁶, X. Han¹¹, S. Hansmann-Menzemer¹¹, N. Harnew⁵⁵, S.T. Harnew⁴⁶, J. Harrison⁵⁴, J. He³⁸, T. Head³⁸, V. Heijne⁴¹, K. Hennessy⁵², P. Henrard⁵, L. Henry⁸, J.A. Hernando Morata³⁷, E. van Herwijnen³⁸, M. Heß⁶², A. Hicheur¹, D. Hill⁵⁵, M. Hoballah⁵, C. Hombach⁵⁴, W. Hulsbergen⁴¹, P. Hunt⁵⁵, N. Hussain⁵⁵, D. Hutchcroft⁵², D. Hynds⁵¹, M. Idzik²⁷, P. Ilten⁵⁶, R. Jacobsson³⁸, A. Jaeger¹¹, J. Jalocha⁵⁵, E. Jans⁴¹, P. Jaton³⁹, A. Jawahery⁵⁸, F. Jing³, M. John⁵⁵, D. Johnson³⁸, C.R. Jones⁴⁷, C. Joram³⁸, B. Jost³⁸, N. Jurik⁵⁹, M. Kaballo⁹, S. Kandybei⁴³, W. Kalso⁶, M. Karacson³⁸, T.M. Karbach³⁸, S. Karodia⁵¹, M. Kelsey⁵⁹, I.R. Kenyon⁴⁵, T. Ketel⁴², B. Khanji²⁰, C. Khurewathanakul³⁹, S. Klaver⁵⁴, K. Klimaszewski²⁸, O. Kochebina⁷, M. Kolpin¹¹, I. Komarov³⁹, R.F. Koopman⁴², P. Koppenburg^{41,38}, M. Korolev³², A. Kozlinskiy⁴¹, L. Kravchuk³³, K. Kreplin¹¹, M. Kreps⁴⁸, G. Krocker¹¹, P. Krokovny³⁴, F. Kruse⁹, W. Kucewicz^{26,o}, M. Kucharczyk^{20,26,k}, V. Kudryavtsev³⁴, K. Kurek²⁸, T. Kvaratskheliya³¹, V.N. La Thi³⁹, D. Lacarrere³⁸, G. Lafferty⁵⁴, A. Lai¹⁵, D. Lambert⁵⁰, R.W. Lambert⁴², G. Lanfranchi¹⁸, C. Langenbruch⁴⁸, B. Langhans³⁸, T. Latham⁴⁸, C. Lazzeroni⁴⁵, R. Le Gac⁶, J. van Leerdam⁴¹, J.-P. Lees⁴, R. Lefèvre⁵, A. Leflat³², J. Lefrançois⁷, S. Leo²³, O. Leroy⁶, T. Lesiak²⁶, B. Leverington¹¹, Y. Li³, T. Likhomanenko⁶³, M. Liles⁵², R. Lindner³⁸, C. Linn³⁸, F. Lionetto⁴⁰, B. Liu¹⁵, S. Lohn³⁸, I. Longstaff⁵¹, J.H. Lopes², N. Lopez-March³⁹, P. Lowdon⁴⁰, H. Lu³, D. Lucchesi^{22,r}, H. Luo⁵⁰, A. Lupato²², E. Luppi^{16,f}, O. Lupton⁵⁵, F. Machefert⁷, I.V. Machikhiliyan³¹, F. Maciuc²⁹, O. Maev³⁰, S. Malde⁵⁵, A. Malinin⁶³, G. Manca^{15,e}, G. Mancinelli⁶, A. Mapelli³⁸, J. Maratas⁵, J.F. Marchand⁴, U. Marconi¹⁴, C. Marin Benito³⁶, P. Marino^{23,t}, R. Märki³⁹, J. Marks¹¹, G. Martellotti²⁵, A. Martens⁸, A. Martín Sánchez⁷, M. Martinelli³⁹, D. Martinez Santos^{42,38}, F. Martinez Vidal⁶⁴, D. Martins Tostes², A. Massafferri¹, R. Matev³⁸, Z. Mathe³⁸, C. Matteuzzi²⁰, A. Mazurov⁴⁵, M. McCann⁵³, J. McCarthy⁴⁵, A. McNab⁵⁴, R. McNulty¹², B. McSkelly⁵², B. Meadows⁵⁷, F. Meier⁹, M. Meissner¹¹, M. Merk⁴¹, D.A. Milanes⁸, M.-N. Minard⁴, N. Moggi¹⁴, J. Molina Rodriguez⁶⁰, S. Monteil⁵, M. Morandin²², P. Morawski²⁷, A. Mordà⁶, M.J. Morello^{23,t}, J. Moron²⁷, A.-B. Morris⁵⁰, R. Mountain⁵⁹, F. Muheim⁵⁰, K. Müller⁴⁰, M. Mussini¹⁴, B. Muster³⁹, P. Naik⁴⁶, T. Nakada³⁹, R. Nandakumar⁴⁹, I. Nasteva², M. Needham⁵⁰, N. Neri²¹, S. Neubert³⁸, N. Neufeld³⁸, M. Neuner¹¹, A.D. Nguyen³⁹, T.D. Nguyen³⁹, C. Nguyen-Mau^{39,q}, M. Nicol⁷, V. Niess⁵, R. Niet⁹, N. Nikitin³², T. Nikodem¹¹, A. Novoselov³⁵, D.P. O'Hanlon⁴⁸, A. Oblakowska-Mucha^{27,38}, V. Obraztsov³⁵, S. Oggero⁴¹, S. Ogilvy⁵¹, O. Okhrimenko⁴⁴, R. Oldeman^{15,e}, G. Onderwater⁶⁵, M. Orlandea²⁹, J.M. Otalora Goicochea², P. Owen⁵³, A. Oyanguren⁶⁴, B.K. Pal⁵⁹, A. Palano^{13,c}, F. Palombo^{21,u}, M. Palutan¹⁸, J. Panman³⁸, A. Papanestis^{49,38}, M. Pappagallo⁵¹, L.L. Pappalardo^{16,f}, C. Parkes⁵⁴, C.J. Parkinson^{9,45}, G. Passaleva¹⁷, G.D. Patel⁵², M. Patel⁵³, C. Patrignani^{19,j}, A. Pazos Alvarez³⁷, A. Pearce⁵⁴, A. Pellegrino⁴¹, M. Pepe Altarelli³⁸, S. Perazzini^{14,d}, E. Perez Trigo³⁷, P. Perret⁵, M. Perrin-Terrin⁶, L. Pescatore⁴⁵, E. Pesen⁶⁶, K. Petridis⁵³, A. Petrolini^{19,j}, E. Picatoste Olloqui³⁶, B. Pietrzyk⁴, T. Pilař⁴⁸, D. Pinci²⁵, A. Pistone¹⁹, S. Playfer⁵⁰, M. Plo Casasus³⁷, F. Polci⁸, A. Poluektov^{48,34}, E. Polycarpo², A. Popov³⁵, D. Popov¹⁰, B. Popovici²⁹, C. Potterat²,

E. Price⁴⁶, J.D. Price⁵², J. Prisciandaro³⁹, A. Pritchard⁵², C. Prouve⁴⁶, V. Pugatch⁴⁴, A. Puig Navarro³⁹, G. Punzi^{23,s}, W. Qian⁴, B. Rachwal²⁶, J.H. Rademacker⁴⁶, B. Rakotomiamanana³⁹, M. Rama¹⁸, M.S. Rangel², I. Raniuk⁴³, N. Rauschmayr³⁸, G. Raven⁴², F. Redi⁵³, S. Reichert⁵⁴, M.M. Reid⁴⁸, A.C. dos Reis¹, S. Ricciardi⁴⁹, S. Richards⁴⁶, M. Rihl³⁸, K. Rinnert⁵², V. Rives Molina³⁶, P. Robbe⁷, A.B. Rodrigues¹, E. Rodrigues⁵⁴, P. Rodriguez Perez⁵⁴, S. Roiser³⁸, V. Romanovsky³⁵, A. Romero Vidal³⁷, M. Rotondo²², J. Rouvinet³⁹, T. Ruf³⁸, H. Ruiz³⁶, P. Ruiz Valls⁶⁴, J.J. Saborido Silva³⁷, N. Sagidova³⁰, P. Sail⁵¹, B. Saitta^{15,e}, V. Salustino Guimaraes², C. Sanchez Mayordomo⁶⁴, B. Sanmartin Sedes³⁷, R. Santacesaria²⁵, C. Santamarina Rios³⁷, E. Santovetti^{24,l}, A. Sarti^{18,m}, C. Satriano^{25,n}, A. Satta²⁴, D.M. Saunders⁴⁶, M. Savrie^{16,f}, D. Savrina^{31,32}, M. Schiller⁴², H. Schindler³⁸, M. Schlupp⁹, M. Schmelling¹⁰, B. Schmidt³⁸, O. Schneider³⁹, A. Schopper³⁸, M.-H. Schune⁷, R. Schwemmer³⁸, B. Sciascia¹⁸, A. Sciubba²⁵, M. Seco³⁷, A. Semennikov³¹, I. Sepp⁵³, N. Serra⁴⁰, J. Serrano⁶, L. Sestini²², P. Seyfert¹¹, M. Shapkin³⁵, I. Shapoval^{16,43,f}, Y. Shcheglov³⁰, T. Shears⁵², L. Shekhtman³⁴, V. Shevchenko⁶³, A. Shires⁹, R. Silva Coutinho⁴⁸, G. Simi²², M. Sirendi⁴⁷, N. Skidmore⁴⁶, T. Skwarnicki⁵⁹, N.A. Smith⁵², E. Smith^{55,49}, E. Smith⁵³, J. Smith⁴⁷, M. Smith⁵⁴, H. Snoek⁴¹, M.D. Sokoloff⁵⁷, F.J.P. Soler⁵¹, F. Soomro³⁹, D. Souza⁴⁶, B. Souza De Paula², B. Spaan⁹, A. Sparkes⁵⁰, P. Spradlin⁵¹, S. Sridharan³⁸, F. Stagni³⁸, M. Stahl¹¹, S. Stahl¹¹, O. Steinkamp⁴⁰, O. Stenyakin³⁵, S. Stevenson⁵⁵, S. Stoica²⁹, S. Stone⁵⁹, B. Storaci⁴⁰, S. Stracka²³, M. Straticiuc²⁹, U. Straumann⁴⁰, R. Stroili²², V.K. Subbiah³⁸, L. Sun⁵⁷, W. Sutcliffe⁵³, K. Swientek²⁷, S. Swientek⁹, V. Syropoulos⁴², M. Szczekowski²⁸, P. Szczypka^{39,38}, D. Szilard², T. Szumlak²⁷, S. T'Jampens⁴, M. Teklishyn⁷, G. Tellarini^{16,f}, F. Teubert³⁸, C. Thomas⁵⁵, E. Thomas³⁸, J. van Tilburg⁴¹, V. Tisserand⁴, M. Tobin³⁹, S. Tolck⁴², L. Tomassetti^{16,f}, S. Topp-Joergensen⁵⁵, N. Torr⁵⁵, E. Tournefier⁴, S. Tourneur³⁹, M.T. Tran³⁹, M. Tresch⁴⁰, A. Tsaregorodtsev⁶, P. Tsoelas⁴¹, N. Tuning⁴¹, M. Ubeda Garcia³⁸, A. Ukleja²⁸, A. Ustyuzhanin⁶³, U. Uwer¹¹, C. Vacca¹⁵, V. Vagnoni^{14,*}, G. Valenti¹⁴, A. Vallier⁷, R. Vazquez Gomez¹⁸, P. Vazquez Regueiro³⁷, C. Vázquez Sierra³⁷, S. Vecchi¹⁶, J.J. Velthuis⁴⁶, M. Veltri^{17,h}, G. Veneziano³⁹, M. Vesterinen¹¹, B. Viaud⁷, D. Vieira², M. Vieites Diaz³⁷, X. Vilasis-Cardona^{36,p}, A. Vollhardt⁴⁰, D. Volyanskyy¹⁰, D. Voong⁴⁶, A. Vorobyev³⁰, V. Vorobyev³⁴, C. Voß⁶², H. Voss¹⁰, J.A. de Vries⁴¹, R. Waldi⁶², C. Wallace⁴⁸, R. Wallace¹², J. Walsh²³, S. Wandernoth¹¹, J. Wang⁵⁹, D.R. Ward⁴⁷, N.K. Watson⁴⁵, D. Websdale⁵³, M. Whitehead⁴⁸, J. Wicht³⁸, D. Wiedner¹¹, G. Wilkinson^{55,38}, M.P. Williams⁴⁵, M. Williams⁵⁶, H.W. Wilschut⁶⁵, F.F. Wilson⁴⁹, J. Wimberley⁵⁸, J. Wishahi⁹, W. Wislicki²⁸, M. Witek²⁶, G. Wormser⁷, S.A. Wotton⁴⁷, S. Wright⁴⁷, K. Wyllie³⁸, Y. Xie⁶¹, Z. Xing⁵⁹, Z. Xu³⁹, Z. Yang³, X. Yuan³, O. Yushchenko³⁵, M. Zangoli¹⁴, M. Zavertyaev^{10,b}, L. Zhang⁵⁹, W.C. Zhang¹², Y. Zhang³, A. Zhelezov¹¹, A. Zhokhov³¹, L. Zhong³, A. Zvyagin³⁸

¹ Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

² Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³ Center for High Energy Physics, Tsinghua University, Beijing, China

⁴ LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵ Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶ CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷ LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸ LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹ Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹⁰ Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹¹ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹² School of Physics, University College Dublin, Dublin, Ireland

¹³ Sezione INFN di Bari, Bari, Italy

¹⁴ Sezione INFN di Bologna, Bologna, Italy

¹⁵ Sezione INFN di Cagliari, Cagliari, Italy

¹⁶ Sezione INFN di Ferrara, Ferrara, Italy

¹⁷ Sezione INFN di Firenze, Firenze, Italy

¹⁸ Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

¹⁹ Sezione INFN di Genova, Genova, Italy

²⁰ Sezione INFN di Milano Bicocca, Milano, Italy

²¹ Sezione INFN di Milano, Milano, Italy

²² Sezione INFN di Padova, Padova, Italy

²³ Sezione INFN di Pisa, Pisa, Italy

²⁴ Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁵ Sezione INFN di Roma La Sapienza, Roma, Italy

²⁶ Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

²⁷ AGH – University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

²⁸ National Center for Nuclear Research (NCBJ), Warsaw, Poland

²⁹ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

³⁰ Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

- ³¹ Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia
³² Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia
³³ Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia
³⁴ Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia
³⁵ Institute for High Energy Physics (IHEP), Protvino, Russia
³⁶ Universitat de Barcelona, Barcelona, Spain
³⁷ Universidad de Santiago de Compostela, Santiago de Compostela, Spain
³⁸ European Organization for Nuclear Research (CERN), Geneva, Switzerland
³⁹ Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
⁴⁰ Physik-Institut, Universität Zürich, Zürich, Switzerland
⁴¹ Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands
⁴² Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands
⁴³ NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine
⁴⁴ Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine
⁴⁵ University of Birmingham, Birmingham, United Kingdom
⁴⁶ H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom
⁴⁷ Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
⁴⁸ Department of Physics, University of Warwick, Coventry, United Kingdom
⁴⁹ STFC Rutherford Appleton Laboratory, Didcot, United Kingdom
⁵⁰ School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
⁵¹ School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
⁵² Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
⁵³ Imperial College London, London, United Kingdom
⁵⁴ School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
⁵⁵ Department of Physics, University of Oxford, Oxford, United Kingdom
⁵⁶ Massachusetts Institute of Technology, Cambridge, MA, United States
⁵⁷ University of Cincinnati, Cincinnati, OH, United States
⁵⁸ University of Maryland, College Park, MD, United States
⁵⁹ Syracuse University, Syracuse, NY, United States
⁶⁰ Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil ^w
⁶¹ Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China ^x
⁶² Institut für Physik, Universität Rostock, Rostock, Germany ^y
⁶³ National Research Centre Kurchatov Institute, Moscow, Russia ^z
⁶⁴ Instituto de Física Corpuscular (IFIC), Universitat de Valencia – CSIC, Valencia, Spain ^{aa}
⁶⁵ KVI – University of Groningen, Groningen, The Netherlands ^{ab}
⁶⁶ Celal Bayar University, Manisa, Turkey ^{ac}

* Corresponding author.

E-mail address: Vincenzo.Vagnoni@bo.infn.it (V. Vagnoni).

^a Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil.

^b P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.

^c Università di Bari, Bari, Italy.

^d Università di Bologna, Bologna, Italy.

^e Università di Cagliari, Cagliari, Italy.

^f Università di Ferrara, Ferrara, Italy.

^g Università di Firenze, Firenze, Italy.

^h Università di Urbino, Urbino, Italy.

ⁱ Università di Modena e Reggio Emilia, Modena, Italy.

^j Università di Genova, Genova, Italy.

^k Università di Milano Bicocca, Milano, Italy.

^l Università di Roma Tor Vergata, Roma, Italy.

^m Università di Roma La Sapienza, Roma, Italy.

ⁿ Università della Basilicata, Potenza, Italy.

^o AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

^p LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

^q Hanoi University of Science, Hanoi, Viet Nam.

^r Università di Padova, Padova, Italy.

^s Università di Pisa, Pisa, Italy.

^t Scuola Normale Superiore, Pisa, Italy.

^u Università degli Studi di Milano, Milano, Italy.

^v Politecnico di Milano, Milano, Italy.

^w Associated to: Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil.

^x Associated to: Center for High Energy Physics, Tsinghua University, Beijing, China.

^y Associated to: Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany.

^z Associated to: Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia.

^{aa} Associated to: Universitat de Barcelona, Barcelona, Spain.

^{ab} Associated to: Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands.

^{ac} Associated to: European Organization for Nuclear Research (CERN), Geneva, Switzerland.