

QUARK BINDING POTENTIAL AND QGP*

SHUKLA PAL^a, APARAJITA BHATTACHARYA^{a,†}, BALLARI CHAKRABARTI^b
RISMITA GHOSH^a^aDepartment of Physics, Jadavpur University, Kolkata 700032, India^bDept. of Physics, Jogamaya Devi College, Kolkata, West Bengal 700026, India*(Received February 27, 2018; accepted September 24, 2018)**Corrected January 2, 2019
according to Erratum 50, 115 (2019)*

The effect of quark–antiquark potential on the dissociation energy and critical screening length of heavy meson such as $b\bar{b}$ and $c\bar{c}$ have been investigated when the respective meson is in Quark–Gluon Plasma (QGP). The different types of interquark potential have been used, which get screened in the QGP medium. The dissociation energy and critical screening length have been studied for both ground and excited states. It has been observed that the form of interquark potential has a substantial effect on the critical screening length when the meson is in QGP. A comparison with other theoretical studies is made.

DOI:10.5506/APhysPolB.49.1865

1. Introduction

It is well-known that the strongly interacting matter with a very high density undergoes a transition to a state of deconfined quarks and gluons. Deconfinement takes place if the color screening dissolves the binding potential between quark and quark or quark and antiquark. The J/ψ or Υ have much smaller radii than the radii of usual mesons and nucleons, due to which the bound state remains unaffected in QGP unless and until the temperature or the density becomes so high that the binding of bound state gets broken. The suppression of J/ψ is one of the signals for quark deconfinement [1]. To investigate quark plasma formation experimentally, it is essential to depend on color screening and deconfinement. In QGP medium, the string tension between a charm c and a charm antiquark \bar{c} disappears and quarks and gluons are deconfined. Only the Coulomb type of color

* Funded by SCOAP³ under Creative Commons License, CC-BY 4.0.

† Corresponding author: pampa@phys.jdvu.ac.in

interaction exists between c and \bar{c} . After the deconfinement of J/ψ , it is impossible to create it by hadronization of plasma. Matsui [2] has discussed the heavy quark J/ψ suppression as a signature of quark–gluon plasma formation. Considering the various attributes of deconfinement test, Satz [3] has concluded that the J/ψ peak in the spectrum of lepton pairs which are emitted during nuclear collisions can give important and required information. At ultra-relativistic energies, dynamical analysis of matter during nuclear collision has been done by Matsui [4]. He also studied J/ψ suppression as a signature of QGP formation. Ruuskanen and Satz [5] have studied the dependence of longitudinal momentum of J/ψ suppression and observed its effects on nuclear collision experiments. Karsch and Petronzio [6] have used the concept of QGP to analyze the nuclear size which depends on J/ψ suppression in heavy-ion collisions and studied the transfer energy related to this heavy-ion collision. Karsch *et al.* [7] have observed the dependence of the dissociation energies, the binding radii and the masses of heavy-quark resonances on the color screening length r_D of the medium and concluded that no binding exists below r_D . Hence, in high-energy heavy-ion collisions, the suppression of J/ψ production may be considered as a symbol for the presence of quark–gluon plasma. Liu and Dong [8] have studied the binding and dissolution for the $c\bar{c}$ and $b\bar{b}$ bound states using different quark potentials in a non-relativistic approximation. They have estimated the critical value of the screening length using the Debye screening effect. They have also estimated the critical temperature T_c of medium. Stubbins [9] has used the generalized variational method to investigate energies for the Yukawa and Hulthen potentials. A number of works have been done on the heavy-meson dissociation in the recent time. Park [10] has investigated the heavy-meson dissociation in light-quark medium and discussed the mechanism of dissociation. He has observed that dissociation length decreases with increase of chemical potential in QGP, whereas in hadronic phase, they behave in an opposite way. Braga *et al.* [11] have studied thermal behavior of $c\bar{c}$ and $b\bar{b}$ by using a holographic model and studied the effect of magnetic field on the thermal spectrum of heavy meson, whereas Blaizot *et al.* [12] investigated the heavy-quark formation, dissociation in QGP in the framework of Langevin equation. They have pointed out that formation of bound states occurs if enough heavy quarks are present in the system, whereas dissociation occurs due to screening of the potential in the plasma. Gau *et al.* [13] studied charmonium $c\bar{c}$ wave function at finite temperature using relativistic Schrödinger equation for spin singlet and triplet. They found that relativistic correction to the J/ψ dissociation temperature in QGP is between 7% to 13%. They have studied wave function S, D states. They have obtained that with temperature both the S and D wave function expands. Adil *et al.* [14] have investigated medium-induced dissociation probability of heavy meson

and found that it is sensitive to the opacity of the quark–gluon plasma and time dependence of its formation and evolution. A comprehensive review of QCD, QGP and heavy quark–meson suppression and production is done by Kisslinger [15].

In the present work, we have investigated the effect of $q\bar{q}$ potential on dissociation energy of $b\bar{b}$ and $c\bar{c}$ mesons in QGP. It is well-known that a number of phenomenological potentials have usually been used to describe the inter-quark potential. The respective potentials get screened in QGP due to interaction with different particles. It is very important to study how the form of the inter-quark potential affects the dissociation energy and, subsequently, affects the critical screening length of the heavy mesons while they are in QGP medium. We use a variational method to study the system and a trial wave function has been used. The dissociation energy and critical screening lengths have been estimated for different form of inter-quark potential. A study with the variation of variational parameter of the trial wave function has also been made. The critical screening length at which dissociation energy vanishes has been extracted for $b\bar{b}$ and $c\bar{c}$ mesons for their ground and excited states.

2. Methodology

The Hamiltonian for a $q\bar{q}$ system in non-relativistic approach can be represented as

$$H(r, T) = \frac{\vec{p}^2}{2m_{re}} + V_{q\bar{q}}(r, T), \tag{1}$$

where m_{re} is the reduced mass of the $q\bar{q}$ system, $V_{q\bar{q}}(r, T)$ is the inter-quark binding potential. The exact nature of the binding potential of $q\bar{q}$ system is not known but a number of phenomenological potentials are suggested which are very successful in describing the binding energy of the system. We have considered four different quark–antiquark potentials such as Cornell potential, Rosener potential, harmonic potential, and a combination of potential for our study. The expression for the potentials run as:

(i) Cornell potential [7]

$$V(r, 0) = -\frac{\alpha_s}{r} + K' r, \tag{2}$$

where K' is the coefficient for confinement and is taken to be 0.92 GeV^2 [16], α_s is the parameter proportional to the strong coupling constant which is 0.471 [7], and r is radius of heavy mesons.

(ii) Rosner potential [17]

$$V(r, 0) = -\frac{A(r^{-\alpha} - 1)}{\alpha} + B \quad (3)$$

with $\alpha = 0.12$, $A = 0.801$ GeV and $B = -0.772$ GeV.

(iii) Harmonic potential [18]

$$V(r, 0) = \delta r^2 \quad (4)$$

with $\delta = 0.08$ GeV³.

(iv) A combination of harmonic, linear and Coulomb potential [19, 20]

$$V(r, 0) = ar^2 + br - c/r \quad (5)$$

with $a = 0.142$ GeV³, $b = 0.465$ GeV² and $c = 0.471$.

In QGP environment of quarks and gluons, the inter-quark potential gets modified due to the color screening so that the $q\bar{q}$ aforesaid potentials can be represented as:

$$V(r, T) = -\frac{Ze^{-\lambda r}}{r} \left[-\frac{\alpha_s}{r} + K' r \right], \quad (6)$$

$$V(r, T) = -\frac{Ze^{-\lambda r}}{r} \left[-\frac{A(r^{-\alpha} - 1)}{\alpha} + B \right], \quad (7)$$

$$V(r, T) = -\frac{Ze^{-\lambda r}}{r} [\delta r^2], \quad (8)$$

$$V(r, T) = -\frac{Ze^{-\lambda r}}{r} [ar^2 + br - c/r], \quad (9)$$

where Z is a constant equal to 1 GeV^{-1} and λ is a temperature-dependent screening parameter. We have parameterized the temperature dependence of λ as $\lambda(T) = \lambda(0)[1 - T/T_c]^{-0.2}$ [21]. The value of $\lambda(0)$ has given input as 0.2 GeV from [7]. Solution of equation (1) with the potential in (2), (3), (4) and (5) will lead to a temperature-dependent binding energy. To get an expression for binding energy, we have considered a trial wave function from the work of Stubbins [9] with a spherical component which runs as

$$\Psi_k = B_k r^k e^{-(\beta/2)} Y_{l,m}(\theta, \phi), \quad (10)$$

where $k = 0, 1, 2, \dots$ and $l = 0, 1, \dots$, and B_k is normalization constant and can be represented as

$$B_k = \left[\frac{\beta^{2k+3}}{(2k+2)!} \right]^{1/2}, \quad (11)$$

where β is variational parameter [9]. Schödinger equation with related eigenvalue can be represented as [7]

$$[H(r, \lambda(T)) - E_{n,l}(\lambda(T))]\Psi_K(r, \lambda(T)) = 0, \tag{12}$$

where n is the principal quantum number and l is the orbital quantum number such that $l \leq (n - 1)$. Dependence of λ on temperature leads to a temperature-dependent eigenenergy.

The dissolution energy is the quantity which accounts the vanishing of bound states. At a fixed value of $\lambda(T)$, the dissolution energy of the bound state can be defined as [7]

$$E_{\text{dis}}^{n,l}(\lambda(T)) = V_{q\bar{q}}(r \rightarrow \infty, \lambda(T)) + E_{n,l}(\lambda(T)). \tag{13}$$

At $r \rightarrow \infty$, equation (13) reduces to

$$E_{\text{dis}}^{n,l}(\lambda(T)) = E_{n,l}(\lambda(T)). \tag{14}$$

The value of dissolution energy is positive for bound states and is negative for continuum, and leads to the condition

$$E_{\text{dis}}^{n,l}(\lambda_c(T)) = 0. \tag{15}$$

Equation (15) gives the critical value of $\lambda(T)$, beyond which for given quantum number, there are no bound states. We have considered first three radial excitation corresponding to J/ψ and Υ for $n = 1, l = 0, \psi'$ and Υ' for $n = 2, l = 0$ and ψ'' and Υ'' for $n = 3, l = 0$ and χ_c and χ_b for $n = 2, l = 1$. We have derived the binding energies of the above-mentioned states and the expressions for binding energies of ground states, first and second excited states for four different potentials are obtained as

(i) Cornell potential

(a) 1s-State

$$E_{\text{BE}} = -2\pi\beta^3 \left[\frac{2K'}{(\beta + \lambda)^3} - \frac{\alpha_s}{(\beta + \lambda)} \right] + \frac{\pi\beta^2}{2m_r}, \tag{16}$$

(b) 2s-State

$$E_{\text{BE}} = -\frac{\pi\beta^5}{3} \left[\frac{12K'}{(\beta + \lambda)^5} - \frac{\alpha_s}{(\beta + \lambda)^3} \right] + \frac{\pi\beta^2}{6m_r}, \tag{17}$$

(c) 3s-State

$$E_{\text{BE}} = -\frac{2\pi\beta^7}{15} \left[\frac{30K'}{(\beta + \lambda)^7} - \frac{\alpha_s}{(\beta + \lambda)^5} \right] + \frac{\pi\beta^2}{10m_r}, \tag{18}$$

(ii) Rosner potential

(a) 1s-State

$$E_{\text{BE}} = -2\pi\beta^3 \left[\frac{(A/\alpha) + C}{(\beta + \lambda)^2} \right] + \frac{\pi\beta^2}{2m_r}, \quad (19)$$

(b) 2s-State

$$E_{\text{BE}} = -\frac{\pi\beta^5}{3} \left[\frac{(A/\alpha) + C}{(\beta + \lambda)^3} \right] + \frac{\pi\beta^2}{6m_r}, \quad (20)$$

(c) 3s-State

$$E_{\text{BE}} = -\frac{2\pi\beta^7}{3} \left[\frac{(A/\alpha) + C}{(\beta + \lambda)^5} \right] + \frac{\pi\beta^2}{10m_r}, \quad (21)$$

(iii) Harmonic potential

(a) 1s-State

$$E_{\text{BE}} = - \left[\frac{12\pi\beta^3 a}{(\beta + \lambda)^4} \right] + \frac{\pi\beta^2}{2m_r}, \quad (22)$$

(b) 2s-State

$$E_{\text{BE}} = -\frac{20\pi\beta^5 a}{(\beta + \lambda)^6} + \frac{\pi\beta^2}{6m_r}, \quad (23)$$

(c) 3s-State

$$E_{\text{BE}} = -\frac{28\pi\beta^7 a}{(\beta + \lambda)^8} + \frac{\pi\beta^2}{10m_r}, \quad (24)$$

and

(iv) A Combination of harmonic, linear and Coulomb potential

(a) 1s-State

$$E_{\text{BE}} = -\frac{2\pi\beta^3}{(\beta + \lambda)} \left[\frac{6a}{(\beta + \lambda)^3} + \frac{2b}{(\beta + \lambda)^2} - c \right] + \frac{\pi\beta^2}{2m_r}, \quad (25)$$

(b) 2s-State

$$E_{\text{BE}} = -\pi\beta^5 \left[\frac{20a}{(\beta + \lambda)^6} \frac{4b}{(\beta + \lambda)^5} - \frac{c}{3(\beta + \lambda)^3} \right] + \frac{\pi\beta^2}{6m_r}, \quad (26)$$

(c) 3s-State

$$E_{\text{BE}} = -2\pi\beta^7 \left[\frac{14a}{(\beta + \lambda)^8} + \frac{2b}{(\beta + \lambda)^7} - \frac{c}{15(\beta + \lambda)^5} \right] + \frac{\pi\beta^2}{10m_r}. \quad (27)$$

We have estimated the critical $\lambda(\lambda_c(T))$ and critical radius using equation (15) with different values of variational parameter β and the results are furnished in Tables I, II, III and IV.

Table I–IV display our results of variation of screening length with variational parameter. Variational technique has provided an effective framework for spectroscopic studies of full-hadron spectrum and a good candidate for investigation of strongly interacting system and gauge theories. The variational method has seen significant success in spectroscopic studies of hadronic system. A number of work have been done in QCD, lattice QCD applying the variational approach [22–24], and found to offer a more efficient method for the determination of nuclear matrix element [25]. Vega *et al.* [26] have studied the Cornell potential using a trial wave function and super symmetric quantum mechanics. The parameters are changed applying successive transformation to obtain the wave function at the origin. Ghalveni *et al.* [27] have studied baryon–meson properties using trial wave function, whereas Chot *et al.* [28] have studied ground state masses with hyperfine interaction in QCD-motivated effective Hamiltonian using a trial wave function. We have used a trial wave function to estimate the binding energy of mesons in QGP and studied the variation of λ_c with variational parameter.

TABLE I

Temperature-dependent critical screening lengths λ_c in GeV with different values of variational parameter (β) in GeV using the Cornell potential.

States		$\beta = 0.1$ [GeV]	$\beta = 0.2$ [GeV]	$\beta = 0.3$ [GeV]	$\beta = 0.4$ [GeV]	$\beta = 0.5$ [GeV]
J/ψ ($n = 1, l = 0$)	λ_c [GeV]	0.6391	0.7048	0.7127	0.6924	0.6577
	r_c [fm]	0.3139	0.2873	0.2806	0.2888	0.3041
ψ' ($n = 2, l = 0$)	λ_c [GeV]	0.3274	0.4437	0.5189	0.5706	0.5975
	r_c [fm]	0.6108	0.4509	0.3854	0.3503	0.3347
ψ'' ($n = 3, l = 0$)	λ_c [GeV]	0.2042	0.2963	0.3632	0.4140	0.4466
	r_c [fm]	0.979	0.6749	0.5506	0.3545	0.4478
χ_c ($n = 2, l = 1$)	λ_c [GeV]	0.3641	0.4395	0.5153	0.5682	0.5771
	r_c [fm]	0.5493	0.4550	0.3881	0.3519	0.3465
Υ ($n = 1, l = 0$)	λ_c [GeV]	0.9341	1.0353	1.0506	1.0315	0.9938
	r_c [fm]	0.2141	0.1932	0.1903	0.1939	0.2012
Υ' ($n = 2, l = 0$)	λ_c [GeV]	0.4331	0.6197	0.7342	0.8244	0.9146
	r_c [fm]	0.4618	0.3227	0.2724	0.2426	0.2187
Υ'' ($n = 3, l = 0$)	λ_c [GeV]	0.2639	0.3910	0.4860	0.5641	0.6197
	r_c [fm]	0.8442	0.5115	0.4115	0.4831	0.3227
χ_b ($n = 2, l = 1$)	λ_c [GeV]	0.5023	0.5893	0.7160	0.8132	0.8551
	r_c [fm]	0.3981	0.3394	0.2793	0.2459	0.2339

TABLE II

Temperature-dependent critical screening lengths λ_c in GeV with different values of variational parameter (β) in GeV using the Rosner potential.

States		$\beta = 0.1$ [GeV]	$\beta = 0.2$ [GeV]	$\beta = 0.3$ [GeV]	$\beta = 0.4$ [GeV]	$\beta = 0.5$ [GeV]
J/ψ ($n = 1, l = 0$)	λ_c [GeV]	1.12	1.536	1.827	2.057	2.248
	r_c [fm]	0.178	0.130	0.109	0.097	0.088
ψ' ($n = 2, l = 0$)	λ_c [GeV]	0.095	0.191	0.287	0.384	0.480
	r_c [fm]	2.105	1.047	0.697	0.521	0.417
ψ'' ($n = 3, l = 0$)	λ_c [GeV]	0.089	0.180	0.271	0.362	0.453
	r_c [fm]	2.247	1.111	0.738	0.552	0.441
χ_c ($n = 2, l = 1$)	λ_c [GeV]	0.07	0.182	0.283	0.379	0.478
	r_c [fm]	2.857	1.099	0.707	0.528	0.418
Υ ($n = 1, l = 0$)	λ_c [GeV]	2.05	2.93	3.55	4.05	4.475
	r_c [fm]	0.0975	0.0682	0.056	0.049	0.044
Υ' ($n = 2, l = 0$)	λ_c [GeV]	0.182	0.347	0.573	0.762	0.955
	r_c [fm]	1.099	0.535	0.349	0.262	0.209
Υ'' ($n = 3, l = 0$)	λ_c [GeV]	0.137	0.279	0.423	0.566	0.708
	r_c [fm]	1.459	0.717	0.473	0.353	0.282
χ_b ($n = 2, l = 1$)	λ_c [GeV]	0.145	0.341	0.51	0.743	0.936
	r_c [fm]	1.379	0.586	0.392	0.269	0.213

TABLE III

Temperature-dependent critical screening lengths λ_c in GeV with different values of variational parameter (β) in GeV using the harmonic potential.

States		$\beta = 0.1$ [GeV]	$\beta = 0.2$ [GeV]	$\beta = 0.3$ [GeV]	$\beta = 0.4$ [GeV]	$\beta = 0.5$ [GeV]
J/ψ ($n = 1, l = 0$)	λ_c [GeV]	0.4893	0.5034	0.4789	0.4371	0.3846
	r_c [fm]	0.4087	0.3973	0.4176	0.4575	0.52
ψ' ($n = 2, l = 0$)	λ_c [GeV]	0.3237	0.4041	0.4409	0.4556	0.4567
	r_c [fm]	0.6178	0.4949	0.4536	0.4389	0.4379
ψ'' ($n = 3, l = 0$)	λ_c [GeV]	0.2297	0.3093	0.3567	0.3863	0.4041
	r_c [fm]	0.8707	0.6466	0.5607	0.5177	0.4949
χ_c ($n = 2, l = 1$)	λ_c [GeV]	0.2923	0.4006	0.4343	0.4518	0.4539
	r_c [fm]	0.6842	0.4992	0.4573	0.4427	0.4406
Υ ($n = 1, l = 0$)	λ_c [GeV]	0.6931	0.7444	0.7476	0.7266	0.6911
	r_c [fm]	0.2885	0.2687	0.2675	0.2752	0.2894
Υ' ($n = 2, l = 0$)	λ_c [GeV]	0.4118	0.5335	0.6004	0.6413	0.6656
	r_c [fm]	0.4857	0.3749	0.3331	0.3118	0.3005
Υ'' ($n = 3, l = 0$)	λ_c [GeV]	0.2759	0.3899	0.4602	0.5108	0.5475
	r_c [fm]	0.7249	0.5129	0.4345	0.3915	0.3653
χ_b ($n = 2, l = 1$)	λ_c [GeV]	0.3511	0.5023	0.5806	0.6305	0.6508
	r_c [fm]	0.5696	0.3981	0.3445	0.3172	0.3073

TABLE IV

Temperature-dependent critical screening lengths λ_c in GeV with different values of variational parameter (β) in GeV using potential (*iv*).

States		$\beta = 0.1$ [GeV]	$\beta = 0.2$ [GeV]	$\beta = 0.3$ [GeV]	$\beta = 0.4$ [GeV]	$\beta = 0.5$ [GeV]
J/ψ ($n = 1, l = 0$)	λ_c [GeV]	0.6656	0.7095	0.7014	0.6679	0.6204
	r_c [fm]	0.3005	0.2819	0.2851	0.2994	0.3224
ψ' ($n = 2, l = 0$)	λ_c [GeV]	0.3910	0.5075	0.5759	0.6194	0.6464
	r_c [fm]	0.5115	0.3941	0.3473	0.3229	0.3094
ψ'' ($n = 3, l = 0$)	λ_c [GeV]	0.2599	0.3626	0.4319	0.4813	0.5185
	r_c [fm]	0.7695	0.5516	0.4631	0.4155	0.3857
χ_c ($n = 2, l = 1$)	λ_c [GeV]	0.3612	0.5023	0.5721	0.6162	0.6435
	r_c [fm]	0.5537	0.3081	0.3496	0.3245	0.3108
Υ ($n = 1, l = 0$)	λ_c [GeV]	0.9183	0.9822	0.9767	0.9404	0.8878
	r_c [fm]	0.2178	0.2036	0.2048	0.2127	0.2253
Υ' ($n = 2, l = 0$)	λ_c [GeV]	0.4928	0.6683	0.7784	0.8551	0.9112
	r_c [fm]	0.4058	0.2997	0.2569	0.2339	0.2195
Υ'' ($n = 3, l = 0$)	λ_c [GeV]	0.3169	0.4507	0.5510	0.6271	0.6877
	r_c [fm]	0.6311	0.4437	0.3629	0.3189	0.2908
χ_b ($n = 2, l = 1$)	λ_c [GeV]	0.4632	0.6271	0.7677	0.8325	0.8973
	r_c [fm]	0.4318	0.3189	0.2605	0.2402	0.2228

3. Conclusions

In the present work, we have investigated the dissolution energy of heavy quarkonia $b\bar{b}$ and $c\bar{c}$ considering the effect of QGP medium in the interquark potential of the heavy mesons. We have used variational method to get the expression for energy and studied the critical parameter with the variation of the variational parameter of the trial wave function. We have also suggested an empirical form of temperature-dependent screening parameter by the relation $\lambda(T) = \lambda(0)[1 - T/T_c]^{-0.2}$ with the critical exponent 0.2 in our work. The study of critical phenomena and corresponding scaling behavior of phase transition get a new impetus with the recent experimental development of studying low temperature physics. The quasi-particle effective mass have been studied by parameterizing the behavior as $m^* = m(0)[1 - \frac{T}{T_c}]^\beta$ [21]. Usually, the critical point is reached by tuning the thermodynamic parameters. The renormalization group theory does not restrict the continuous variation of critical exponent which leads to the weak universality. Crystal behavior shows asymmetry in critical exponent (α) with $(\alpha) = -0.2 \pm 0.3$. The critical exponent for $T < T_c$ for fluid varies from 0.1 to 0.2 [29]. The mean-field prediction of $(\alpha) = 0.5$ does not match with experimental value of 0.3 for fluid. The critical exponent is suggested to be 0.1 for CO₂, whereas

for Xe, the value is ~ 0.2 which does not violate the Rushbrooke or Griffith inequality. We have used critical exponent as treating the QGP as a fluid and have studied the phase transition to estimate the critical screening length for heavy-meson dissociation in QGP. The variation of dissociation energy with the variational parameter have been studied for ground states, first and second excited states of the heavy mesons. We have estimated the critical values of the screening parameters and critical radii for the states considering variational parameter β varying from 0.1 GeV to 0.5 GeV. Liu *et al.* [8] have studied the quark binding potential in QGP for various values of power of potential between quark and antiquark and studied J/ψ suppression. They have estimated the screening masses and Debye screening radii with temperature-dependent different types of potentials and studied the values with different values of screening parameter. In the current work, we have studied screening lengths and screening radii with the variation of variable parameter. Variational method is a useful tool to estimate the ground state energies and also excited states. It may be mentioned that the variational method together with physically motivated trial wave function provide a powerful tool to study the systems under extreme condition and can be more robust in situation where it is difficult to determine a good unperturbed Hamiltonian. In the current work, it has been observed that the value of critical screening lengths with variational parameter $\beta = 0.2$ GeV estimated in the present work with Cornell potential agree well with the estimation of Liu *et al.* [8] with the value of screening parameter equal to 1.0 with model (i). More comparative study will be done with this variational approach in our future work.

The authors are thankful to the University Grants Commission, New Delhi, India, for financial support.

REFERENCES

- [1] T. Matsui, H. Satz, *Phys. Lett. B* **178**, 416 (1986).
- [2] T. Matsui, *Z. Phys. C* **38**, 245 (1988).
- [3] H. Satz, *Nucl. Phys. A* **488**, 511c (1988).
- [4] T. Matsui, *Nucl. Phys. A* **488**, 535c (1988).
- [5] P.V. Ruuskanen, H. Satz, *Z. Phys. C* **37**, 623 (1988).
- [6] F. Karsch, R. Petronzio, *Phys. Lett. B* **212**, 255 (1988).
- [7] F. Karsch, M.T. Mehr, H. Satz, *Z. Phys. C* **37**, 617 (1988).
- [8] B. Liu, Y.B. Dong, *Commun. Theor. Phys.* **26**, 425 (1996).
- [9] C. Stubbins, *Phys. Rev. A* **48**, 220 (1993).

- [10] C. Park, *Phys. Rev. D* **81**, 045009 (2010).
- [11] N.R.F. Braga, L.F. Ferreira, *Phys. Lett. B* **783**, 186 (2018).
- [12] J.P. Blaizot, D. De Boni, P. Faccioli, G. Garberoglio, *Nucl. Phys. A* **946**, 49 (2016).
- [13] X. Guo, S. Shi, P. Zhuang, *Phys. Lett. B* **718**, 143 (2012).
- [14] A. Adil, I. Vitev, *Phys. Lett. B* **649**, 139 (2007).
- [15] L.S. Kisslinger, *Int. J. Mod. Phys. A* **173**, 00083 (2017).
- [16] K.K. Seth, arXiv:0912.2776 [hep-ex].
- [17] A.K. Grant, J.L. Rosner, E. Rynes, *Phys. Rev. D* **47**, 1981 (1993).
- [18] M.R. Shojaei, A.A. Rajabi, *Mod. Phys. Lett. A* **23**, 3411 (2009).
- [19] R. Kumar, D. Kumar, F. Chand, *Proceedings of DAE Symp. Nucl. Phys.* **57**, 664 (2012).
- [20] M.R. Shojaei, H.T. Anbaran, *Appl. Phys. Res.* **2**, 148 (2010).
- [21] A. Chandra, A. Bhattacharya, B. Chakrabarti, *J. Mod. Phys.* **4**, 945 (2013).
- [22] D.S. Roberts, W. Kamleh, D.B. Leinweber, arXiv:1304.0325 [hep-lat].
- [23] I.I. Kogan, A. Kovner, *Phys. Rev. D* **52**, 3719 (1995).
- [24] T.A. DeGrand, R.D. Loft, *Comput. Phys. Commun.* **65**, 84 (1991).
- [25] J. Dragos *et al.*, *Phys. Rev. D* **94**, 074505 (2016).
- [26] A. Vega, J. Flores, *Pramana — J. Phys.* **87**, 1 (2016).
- [27] Z. Ghalehovi, A.A. Rajabi, *Acta. Phys. Pol. B* **42**, 1849 (2011).
- [28] H.M. Choi *et al.*, *Acta. Phys. Pol. B Proc. Suppl.* **6**, 281 (2013).
- [29] NPTEL-Phase II, Advanced Statistical Physics.