Erratum: CP violation in a model with Higgs triplets

Ting-Kuo Chen,[a,b] **Cheng-Wei Chiang**[a,c] **and Kei Yagyu**[d]

[a]*Department of Physics and Center for Theoretical Physics, National Taiwan University,*
Taipei, Taiwan 10617, R.O.C.

[b]*Department of Physics, University of Wisconsin-Madison,*
*Madison, WI 53706, U.S.A.*

[c]*Physics Division, National Center for Theoretical Sciences,*
Taipei, Taiwan 10617, R.O.C.

[d]*Department of Physics, Osaka University,*
Toyonaka, Osaka 560-0043, Japan

E-mail: tchen463@wisc.edu, chengwei@phys.ntu.edu.tw,
yagyu@het.phys.sci.osaka-u.ac.jp

Figure 1. Branching ratios of the most dominant decay channels of H_1 (upper row) and H_2 (lower row) for benchmark 1 (left column) and benchmark 2 (right column) as a function of $|\Im\sigma_4|$. The gray hatched regions are excluded by the eEDM constraint at 90% CL, and the brown hatched regions by the other theoretical and collider measurement constraints at 95% CL. The bound set by the eEDM constraint for benchmark 2 is beyond the plotting range.

In figure 7, each of the upper two rows of plots is missing one curve, the branching ratio of the $t\bar{t}$ mode. The updated plots are given in figure 1.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.