

Lepton mass correction in partial wave analyses of charmed meson semi-leptonic decays*

Han Zhang (张晗)¹ Bai-Cian Ke (柯百谦)^{1†} Yao Yu (余耀)^{2‡} En Wang (王恩)¹

¹School of Physics and Microelectronics, Zhengzhou University, Zhengzhou 450001, China

²Chongqing University of Posts & Telecommunications, Chongqing, 400065, China

Abstract: We derive a parameterization formula for the partial wave analyses of charmed meson semi-leptonic decays while considering the effects of lepton mass. Because the proposed super-tau-charm factory will reach a significantly enhanced luminosity and BESIII is collecting new $\psi(3770) \rightarrow D\bar{D}$ data, our results will help improve the measurement precision of future partial wave analyses of charmed meson semi-muonic decays.

Keywords: partial wave analysis, charmed meson, semi-leptonic decay

DOI: 10.1088/1674-1137/acc642

I. INTRODUCTION

The standard model, as a theory of elementary particle interactions, has achieved great success in describing experimental data with various energies. Nevertheless, the ~ 2 GeV masses of charmed mesons place them in the region where perturbative QCD is not applicable, which raises the challenges faced by both theory and experiments. In recent years, standard model testing with high precision measurements has become one of the hottest topics in the charm sector.

The semi-leptonic decays of charmed mesons, in which hadronic and weak currents can be well separated, provide a clean platform for studying the mechanism of the c quark to $d(s)$ quark transition and play an important role in understanding strong and weak interactions. Their partial decay width accesses the product of the hadronic form factor, which describes the strong-interaction in the hadronic current connecting initial and final hadrons, and the Cabibbo-Kobayashi-Maskawa matrix element $|V_{cs}|$ or $|V_{cd}|$, which parameterizes the weak interaction between different quark flavors. Partial wave analyses of the four-body semi-leptonic decays of charmed mesons allow us

to extract the form factors in the $D \rightarrow V\ell^+\nu_\ell$ and $D \rightarrow S\ell^+\nu_\ell$ transitions (where $\ell = e, \mu$, V and S denote vector and scalar mesons, respectively). $K^*(892)^{(0)}$ resonance has been studied in the $D^{0(+)} \rightarrow K^{0(-)}\pi^+e^+\nu_e$ decay by the CLEO, BABAR, and BESIII collaborations [1–4]. BESIII has also studied ρ^- resonance in $D^0 \rightarrow \pi^-\pi^0e^+\nu_e$ [5], and ρ^0 , ω , and $f_0(500)$ resonances in $D^+ \rightarrow \pi^+\pi^-e^+\nu_e$ [5]. In the near future, more partial wave analyses are expected to be performed with high-statistics datasets, including $D_{(s)}$ semi-muonic decays¹⁾.

However, lepton mass is neglected in the parameterization formula used for the amplitude analyses of charmed meson semi-leptonic decays. This should be manageable in the case of semi-electronic decays, but will cause bias in semi-muonic decays, which downgrades the advantage of high-statistics from the new data and conflicts with the purpose of precision measurement. In this study, we derive the formula while considering the mass of leptons based on Refs. [6, 7]. The results are presented in the format used in experimental analyses; therefore, researchers can easily adopt our results to their experimental analyses. Charge-conjugated decay modes are implied throughout this paper.

Received 22 February 2023; Accepted 22 March 2023; Published online 23 March 2023

* HZ and BCK were supported in part by the Joint Large-Scale Scientific Facility Fund of the National Natural Science Foundation of China (NSFC) (11875054, 12192263) and the Chinese Academy of Sciences (U2032104); YY was supported in part by the NSFC (11905023, 12047564, 12147102), the Natural Science Foundation of Chongqing, China (cstc2020jcyj-msxmX0555), and the Science and Technology Research Program of Chongqing Municipal Education Commission (KJQN202200605, KJQN202200621); EW and HZ were supported in part by the Natural Science Foundation of Henan, China (222300420554, 232300421140), the Project of Youth Backbone Teachers of Colleges and Universities of Henan Province, China (2020GGJS017), the Youth Talent Support Project of Henan Province, China (2021HYTP002), and the Open Project of Guangxi Key Laboratory of Nuclear Physics and Nuclear Technology (NLK2021-08)

[†] E-mail: baiciank@ihep.ac.cn

[‡] E-mail: yuyao@cqpt.edu.cn

1) At present, there is no partial wave analyses of $D_{(s)}$ semi-muonic decays reported yet because low statistics and high-level background caused by μ - π misidentification.



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

II. FORMALISM

First, we define the kinematic variables and discuss their relations. The four-body semi-leptonic decay $D \rightarrow M_1 M_2 \ell^+ \nu_\ell$ is considered, where D is the parent meson, $M_{1(2)}$ is the product meson, and $\ell = e, \mu$. The momentum four-vectors and invariant masses are denoted by p and m , respectively. For convenience, the independent four-vector combinations are defined as

$$\begin{aligned} P^\mu &= p_{M_1}^\mu + p_{M_2}^\mu, & Q^\mu &= p_{M_1}^\mu - p_{M_2}^\mu, \\ L^\mu &= p_\ell^\mu + p_{\nu_\ell}^\mu, & N^\mu &= p_\ell^\mu - p_{\nu_\ell}^\mu. \end{aligned} \quad (1)$$

A four-body decay can be uniquely described via kinematic parameterization with five variables (besides spin). The squared masses of the hadronic system $M_1 M_2$ and leptonic system $\ell^+ \nu_\ell$ are chosen as two of the five variables,

$$s_M = P^2, \quad s_L = L^2, \quad (2)$$

and the following relations can be easily derived:

$$Q^2 = 2m_{M_1}^2 + 2m_{M_2}^2 - s_M, \quad N^2 = 2m_\ell^2 + 2m_{\nu_\ell}^2 - s_L, \quad (3)$$

$$L \cdot P = \frac{m_D^2 - s_M - s_L}{2}, \quad L \cdot N = m_\ell^2 - m_{\nu_\ell}^2, \quad P \cdot Q = m_{M_1}^2 - m_{M_2}^2. \quad (4)$$

The other three variables are chosen as the angle between the M_2 three-momentum and the D direction in the $M_1 M_2$ rest frame (θ_M), the angle between ν_ℓ and the D direction in the $\ell^+ \nu_\ell$ rest frame (θ_L), and the angle between the two decay planes (ϕ)¹⁾. The angles θ_M , θ_L , and ϕ are illustrated in Fig. 1. The various relationships between scalar-product invariants can be written as

$$\begin{aligned} L \cdot Q &= L \cdot P \frac{m_{M_1}^2 - m_{M_2}^2}{s_M} + X \beta_M \cos \theta_M, \\ N \cdot P &= L \cdot P \frac{m_\ell^2 - m_{\nu_\ell}^2}{s_L} + X \beta_L \cos \theta_L, \\ N \cdot Q &= L \cdot P \beta_M \beta_L \cos \theta_M \cos \theta_L + \frac{m_\ell^2 - m_{\nu_\ell}^2}{s_L} X \beta_M \cos \theta_M \\ &\quad + \frac{m_{M_1}^2 - m_{M_2}^2}{s_M} X \beta_L \cos \theta_L + L \cdot P \frac{m_{M_1}^2 - m_{M_2}^2}{s_M} \frac{m_\ell^2 - m_{\nu_\ell}^2}{s_L} \\ &\quad - \sqrt{s_M} \sqrt{s_L} \beta_M \beta_L \sin \theta_M \sin \theta_L \cos \phi, \\ \epsilon_{\mu\nu\rho\sigma} L^\mu N^\nu P^\rho Q^\sigma &= X \sqrt{s_M} \sqrt{s_L} \beta_M \beta_L \sin \theta_M \sin \theta_L \sin \phi, \end{aligned} \quad (5)$$




Fig. 1. (color online) Definition of angles θ_L , θ_M , and ϕ in the cascade decay $D \rightarrow M_1 M_2 \ell^+ \nu_\ell$.

where β_M is the three-momentum modulus of the meson in the center-of-mass frame of the meson-meson system, β_L the three-momentum modulus of the lepton in the center-of-mass frame of the lepton-neutrino system, and X an element of phase space,

$$\begin{aligned} \beta_M &= \sqrt{(s_M - m_+^2)(s_M - m_-^2)} / s_M, \\ \beta_L &= \sqrt{(s_L - m_{L^+}^2)(s_L - m_{L^-}^2)} / s_L, \\ X &= \sqrt{m_D^4 + s_L^2 + s_M^2 - 2s_D m_L^2 - 2s_M m_D^2 - 2s_M s_L} / 2, \end{aligned} \quad (6)$$

with

$$\begin{aligned} m_{L^+} &= m_\ell + m_{\nu_\ell}; & m_{L^-} &= m_\ell - m_{\nu_\ell}; \\ m_+ &= m_{M_1} + m_{M_2}; & m_- &= m_{M_1} - m_{M_2}. \end{aligned} \quad (7)$$

Compared to Refs. [6, 7], we do not neglect the lepton mass.

Next, from the effective Hamiltonian at the quark level for $D \rightarrow M_1 M_2 \ell^+ \nu_\ell$, the decay amplitude is given by

$$\begin{aligned} \mathcal{A}(D \rightarrow M_1 M_2 \ell^+ \nu_\ell) &= \frac{G_F}{\sqrt{2}} V_{q_1 q_2} \langle M_2 M_1 | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | D \rangle \bar{u}(p_\ell) \gamma^\mu (1 - \gamma_5) v(p_{\nu_\ell}), \end{aligned} \quad (8)$$

where G_F is the Fermi constant, and $V_{q_1 q_2}$ is the element of the Cabibbo-Kobayashi-Maskawa matrix. The hadronic matrix element can be written in terms of four form factors, w_\pm , r , and h , which are defined by

$$\begin{aligned} \langle M_2 M_1 | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | D \rangle &= h \epsilon^{\mu\nu\alpha\beta} p_D^\nu P^\alpha Q^\beta + i r L^\mu + i w_+ P^\mu + i w_- Q^\mu, \end{aligned} \quad (9)$$

1) ϕ in this paper is defined as $-\phi$ in [6, 7]

where the form factors w_{\pm} , r , and h are functions of s_M , s_L , and $\cos\theta_M$, and $\epsilon^{\mu\nu\alpha\beta}$ is the Levi-Civita symbol.

The differential decay rate takes the form

$$d\Gamma = \frac{G_F^2 |V_{q_1 q_2}|^2}{(4\pi)^6 m_D^3} X \beta_M I(s_M, s_L, \theta_M, \theta_L, \phi) \times ds_M ds_L d\cos\theta_M d\cos\theta_L d\phi. \quad (10)$$

To study the structure of the hadron system, that is, the form factors of the $M_1 M_2$ system, the decay intensity I is decomposed with respect to θ_L and ϕ , which is written as

$$I = I_1 + I_2 \cos 2\theta_L + I_3 \sin^2 \theta_L \cos 2\phi + I_4 \sin 2\theta_L \cos \phi + I_5 \sin \theta_L \cos \phi + I_6 \cos \theta_L + I_7 \sin \theta_L \sin \phi + I_8 \sin 2\theta_L \sin \phi + I_9 \sin^2 \theta_L \sin 2\phi, \quad (11)$$

where $I_{1,2,\dots,9}$ depend only on s_M , s_L , and ϕ . We can further express $I_{1,2,\dots,9}$ in terms of form factors,

$$\begin{aligned} I_1 &= \frac{1}{4}(2 - \beta_L)\beta_L |F_1|^2 + \left(\frac{\beta_L}{2} - \frac{\beta_L^2}{8}\right) \sin^2 \theta_M (|F_2|^2 + |F_3|^2) \\ &\quad + \frac{1}{2}(1 - \beta_L)\beta_L |F_4|^2, \\ I_2 &= -\frac{\beta_L^2}{4} \left[|F_1|^2 - \frac{1}{2} \sin^2 \theta_M (|F_2|^2 + |F_3|^2) \right], \\ I_3 &= -\frac{\beta_L^2}{4} \left[\sin^2 \theta_M (|F_2|^2 - |F_3|^2) \right], \\ I_4 &= \frac{\beta_L^2}{2} \sin \theta_M \operatorname{Re}(F_1 F_2^*), \\ I_5 &= -\beta_L \sin \theta_M \left\{ \operatorname{Re}(F_1 F_3^*) - (1 - \beta_L) \operatorname{Re}(F_2 F_4^*) \right\}, \\ I_6 &= -\beta_L \sin^2 \theta_M \operatorname{Re}(F_2 F_3^*) - \beta_L (1 - \beta_L) \operatorname{Re}(F_1 F_4^*), \\ I_7 &= \beta_L \sin \theta_M \left\{ \operatorname{Im}(F_1 F_2^*) + (1 - \beta_L) \operatorname{Im}(F_3 F_4^*) \right\}, \\ I_8 &= -\frac{\beta_L^2}{2} \sin \theta_M \operatorname{Im}(F_1 F_3^*), \\ I_9 &= \frac{\beta_L^2}{2} \sin^2 \theta_M \operatorname{Im}(F_2 F_3^*), \end{aligned} \quad (12)$$

where $F_{1,2,3,4}$ are the form factors

$$\begin{aligned} F_1 &= X w_+ + (\beta_M P \cdot L \cos \theta_M + \frac{m_+ m_-}{s_M} X) w_-, \\ F_2 &= \beta_M \sqrt{s_M} \sqrt{s_L} w_-, \\ F_3 &= X \beta_M \sqrt{s_M} \sqrt{s_L} h, \\ F_4 &= s_L r + P \cdot L w_+ + (X \beta_M \cos \theta_M + \frac{m_+ m_-}{s_M} P \cdot L) w_-. \end{aligned} \quad (13)$$

For the purpose of discussing the angular momentum of $M_1 M_2$, for example, S - and P -waves, the partial wave expansions in spherical harmonics for the form factors $F_{1,2,3,4}$ are written as

$$\begin{aligned} F_1(s_M, s_L, \cos \theta_M) &= \sum_{l=0}^{\infty} F_{1l}(s_M, s_L) P_l(\cos \theta_M), \\ F_2(s_M, s_L, \cos \theta_M) &= \sum_{l=1}^{\infty} \frac{1}{\sqrt{l(l+1)}} F_{2l}(s_M, s_L) \frac{dP_l(\cos \theta_M)}{d\cos \theta_M}, \\ F_3(s_M, s_L, \cos \theta_M) &= \sum_{l=1}^{\infty} \frac{1}{\sqrt{l(l+1)}} F_{3l}(s_M, s_L) \frac{dP_l(\cos \theta_M)}{d\cos \theta_M}, \\ F_4(s_M, s_L, \cos \theta_M) &= \sum_{l=0}^{\infty} F_{4l}(s_M, s_L) P_l(\cos \theta_M). \end{aligned} \quad (14)$$

Moreover, the decay $D \rightarrow M_1 M_2 \ell^+ \nu_\ell$ may occur via intermediate states, such as vector or scalar mesons, which provides information about the intermediate resonances [8, 9]. The amplitudes of $D \rightarrow (S \rightarrow M_1 M_2) \ell^+ \nu_\ell$ and $D \rightarrow (V \rightarrow M_1 M_2) \ell^+ \nu_\ell$ can be given by

$$\begin{aligned} \mathcal{A}(D \rightarrow (V \rightarrow M_1 M_2) \ell^+ \nu_\ell) &= \langle V | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | D \rangle \epsilon \cdot Q g_{V M_1 M_2} D_{FV} \bar{u}(p_\ell) \gamma^\mu (1 - \gamma_5) v(p_\nu), \\ \mathcal{A}(D \rightarrow (S \rightarrow M_1 M_2) \ell^+ \nu_\ell) &= \langle S | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | D \rangle g_{S M_1 M_2} D_{FS} \bar{u}(p_\ell) \gamma^\mu (1 - \gamma_5) v(p_\nu), \end{aligned} \quad (15)$$

where

$$\begin{aligned} \langle V | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | D \rangle &= -\epsilon_{\mu\nu\alpha\beta} \epsilon^{\nu*} P_D^\alpha P_D^\beta \frac{2V_0(s_L)}{m_V + m_D} - i \left(\epsilon_\mu^* - \frac{\epsilon^* \cdot L}{L^2} L_\mu \right) (m_V + m_D) A_1(s_L) \\ &\quad + i \left(P_{D\mu} + P_\mu - \frac{m_D^2 - m_V^2}{L^2} L_\mu \right) \epsilon^* \cdot L \frac{A_2(s_L)}{m_V + m_D} \\ &\quad - i \frac{2m_V \epsilon^* \cdot L}{L^2} L_\mu A_0(s_L), \\ \langle S | \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2 | D \rangle &= i(f^+(s_L) P_\mu + f^-(s_L) L_\mu), \end{aligned} \quad (16)$$

with

$$\sum \epsilon^{\mu*} \epsilon^\nu = -g^{\mu\nu} + \frac{P^\mu P^\nu}{P^2},$$

$$\begin{aligned} \mathcal{A}(D \rightarrow M_1 M_2 \ell^+ \nu_\ell) &= \mathcal{A}(D \rightarrow (V \rightarrow M_1 M_2) \ell^+ \nu_\ell) \\ &\quad + \mathcal{A}(D \rightarrow (S \rightarrow M_1 M_2) \ell^+ \nu_\ell). \end{aligned} \quad (17)$$

Here, $f^\pm(s_L)$ is the $D \rightarrow S$ form factor, $A_{0,1,2}$ are the

$D \rightarrow V$ axial-vector form factors, and V_0 is the $D \rightarrow V$ vector form factor [10, 11]. Finally, we can obtain F_{1-4} in the helicity basis (for S - and P -waves only),

$$\begin{aligned}
F_1(s_M, s_L, \cos\theta_M) &= X f^+(s_L) g_{SM_1 M_2} D_{FS} \\
&\quad + \cos\theta_M \beta_M g_{VM_1 M_2} D_{FV} \sqrt{s_L s_M} H_0(s_L), \\
F_2(s_M, s_L, \cos\theta_M) &= \frac{1}{2} \beta_M g_{VM_1 M_2} D_{FV} \sqrt{s_L s_M} \\
&\quad \times (H_+(s_L) + H_-(s_L)), \\
F_3(s_M, s_L, \cos\theta_M) &= \frac{1}{2} \beta_M g_{VM_1 M_2} D_{FV} \sqrt{s_L s_M} \\
&\quad \times (H_+(s_L) - H_-(s_L)), \\
F_4(s_M, s_L, \cos\theta_M) &= s_L f^-(s_L) g_{SM_1 M_2} D_{FS} \\
&\quad + 2 \cos\theta_M \beta_M g_{VM_1 M_2} D_{FV} \sqrt{s_L s_M} H_t(s_L),
\end{aligned} \tag{18}$$

with

$$\begin{aligned}
H_0(s_L) &= \frac{1}{\sqrt{s_L s_M}} \left[P \cdot L(m_V + m_D) A_1(s_L) - 2 \frac{X^2}{m_V + m_D} A_2(s_L) \right] \\
H_{\pm}(s_L) &= (m_V + m_D) A_1(s_L) \mp \frac{2X}{m_V + m_D} V_0(s_L), \\
H_t(s_L) &= \frac{X}{\sqrt{s_L}} [A_0(s_L) + A_1(s_L) + A_2(s_L)],
\end{aligned} \tag{19}$$

where $g_{SM_1 M_2} (g_{VM_1 M_2})$ is the coupling constant, $D_{FS} (D_{FV})$ is derived from the propagator for $S (V)$. In the case of Breit-Wigner lineshapes, $D_{FS} = 1/(s_M - m_S^2 + i m_S \Gamma_S)$ and

$D_{FV} = 1/(s_M - m_V^2 + i m_V \Gamma_V)$, or $D_{FS} = 1/(s_M - m_S^2 + i \frac{s_M}{m_S} \Gamma_S)$ and $D_{FV} = 1/(s_M - m_V^2 + i \frac{s_M}{m_V} \Gamma_V)$ if decay width $\Gamma_S (\Gamma_V)$ is not negligible [12]. For the scalar meson $f_0(980)$, it couples to the channels $K\bar{K}$ and $\pi\pi$ strongly. Likewise, $a_0(980)$ couples to $K\bar{K}$ and $\pi\eta$ strongly. One needs to replace the usual Breit-Wigner lineshape by the Flatté formula [13] to take into account the coupled channel effect. For the broad $\rho(770)$ resonance, it is customary to adapt the Gounaris-Sakurai model [14] to account for the $\pi\pi$ rescattering.

III. DISCUSSION AND CONCLUSION

In this study, we derive the parameterization formula of a four-body semi-leptonic decay while considering the effects of lepton mass and express the differential decay width in the format used in partial wave analyses. We can obtain the parameterization formula used in Refs. [3–5], which neglects lepton mass (or set $\beta_L = 1$); however, while this does not significantly influence semi-electronic decays, neglecting the lepton mass can result in a bias of up to $\sim 1\%$ in partial wave analyses for charmed meson semi-muonic decays.

BESIII is accumulating data samples with an integrated luminosity of 20 fb^{-1} at a center-of-mass energy of 3.773 GeV (for D^0 and D^\pm mesons) and has collected 7.33 fb^{-1} data samples at $4.128\text{--}4.226 \text{ GeV}$ (for D_s^\pm mesons) [15]. In addition, the proposed super-tau-charm factory will be able to reach significantly enhanced luminosities. These aim at testing the standard model with high precision in the charm sector; however, precise theoretical parameterization is also needed. With the correction for lepton mass presented in this paper, partial wave analyses can be performed to study the form factors in the $D_{(s)} \rightarrow V \mu^+ \nu_\mu$ and $D_{(s)} \rightarrow S \mu^+ \nu_\mu$ decays more precisely.

References

- [1] R. A. Briere *et al.* (CLEO Collaboration), *Phys. Rev. D* **81**, 112001 (2010)
- [2] P. del Amo Sanchez *et al.* (BABAR Collaboration), *Phys. Rev. D* **83**, 072001 (2011)
- [3] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **94**, 032001 (2016)
- [4] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **99**, 011103(R) (2019)
- [5] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **122**, 062001 (2019)
- [6] C. L. Y. Lee, M. Lu, and M. B. Wise, *Phys. Rev. D* **46**, 5040 (1992)
- [7] A. Pais and S. B. Treiman, *Phys. Rev.* **168**, 1858 (1968)
- [8] G. Y. Wang, L. Roca, E. Wang *et al.*, *Eur. Phys. J. C* **80**(5), 388 (2020)
- [9] L. R. Dai, X. Zhang, and E. Oset, *Phys. Rev. D* **98**, 036004 (2018)
- [10] A. Ali, G. Kramer, and C. D. Lu, *Phys. Rev. D* **58**, 094009 (1998)
- [11] H. Y. Cheng, C. Y. Cheung, and C. W. Hwang, *Phys. Rev. D* **55**, 1559 (1997)
- [12] A. R. Bohm and Y. Sato, *Phys. Rev. D* **71**, 085018 (2005)
- [13] Z. L. Wang, and B. S. Zou, *Eur. Phys. J. C* **82**, 509 (2022)
- [14] G. J. Gounaris and J. J. Sakurai, *Phys. Rev. Lett.* **21**, 244 (1968)
- [15] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **44**, 040001 (2020)