



Generalised CP and trimaximal TM_1 lepton mixing in S_4 family symmetry

Cai-Chang Li, Gui-Jun Ding*

Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China

Received 20 December 2013; received in revised form 18 January 2014; accepted 5 February 2014

Available online 13 February 2014

Abstract

We construct two flavor models based on S_4 family symmetry and generalised CP symmetry. In both models, the S_4 family symmetry is broken down to the Z_2^{SU} subgroup in the neutrino sector, as a consequence, the trimaximal TM_1 lepton mixing is produced. Depending on the free parameters in the flavon potential, the Dirac CP is predicted to be either conserved or maximally broken, and the Majorana CP phases are trivial. The two models differ in the neutrino sector. The flavon fields are involved in the Dirac mass terms at leading order in the first model, and the neutrino mass matrix contains three real parameters such that the absolute neutrino masses are fixed. Nevertheless, the flavon fields enter into the Majorana mass terms at leading order in the second model. The leading order lepton mixing is of the tri-bimaximal form which is broken down to TM_1 by the next to leading order contributions.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Introduction

In the past years, the Daya Bay [1], RENO [2] and Double Chooz [3] experiments, together with the long-baseline experiments T2K [4] and MINOS [5], have provided an accurate determination of the last unknown lepton mixing angle θ_{13} , with the latest central value measured by Daya Bay being $\theta_{13} \simeq 8.7^\circ$ [6]. The measurement of the reactor angle excluded many neutrino mass models, and led to new model building strategies based on family symmetries [7–10]. So far all the three lepton mixing angles and both mass-squared differences Δm_{sol}^2 and Δm_{atm}^2 have

* Corresponding author.

E-mail addresses: lcc0915@mail.ustc.edu.cn (C.-C. Li), dinggj@ustc.edu.cn (G.-J. Ding).

been measured to reasonably good accuracy. However, barely nothing is known on the leptonic CP phases, which contain one Dirac phase δ_{CP} and two Majorana phases α_{21} and α_{31} . The global analysis of the current neutrino oscillation data gives that the 3σ range of δ_{CP} is $[0, 2\pi)$ [11–13], although there is some indications for non-zero δ_{CP} . Therefore we still don't know whether CP violation occurs in the lepton sector and how large it is if the CP symmetry is really violated. Measuring the leptonic CP violation is one of important goals of future long-baseline neutrino oscillation experiments [14].

Family symmetry and its spontaneous breaking have turned out to be able to naturally derive some mass independent textures, please see Ref. [15] for a review. In order to explain the observed lepton mixing angles and predict CP phases at the same time, it is natural to extend the family symmetry to include a generalised CP symmetry [16–22]. In this setup, the symmetries are spontaneously broken by the flavon vacuum expectation values (VEVs) which take specific discrete complex phases. As in the paradigm of family symmetry, the whole symmetry including both family and CP symmetries are generally broken into different subgroups in the neutrino and charged lepton sectors, and the mismatch between the two remnant subgroups gives rise to particular predictions for lepton mixing angles and CP phases.

Combining family symmetry with generalised CP symmetry is a promising framework to predict the values of CP violating phases. It has arisen some interesting discussions in the past years. Imposing generalised CP symmetry within the context of simple μ - τ interchange symmetry [23], A_4 [22], S_4 [19–21,24,25] and T' [26] family symmetries have been explored (other approaches to discrete symmetry and CP violation can be found in Refs. [27–29]). In such scenario, the mixing angles and CP phases are generally predicted to be strongly correlated with each other because of the constraint of the family and CP symmetries. The so-called trimaximal TM_2 neutrino mixing, whose second column of the mixing matrix is of the form $(1, 1, 1)^T/\sqrt{3}$, is frequently produced. In Refs. [19,22], the TM_2 mixing is a natural consequence of the preserved Z_2^S family symmetry in the neutrino sector. In the present work, we shall focus on the trimaximal TM_1 mixing whose first column of the mixing matrix takes the form $(2, -1, -1)^T/\sqrt{6}$, since the TM_1 mixing leads to better agreement of solar mixing angle θ_{12} with the measured value than the TM_2 pattern. We shall construct two typical models based on S_4 family symmetry and the corresponding generalised CP symmetry. Both models predict TM_1 mixing due to the remnant Z_2^{SU} symmetry in the neutrino sector and the Dirac CP is conserved or maximally broken. In the first model (Model 1), the flavon fields enter into the neutrino Dirac couplings instead of the Majorana mass terms of the right-handed neutrinos at leading order (LO), and the TM_1 mixing is generated at LO. After taking into account the measured solar and atmospheric neutrino mass squared differences and the reactor mixing angle θ_{13} , the absolute neutrino masses and the effective mass $|m_{\beta\beta}|$ for the neutrinoless double beta decay are fixed completely. For the second model (Model 2), the lepton mixing is of the tri-bimaximal form at LO with the remnant $Z_2^S \times Z_2^{SU}$ family symmetry in the neutrino sector, and the next-to-leading order (NLO) corrections further breaks $Z_2^S \times Z_2^{SU}$ into Z_2^{SU} such that TM_1 pattern is obtained. Since the non-zero θ_{13} arises from the NLO contributions, the relative smallness of the reactor angle with respect to the solar and atmospheric mixing angles are explained.

The paper is organized as follows. In Section 2, we briefly review the concept of generalised CP symmetry and the generalised CP transformation compatible with S_4 family symmetry. Moreover, the possible residual CP symmetries consistent with the remnant Z_2^{SU} family symmetry in the neutrino sector and the corresponding phenomenological predictions for the lepton mixing parameters are investigated. In Section 3, we present the first model, and we show that the desired vacuum configuration with their phase structure can be realized in a supersymmetric context. In

Section 4, we specify our second model, the LO structure of the model, the vacuum alignment and the NLO corrections induced by higher dimensional operators are discussed. We summarize and conclude in Section 5. The details of the group theory of S_4 are given in Appendix A, where the explicit representation matrices and the Clebsch–Gordan coefficients are listed.

2. General analysis of lepton mixing with residual Z_2^{SU} family symmetry and CP symmetry

2.1. Generalised CP transformations consistent with S_4

It is highly non-trivial to combine a family symmetry G_f with the generalised CP symmetry together [17–19]. Let us consider a generic multiplet of fields $\varphi(x)$ in the irreducible representation \mathbf{r} of G_f . Under the action of G_f , $\varphi(x)$ transforms as

$$\varphi \xrightarrow{G_f} \rho_{\mathbf{r}}(g)\varphi(x), \quad g \in G_f, \quad (2.1)$$

where $\rho_{\mathbf{r}}(g)$ is the representation matrix for the element g in the irreducible representation \mathbf{r} . The generalised CP transformation should leave the kinetic term $|\partial\varphi|^2$ invariant and it acts on $\varphi(x)$ as

$$\varphi(x) \xrightarrow{CP} X_{\mathbf{r}}\varphi^*(x'), \quad (2.2)$$

where $X_{\mathbf{r}}$ is a unitary matrix, $x' = (t, -\mathbf{x})$ and we have omit the action of CP on spinor indices for the case that φ is a spinor. Notice that we are considering the “minimal” theory in which the generalised CP transformation maps the field $\varphi \sim \mathbf{r}$ into its complex conjugate $\varphi^* \sim \mathbf{r}^*$. The generalised CP transformation $X_{\mathbf{r}}$ has to be consistently defined to be compatible with the family symmetry G_f . Hence the so-called consistency condition [17–19,30] must be satisfied

$$X_{\mathbf{r}}\rho_{\mathbf{r}}^*(g)X_{\mathbf{r}}^{-1} = \rho_{\mathbf{r}}(g'), \quad g, g' \in G_f. \quad (2.3)$$

Note that Eq. (2.3) should be fulfilled for all the irreducible representations of G_f . Moreover, Eq. (2.3) implies that the generalised CP transformation $X_{\mathbf{r}}$ maps the group element g into g' , and this mapping preserves the family symmetry group structure [18,19]. Therefore Eq. (2.3) defines a homomorphism of the family symmetry group G_f . It is now established that there is one to one correspondence between the generalised CP transformations and the automorphism group of the family symmetry group [30].

In the present work, we shall concentrate on the family symmetry $G_f = S_4$, which can be generated by three generators S , T and U . It is convenient to work in the T generator diagonal basis, the representation matrices for the three generators in different S_4 irreducible representations are summarized in Table 5. The corresponding Clebsch–Gordan coefficients are listed in Appendix A. The automorphism structure of S_4 is rather simple, since it doesn't have non-trivial outer automorphism. Therefore the automorphism of S_4 is exactly its inner automorphism, and the automorphism group of S_4 is isomorphic to S_4 itself. For the representative automorphism element $\text{conj}(U) : (S, T, U) \rightarrow (S, T^2, U)$, where $\text{conj}(h)$ denotes a group conjugation with an element h , i.e. $\text{conj}(h) : g \rightarrow hgh^{-1}$ with $h, g \in S_4$, the associated generalised CP transformation $X_{\mathbf{r}}^0$ is determined by the consistency equations

$$\begin{aligned} X_{\mathbf{r}}^0\rho_{\mathbf{r}}^*(S)(X_{\mathbf{r}}^0)^{-1} &= \rho_{\mathbf{r}}(S), & X_{\mathbf{r}}^0\rho_{\mathbf{r}}^*(T)(X_{\mathbf{r}}^0)^{-1} &= \rho_{\mathbf{r}}(T^2), \\ X_{\mathbf{r}}^0\rho_{\mathbf{r}}^*(U)(X_{\mathbf{r}}^0)^{-1} &= \rho_{\mathbf{r}}(U). \end{aligned} \quad (2.4)$$

From the explicit form of the representation matrices shown in Table 5, we see that for any irreducible representations \mathbf{r} of S_4 , the following relations are fulfilled

$$\rho_{\mathbf{r}}^*(S) = \rho_{\mathbf{r}}(S), \quad \rho_{\mathbf{r}}^*(U) = \rho_{\mathbf{r}}(U), \quad \rho_{\mathbf{r}}^*(T) = \rho_{\mathbf{r}}(T^2). \quad (2.5)$$

Therefore $X_{\mathbf{r}}^0$ is fixed to be equal to identity (up to an arbitrary overall phase), i.e.

$$X_{\mathbf{r}}^0 = 1. \quad (2.6)$$

Including the family symmetry transformation, the generalised CP transformation consistent with the S_4 family symmetry is given by

$$X_{\mathbf{r}} = \rho_{\mathbf{r}}(g)X_{\mathbf{r}}^0 = \rho_{\mathbf{r}}(g), \quad g \in S_4. \quad (2.7)$$

Hence the generalised CP transformation consistent with an S_4 family symmetry is of the same form as the family group transformation in the chosen basis. We confirm the results in Refs. [18, 19] that the generalised CP transformation group is the identity up to inner automorphism. Since we have found all generalised CP transformations consistent with the S_4 family symmetry, we turn to investigate their phenomenological implications on lepton masses and flavor mixings in the following.

2.2. Lepton mixing from $S_4 \rtimes H_{CP}$ breaking into $G_{CP}^l \cong Z_3^T \rtimes H_{CP}^l$ and $G_{CP}^v \cong Z_2^{SU} \times H_{CP}^v$

In this work we shall introduce the family symmetry S_4 together with the corresponding generalised CP symmetry H_{CP} at high energy scale, where H_{CP} is the collection of the generalised CP transformations $X_{\mathbf{r}}$. Hence the original symmetry of the theory is $S_4 \rtimes H_{CP}$. To obtain phenomenologically acceptable lepton masses and mixings, the original symmetry should be broken in both charged lepton and neutrino sectors. The mismatch between the symmetry breaking patterns in the neutrino and charged lepton sectors leads to particular predictions for lepton mixing angles and CP phases. In Ref. [19], the symmetry is broken down to $Z_2^S \times H_{CP}^v$ in the neutrino sector, and the residual family symmetry $Z_2^S = \{1, S\}$ enforces that the lepton mixing is the trimaximal TM_2 pattern [31–33], where the second column of the Pontecorvo–Maki–Nakagawa–Sakata (PMNS) matrix is proportional to $(1, 1, 1)^T$. In the present work, we shall investigate another case that $S_4 \rtimes H_{CP}$ is broken to $G_{CP}^l \cong Z_3^T \rtimes H_{CP}^l$ and $G_{CP}^v \cong Z_2^{SU} \times H_{CP}^v$ in the charged lepton and the neutrino sectors respectively, where $Z_3^T = \{1, T, T^2\}$ and $Z_2^{SU} = \{1, SU\}$. The remnant Z_2^{SU} symmetry would lead to the trimaximal TM_1 mixing pattern [21,34,35], where the first column of the PMNS matrix is proportional to $(2, -1, -1)^T$. General phenomenology analysis has shown that TM_1 mixing can lead to excellent agreement with the present data [33]. Furthermore, if the residual family symmetry in the neutrino sector chosen to be $Z_2^U = \{1, U\}$, the third column of the mixing matrix would be proportional to $(0, 1, -1)^T$. The reactor mixing angle would be predicted to be zero, and it is not consistent with both the experimental measurements [1–6] and the global data fitting [11–13]. Hence we don't consider this scenario.

In the charged lepton sector, the full symmetry $S_4 \rtimes H_{CP}$ is broken to $G_{CP}^l \cong Z_3^T \rtimes H_{CP}^l$. For G_{CP}^l to be a well-defined symmetry, the consistency condition of Eq. (2.3) should be satisfied for the residual family symmetry subgroup Z_3^T , i.e. the element $X_{\mathbf{r}l}$ of H_{CP}^l should fulfill

$$X_{\mathbf{r}l}\rho_{\mathbf{r}}^*(T)X_{\mathbf{r}l}^{-1} = \rho_{\mathbf{r}}(T'), \quad T' \in Z_3^T = \{1, T, T^2\}. \quad (2.8)$$

It is easy to check that the remnant CP symmetry H_{CP}^l can take the value

$$H_{CP}^l = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(T), \rho_{\mathbf{r}}(T^2), \rho_{\mathbf{r}}(U), \rho_{\mathbf{r}}(TU), \rho_{\mathbf{r}}(T^2U)\}. \quad (2.9)$$

Without loss of generality, we assume that the three generations of the left-handed lepton doublets are unified into the three-dimensional representation $\mathbf{3}$. The same results would be obtained if the lepton doublets were assigned to $\mathbf{3}'$ of S_4 , since the representation $\mathbf{3}'$ differs from $\mathbf{3}$ only in the overall sign of the generator U . The charged lepton mass matrix m_l is constrained by the remnant family symmetry Z_3^T and the remnant CP symmetry H_{CP}^l as

$$\rho_{\mathbf{3}}^\dagger(T) m_l m_l^\dagger \rho_{\mathbf{3}}(T) = m_l m_l^\dagger, \quad (2.10a)$$

$$X_{\mathbf{3}l}^\dagger m_l m_l^\dagger X_{\mathbf{3}l} = (m_l m_l^\dagger)^*, \quad (2.10b)$$

where the charged lepton mass matrix m_l is given in the convention in which the left-handed (right-handed) fields are on the left-hand (right-hand) side of m_l . Since the representation matrix $\rho_{\mathbf{3}}(T)$ is diagonal, the invariant condition Eq. (2.10a) under Z_3^T implies that $m_l m_l^\dagger$ is diagonal with

$$m_l m_l^\dagger = \text{diag}(m_e^2, m_\mu^2, m_\tau^2), \quad (2.11)$$

where m_e , m_μ and m_τ denote the electron, muon and tau masses, respectively. For the case of $X_{\mathbf{r}l} = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(T), \rho_{\mathbf{r}}(T^2)\}$, the conditions of Eq. (2.10b) is satisfied automatically, and therefore no additional constraints are required. For the remaining values $X_{\mathbf{r}l} = \{\rho_{\mathbf{r}}(U), \rho_{\mathbf{r}}(TU), \rho_{\mathbf{r}}(T^2U)\}$, the residual CP invariant condition of Eq. (2.10b) implies $m_\mu = m_\tau$. Hence this case is not viable phenomenologically. We note that in the models constructed in Sections 3 and 4, the Z_3^T remnant symmetry is broken by the flavon VEVs in order to facilitate the generation of the charged lepton mass hierarchies without fine tuning. However, we properly arrange the breaking such that the resulting charged lepton mass matrix remains diagonal. As a consequence, the hermitian product $m_l m_l^\dagger$ is invariant under the action of Z_3^T elements and the generalised CP transformations $X_{\mathbf{r}l} = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(T), \rho_{\mathbf{r}}(T^2)\}$, i.e. Eqs. (2.10a), (2.10b) are satisfied. Therefore the following general analysis is still meaningful and valid, and in particular it guides our model building.

Now we turn to the neutrino sector. In order to reproduce the TM_1 mixing pattern, the symmetry $S_4 \times H_{CP}$ is spontaneously broken to $G_{CP}^v = Z_2^{SU} \times H_{CP}^v$. The residual CP symmetry H_{CP}^v should be consistent with the residual family symmetry Z_2^{SU} , and therefore its element $X_{\mathbf{r}v}$ has to fulfill the consistency equation

$$X_{\mathbf{r}v} \rho_{\mathbf{r}}^*(SU) X_{\mathbf{r}v}^{-1} = \rho_{\mathbf{r}}(SU). \quad (2.12)$$

One can easily check that there are only 4 possible choices for $X_{\mathbf{r}v}$, i.e.

$$H_{CP}^v = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(S), \rho_{\mathbf{r}}(U), \rho_{\mathbf{r}}(SU)\}. \quad (2.13)$$

The light neutrino mass matrix m_ν is constrained by the residual family symmetry Z_2^{SU} and residual CP symmetry H_{CP}^v as [19]

$$\rho_{\mathbf{3}}^T(SU) m_\nu \rho_{\mathbf{3}}(SU) = m_\nu, \quad (2.14a)$$

$$X_{\mathbf{3}v}^T m_\nu X_{\mathbf{3}v} = m_\nu^*. \quad (2.14b)$$

The most general neutrino mass matrix which satisfies Eq. (2.14a) is of the form

$$m_\nu = \alpha \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + \beta \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + \gamma \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + \delta \begin{pmatrix} 0 & 1 & -1 \\ 1 & 2 & 0 \\ -1 & 0 & -2 \end{pmatrix}, \tag{2.15}$$

where the four parameters α , β , γ and δ are generally complex, and the remnant CP invariant condition of Eq. (2.14b) would further constrain these parameters to be real or purely imaginary.

In order to diagonalize light neutrino mass matrix m_ν in Eq. (2.15), it is useful to first perform a tri-bimaximal transformation U_{TB}

$$m'_\nu = U_{TB}^T m_\nu U_{TB} = \begin{pmatrix} 3\alpha + \beta - \gamma & 0 & 0 \\ 0 & \beta + 2\gamma & -\sqrt{6}\delta \\ 0 & -\sqrt{6}\delta & 3\alpha - \beta + \gamma \end{pmatrix}, \tag{2.16}$$

with

$$U_{TB} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \tag{2.17}$$

Then we investigate the implication of the remnant CP invariant condition of Eq. (2.14b). Two distinct phenomenological predictions arise for $X_{\mathbf{r}\nu} = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(SU)\}$ and $X_{\mathbf{r}\nu} = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(SU)\}$. We shall discuss the two cases in detail in the following.

(1) $X_{\mathbf{r}\nu} = \rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(SU)$

In this case, we can straightforwardly find that all the four parameters α , β , γ and δ are constrained to be real. As a result, m'_ν becomes a real symmetry matrix and can be diagonalized by a rotation matrix $R(\theta)$ in the (2, 3) sector with

$$R(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta & \sin\theta \\ 0 & -\sin\theta & \cos\theta \end{pmatrix}, \tag{2.18}$$

where

$$\tan 2\theta = \frac{-2\sqrt{6}\delta}{3\alpha - 2\beta - \gamma}. \tag{2.19}$$

Hence we have

$$U_\nu'^T m'_\nu U'_\nu = \text{diag}(m_1, m_2, m_3), \quad U'_\nu = R(\theta)P, \tag{2.20}$$

where P is a unitary diagonal matrix with entries ± 1 or $\pm i$, which encode the CP parity of the neutrino state. Furthermore, the light neutrino masses $m_{1,2,3}$ are determined to be

$$\begin{aligned} m_1 &= |3\alpha + \beta - \gamma|, \\ m_2 &= \frac{1}{2} |3(\alpha + \gamma) - \text{sign}((3\alpha - 2\beta - \gamma) \cos 2\theta) \sqrt{24\delta^2 + (3\alpha - 2\beta - \gamma)^2}|, \\ m_3 &= \frac{1}{2} |3(\alpha + \gamma) + \text{sign}((3\alpha - 2\beta - \gamma) \cos 2\theta) \sqrt{24\delta^2 + (3\alpha - 2\beta - \gamma)^2}|. \end{aligned} \tag{2.21}$$

We see that the three neutrino masses depend on four real parameters, and therefore any neutrino mass spectrum can be realized in this scenario. Since the charged lepton mass matrix

is diagonal, the lepton mixing matrix U_{PMNS} is fixed by neutrino sector completely, and we have

$$U_{PMNS} = U_{TB}U'_\nu = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{\cos\theta}{\sqrt{3}} & \frac{\sin\theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos\theta}{\sqrt{3}} + \frac{\sin\theta}{\sqrt{2}} & -\frac{\cos\theta}{\sqrt{2}} + \frac{\sin\theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos\theta}{\sqrt{3}} - \frac{\sin\theta}{\sqrt{2}} & \frac{\cos\theta}{\sqrt{2}} + \frac{\sin\theta}{\sqrt{3}} \end{pmatrix} P. \quad (2.22)$$

The three lepton mixing angles θ_{13} , θ_{12} and θ_{23} are predicted to be

$$\begin{aligned} \sin^2\theta_{13} &= \frac{1}{3}\sin^2\theta, & \sin^2\theta_{12} &= \frac{\cos^2\theta}{2 + \cos^2\theta} = \frac{1}{3} - \frac{2}{3}\tan^2\theta_{13}, \\ \sin^2\theta_{23} &= \frac{1}{2} - \frac{\sqrt{6}\sin\theta\cos\theta}{3 - \sin^2\theta} = \frac{1}{2} \pm \tan\theta_{13}\sqrt{2(1 - 2\tan^2\theta_{13})}. \end{aligned} \quad (2.23)$$

For the best fitting value of the reactor angle $\theta_{13} = 8.71^\circ$ [13], the remaining two mixing angles are determined to be $\theta_{12} \simeq 34.31^\circ$ and $\theta_{23} \simeq 32.49^\circ$ or $\theta_{23} \simeq 57.51^\circ$, which are compatible with the preferred values from global fits. Adopting the PDG parameterization [36], the Dirac CP violating phase δ_{CP} and two Majorana CP violating phases α_{21} and α_{31} take the values

$$\sin\delta_{CP} = \sin\alpha_{21} = \sin\alpha_{31} = 0, \quad (2.24)$$

which implies

$$\delta_{CP}, \alpha_{21}, \alpha_{31} = 0, \pi. \quad (2.25)$$

Hence there is no CP violation in this case.

(2) $X_{\nu\nu} = \rho_r(S), \rho_{\nu\nu}(U)$

Solving the residual CP invariant equation of Eq. (2.14b), we find the three parameters α , β and γ are real, and δ is purely imaginary. The unitary transformation U'_ν diagonalizing the neutrino mass matrix m'_ν is of the form

$$U'_\nu = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta & \sin\theta \\ 0 & -i\sin\theta & i\cos\theta \end{pmatrix} P, \quad (2.26)$$

with

$$\tan 2\theta = \frac{2i\sqrt{6}\delta}{3(\alpha + \gamma)}. \quad (2.27)$$

The resulting PMNS matrix is

$$U_{PMNS} = U_{TB}U'_\nu = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{\cos\theta}{\sqrt{3}} & \frac{\sin\theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos\theta}{\sqrt{3}} + \frac{i\sin\theta}{\sqrt{2}} & -\frac{i\cos\theta}{\sqrt{2}} + \frac{\sin\theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos\theta}{\sqrt{3}} - \frac{i\sin\theta}{\sqrt{2}} & \frac{i\cos\theta}{\sqrt{2}} + \frac{\sin\theta}{\sqrt{3}} \end{pmatrix} P. \quad (2.28)$$

The lepton mixing angles and CP phases are determined to be

Table 1

The particle contents and their transformation property under the family symmetry $S_4 \times Z_7$ and $U(1)_R$, where $\omega_7 = e^{\frac{2\pi i}{7}}$.

Field	l	ν^c	e^c	μ^c	τ^c	$h_{u,d}$	φ_T	ϕ	φ_S	η	ξ	φ_T^0	ζ^0	φ_S^0	η^0
S_4	$\mathbf{3}'$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{2}$	$\mathbf{3}$	$\mathbf{2}$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{3}$	$\mathbf{2}$
Z_7	ω_7	1	ω_7^3	ω_7^4	ω_7^5	1	ω_7	ω_7	ω_7^6	ω_7^6	ω_7^6	ω_7^5	ω_7^5	ω_7^2	ω_7^2
$U(1)_R$	1	1	1	1	1	0	0	0	0	0	0	2	2	2	2

$$\begin{aligned}
 |\sin \delta_{CP}| &= 1, & \sin \alpha_{21} &= \sin \alpha_{31} = 0, \\
 \sin^2 \theta_{13} &= \frac{1}{3} \sin^2 \theta, & \sin^2 \theta_{12} &= \frac{\cos^2 \theta}{2 + \cos^2 \theta} = \frac{1}{3} - \frac{2}{3} \tan^2 \theta_{13}, \\
 \sin^2 \theta_{23} &= \frac{1}{2}.
 \end{aligned} \tag{2.29}$$

The predictions for both the solar and reactor mixing angles are the same as the ones in case (I), and the atmospheric mixing angle is maximal. Moreover, we have maximal Dirac CP violation $\delta_{CP} = \pm \frac{\pi}{2}$, and Majorana phases are trivial with $\alpha_{21}, \alpha_{31} = 0, \pi$. Finally, the light neutrino masses are given by

$$\begin{aligned}
 m_1 &= |3\alpha + \beta - \gamma|, \\
 m_2 &= \frac{1}{2} \left| -3\alpha + 2\beta + \gamma + \text{sign}((\alpha + \gamma) \cos 2\theta) \sqrt{9(\alpha + \gamma)^2 - 24\delta^2} \right|, \\
 m_3 &= \frac{1}{2} \left| -3\alpha + 2\beta + \gamma - \text{sign}((\alpha + \gamma) \cos 2\theta) \sqrt{9(\alpha + \gamma)^2 - 24\delta^2} \right|.
 \end{aligned} \tag{2.30}$$

Notice that the above results are exactly the same as those of Ref. [17], although we use a different basis in which the generator T is diagonal. The chosen basis in the present paper is particularly suitable to build TM_1 model, since the charged lepton mass matrix is diagonal in this basis and the lepton mixing completely comes from the neutrino sector. Now that we have finished the general analysis, we proceed to construct models to realize these model independent results. Two typical models would be proposed in the following sections. In the first model, the lepton mixing is the TM_1 pattern at LO. In the second model, tri-bimaximal mixing is produced at LO, and it is broken to TM_1 mixing by the NLO corrections. As a consequence, the relative smallness of θ_{13} with respect to θ_{12} and θ_{23} is explained.

3. Model 1

In this section, we shall present the first TM_1 model (Model 1) based on $S_4 \times H_{CP}$ with the extra symmetry $Z_7 \times U(1)_R$. We shall formulate the model in the framework of type I see-saw mechanism and supersymmetry (SUSY). Both the three generations of left-handed lepton doublets l and the right-handed neutrinos ν^c are assigned to transform as S_4 triplet $\mathbf{3}'$, while the RH charged leptons e^c , μ^c and τ^c are all invariant under S_4 . The involved fields and their transformation rules under the family symmetry $S_4 \times Z_7 \times U(1)_R$ are summarized in Table 1. Notice that the auxiliary Z_7 symmetry separates the flavon fields entering the charged lepton sector at LO from those entering the neutrino sector, and it is also helpful to achieve the charged lepton mass hierarchies and suppress the NLO corrections. Compared with most flavor models in which the flavon fields generally couple to the right-handed neutrinos at LO, the flavons are involved in

the neutrino Dirac mass term instead of the Majorana mass term of right-handed neutrino in the present model.

3.1. Vacuum alignment

We adopt the now-standard F -term alignment mechanism to arrange the vacuum [37]. A continuous $U(1)_R$ symmetry related to R -parity is generally introduced under which the matter fields carry a $+1$ R -charge while the electroweak Higgs and flavon fields are uncharged. In addition, one needs the so-called driving fields carrying two unit of R -charge, and hence each term in the superpotential can contain at most one driving field. In the SUSY limit, the minimization of the flavon potential can be achieved simply by ensuring that the F -terms of the driving fields vanish at the minimum. The required driving fields and their transformation rules are listed in Table 1. The LO driving superpotential w_d invariant under the family symmetry $S_4 \times Z_7$ can be written as

$$w_d = w_d^l + w_d^v, \quad (3.1)$$

where w_d^l is the flavon superpotential which contains the flavons only entering into the charged lepton at LO, i.e.

$$w_d^l = f_1(\varphi_T^0(\varphi_T\varphi_T)\mathbf{3})_{\mathbf{1}} + f_2(\varphi_T^0(\phi\varphi_T)\mathbf{3})_{\mathbf{1}} + f_3\zeta^0(\varphi_T\varphi_T)_{\mathbf{1}} + f_4\zeta^0(\phi\phi)_{\mathbf{1}}. \quad (3.2)$$

w_d^v is the superpotential associated with the flavons in the neutrino sector

$$w_d^v = g_1(\varphi_S^0(\varphi_S\varphi_S)\mathbf{3})_{\mathbf{1}} + g_2(\varphi_S^0(\eta\varphi_S)\mathbf{3})_{\mathbf{1}} + g_3\xi(\varphi_S^0\varphi_S)_{\mathbf{1}} + g_4(\eta^0(\varphi_S\varphi_S)\mathbf{2})_{\mathbf{1}} \\ + g_5(\eta^0(\eta\eta)\mathbf{2})_{\mathbf{1}} + g_6\xi(\eta^0\eta)_{\mathbf{1}}, \quad (3.3)$$

where $(\dots)_{\mathbf{r}}$ denotes the contraction into S_4 irreducible representation \mathbf{r} according to the Clebsch–Gordan coefficients presented in Appendix A. We note that the first term vanishes automatically due to the anti-symmetric property of the contraction $(\varphi_S\varphi_S)\mathbf{3}$. Since we require the theory to be invariant under the generalised CP transformation, then all the couplings f_i and g_i in Eqs. (3.2), (3.3) are constrained to be real. We start from the charged lepton sector, and the F -term conditions obtained from the driving fields φ_T^0 and ζ^0 read

$$\frac{\partial w_d}{\partial \varphi_{T_1}^0} = 2f_1(\varphi_{T_1}^2 - \varphi_{T_2}\varphi_{T_3}) + f_2(\phi_1\varphi_{T_2} + \phi_2\varphi_{T_3}) = 0, \\ \frac{\partial w_d}{\partial \varphi_{T_2}^0} = 2f_1(\varphi_{T_2}^2 - \varphi_{T_1}\varphi_{T_3}) + f_2(\phi_1\varphi_{T_1} + \phi_2\varphi_{T_2}) = 0, \\ \frac{\partial w_d}{\partial \varphi_{T_3}^0} = 2f_1(\varphi_{T_3}^2 - \varphi_{T_1}\varphi_{T_2}) + f_2(\phi_1\varphi_{T_3} + \phi_2\varphi_{T_1}) = 0, \\ \frac{\partial w_d}{\partial \zeta^0} = f_3(\varphi_{T_1}^2 + 2\varphi_{T_2}\varphi_{T_3}) + 2f_4\phi_1\phi_2 = 0. \quad (3.4)$$

We find two possible solutions for the vacuum (up to S_4 transformations). The first one is given by

$$\langle \varphi_T \rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} v_T, \quad \langle \phi \rangle = \begin{pmatrix} 1 \\ -1 \end{pmatrix} v_\phi, \quad \text{with } v_T^2 = \frac{2f_4}{3f_3} v_\phi^2. \quad (3.5)$$

The second solution is

$$\langle \varphi_T \rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} v_T, \quad \langle \phi \rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} v_\phi, \quad \text{with } v_T = -\frac{f_2}{2f_1} v_\phi. \quad (3.6)$$

We shall choose the second solution in this work. We note that the phase of v_ϕ can be absorbed into lepton fields, and therefore we can take both v_ϕ and v_T to be real, since f_1 and f_2 are real. Furthermore, v_ϕ and v_T are expected to be of the same order of magnitude without fine tuning among the parameters f_1 and f_2 . The vacuum expectation values of φ_S , η and ξ , which give rise to TM_1 mixing in the neutrino sector, are determined by the F -terms of the associated driving fields as follows:

$$\begin{aligned} \frac{\partial w_d^v}{\partial \varphi_{S_1}^0} &= g_2(\eta_1 \varphi_{S_2} + \eta_2 \varphi_{S_3}) + g_3 \xi \varphi_{S_1} = 0, \\ \frac{\partial w_d^v}{\partial \varphi_{S_2}^0} &= g_2(\eta_1 \varphi_{S_1} + \eta_2 \varphi_{S_2}) + g_3 \xi \varphi_{S_3} = 0, \\ \frac{\partial w_d^v}{\partial \varphi_{S_3}^0} &= g_2(\eta_1 \varphi_{S_3} + \eta_2 \varphi_{S_1}) + g_3 \xi \varphi_{S_2} = 0, \\ \frac{\partial w_d^v}{\partial \eta_1^0} &= g_4(\varphi_{S_3}^2 + 2\varphi_{S_1} \varphi_{S_2}) + g_5 \eta_1^2 + g_6 \xi \eta_2 = 0, \\ \frac{\partial w_d^v}{\partial \eta_2^0} &= g_4(\varphi_{S_2}^2 + 2\varphi_{S_1} \varphi_{S_3}) + g_5 \eta_2^2 + g_6 \xi \eta_1 = 0. \end{aligned} \quad (3.7)$$

There are two independent solutions to this set of equations up to S_4 family symmetry transformations. The first solution is

$$\langle \varphi_S \rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} v_S, \quad \langle \eta \rangle = \begin{pmatrix} 1 \\ 1 \end{pmatrix} v_\eta, \quad \langle \xi \rangle = v_\xi. \quad (3.8)$$

The VEVs v_S , v_η and v_ξ are related with each other via

$$v_S^2 = \frac{g_3(2g_2g_6 - g_3g_5)}{12g_4g_2^2} v_\xi^2, \quad v_\eta = -\frac{g_3}{2g_2} v_\xi, \quad (3.9)$$

where v_ξ is undetermined and generally complex. With the representation matrix given in [Appendix A](#), we can straightforwardly check that the S_4 family symmetry is broken down to $Z_2^S \times Z_2^{SU}$ subgroup by the vacuum alignment of Eq. (3.8). The second solution reads as

$$\langle \varphi_S \rangle = \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix} v_S, \quad \langle \eta \rangle = \begin{pmatrix} 1 \\ 1 \end{pmatrix} v_\eta, \quad \langle \xi \rangle = v_\xi, \quad (3.10)$$

with

$$v_S^2 = \frac{g_3(g_3g_5 + g_2g_6)}{3g_4g_2^2} v_\xi^2, \quad v_\eta = \frac{g_3}{g_2} v_\xi. \quad (3.11)$$

We find the S_4 family symmetry is spontaneously broken down to Z_2^{SU} subgroup in this case. In order to reproduce the TM_1 pattern, we choose the second solution in the following. Since

all the couplings g_i are real due to the generalised CP invariance, Eq. (3.11) implies that the VEVs v_η and v_ξ have the same phase up to π , and the phase difference between v_S and v_ξ is $0, \pi$ or $\pm\frac{\pi}{2}$ depending on the sign of $g_3g_4(g_3g_5 + g_2g_6)$. In addition, it is natural to expect that the three VEVs v_ξ, v_η and v_S are of the same order of magnitudes. As shall be shown below, the phase of v_ξ turns out to be an overall phase of the light neutrino mass matrix, and hence it can be absorbed into the neutrino fields. That is to say we can take v_ξ to be real without loss of generality. As a consequence, the VEV v_η would be real as well and the VEV v_S is real for the product $g_3g_4(g_3g_5 + g_2g_6) > 0$ or purely imaginary for $g_3g_4(g_3g_5 + g_2g_6) < 0$.

Regarding the order of magnitude of the different VEVs, as we shall find in the following, the charged lepton mass hierarchies can be naturally reproduced if v_ϕ/Λ and v_T/Λ are of order λ^2 , i.e.

$$\frac{v_\phi}{\Lambda} \sim \frac{v_T}{\Lambda} \sim \lambda^2, \quad (3.12)$$

where $\lambda \simeq 0.23$ is the Cabibbo angle. In order to guarantee the stability of the successful LO results under the inclusion of higher dimensional terms, we choose all the VEVs in the model are of the same order $\lambda^2\Lambda$, i.e.

$$\frac{v_S}{\Lambda} \sim \frac{v_\eta}{\Lambda} \sim \frac{v_\xi}{\Lambda} \sim \lambda^2. \quad (3.13)$$

This assumption is frequently used in the family symmetry model building.

3.2. The lepton masses and mixing

The most general superpotential for the charged lepton masses, which is invariant under the family symmetry, is of the form

$$\begin{aligned} w_l = & \frac{y_\tau}{\Lambda} (l\varphi_T)_1 \tau^c h_d + \frac{y_{\mu 1}}{\Lambda^2} (l(\varphi_T\varphi_T)_{3'})_1 \mu^c h_d + \frac{y_{\mu 2}}{\Lambda^2} (l(\phi\varphi_T)_{3'})_1 \mu^c h_d \\ & + \frac{y_{e1}}{\Lambda^3} (l\varphi_T)_1 (\varphi_T\varphi_T)_1 e^c h_d + \frac{y_{e2}}{\Lambda^3} ((l\varphi_T)_2 (\varphi_T\varphi_T)_2)_1 e^c h_d \\ & + \frac{y_{e3}}{\Lambda^3} ((l\varphi_T)_{3'} (\varphi_T\varphi_T)_{3'})_1 e^c h_d \\ & + \frac{y_{e4}}{\Lambda^3} ((l\varphi_T)_3 (\varphi_T\varphi_T)_3)_1 e^c h_d + \frac{y_{e5}}{\Lambda^3} ((l\phi)_{3'} (\varphi_T\varphi_T)_{3'})_1 e^c h_d \\ & + \frac{y_{e6}}{\Lambda^3} ((l\phi)_3 (\varphi_T\varphi_T)_3)_1 e^c h_d + \frac{y_{e7}}{\Lambda^3} ((l\varphi_T)_2 (\phi\phi)_2)_1 e^c h_d \\ & + \frac{y_{e8}}{\Lambda^3} (l\varphi_T)_1 (\phi\phi)_1 e^c h_d + \dots, \end{aligned} \quad (3.14)$$

where dots represent the higher dimensional operators which will be commented later. Generalised CP symmetry enforces the Yukawa couplings to be real. Due to the constraint of the Z_7 symmetry, the electron, muon and tau mass terms are suppressed by $1/\Lambda, 1/\Lambda^2$ and $1/\Lambda^3$ respectively. With the vacuum alignment of Eq. (3.6), we find the resulting charged lepton mass matrix is diagonal with

$$\begin{aligned} m_e = & \left(y_{e2} - 2y_{e3} + 2y_{e5} \frac{v_\phi}{v_T} + y_{e7} \frac{v_\phi^2}{v_T^2} \right) \frac{v_T^3}{\Lambda^3} v_d, \\ m_\mu = & \left(2y_{\mu 1} + y_{\mu 2} \frac{v_\phi}{v_T} \right) \frac{v_T^2}{\Lambda^2} v_d, \quad m_\tau = y_\tau \frac{v_T}{\Lambda} v_d, \end{aligned} \quad (3.15)$$

in which $v_d = \langle h_d \rangle$ is the VEV of the electroweak Higgs field h_d . We see that the observed mass hierarchies among the charged leptons can be generated for $v_\phi/\Lambda \sim v_T/\Lambda \sim \lambda^2$. For the vacuum of φ_T and ϕ in Eq. (3.6), we can check that the S_4 family symmetry is broken completely in the charged lepton sector, since $T\langle\varphi_T\rangle = \omega^2\langle\varphi_T\rangle$ and $T\langle\phi\rangle = \omega^2\langle\phi\rangle$. However, the lepton flavor mixing is associated with the hermitian combination $m_l m_l^\dagger$, which is obviously invariant under the action of T , i.e., $T^\dagger m_l m_l^\dagger T = m_l m_l^\dagger$. Consequently there is still a remnant Z_3^T symmetry in the charged lepton sector if we concentrate on lepton flavor mixing. Furthermore, we can check that only three of the 24 generalised CP symmetries are preserved by $m_l m_l^\dagger$ and $H_{CP}^l = \{\rho_r(1), \rho_r(T), \rho_r(T^2)\}$.

Neutrino masses are generated by type I see-saw mechanism. The LO superpotential is given by

$$w_\nu = \frac{y_1}{\Lambda} (lv^c)_1 \xi h_u + \frac{y_2}{\Lambda} ((lv^c)_2 \eta)_1 h_u + \frac{y_3}{\Lambda} ((lv^c)_3 \varphi_S)_1 h_u + M (v^c v^c)_1, \tag{3.16}$$

where the first three terms contribute to the neutrino Dirac mass whereas the last one is the Majorana mass terms for the right-handed neutrinos. All the couplings are again real because of the imposed generalised CP symmetry. Given the vacuum configuration of Eq. (3.10), we can read out the Dirac and Majorana mass matrices as follows

$$m_D = y_1 v_u \frac{v_\xi}{\Lambda} \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + x \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + y \begin{pmatrix} 0 & -1 & 1 \\ 1 & 0 & 2 \\ -1 & -2 & 0 \end{pmatrix} \right],$$

$$m_M = M \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \tag{3.17}$$

where $v_u = \langle h_u \rangle$ and the parameters x, y are

$$x = \frac{y_2 v_\eta}{y_1 v_\xi}, \quad y = \frac{y_3 v_S}{y_1 v_\xi}. \tag{3.18}$$

After extracting the common phase of the VEVs v_S, v_η and v_ξ , the parameter x is real, while y is real or purely imaginary. The light neutrino mass matrix is given by the see-saw formula

$$m_\nu = -m_D m_M^{-1} m_D^T$$

$$= \alpha \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + \beta \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$+ \gamma \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + \delta \begin{pmatrix} 0 & 1 & -1 \\ 1 & 2 & 0 \\ -1 & 0 & -2 \end{pmatrix}. \tag{3.19}$$

It is the most general neutrino mass matrix consistent with the residual Z_2^{SU} flavor symmetry, as is shown in Eq. (2.15). The four parameters α, β, γ and δ are given by

$$\alpha = -y^2 m_0, \quad \beta = (4y^2 - 2x^2 - 1)m_0,$$

$$\gamma = (y^2 - x^2 - 2x)m_0, \quad \delta = -3xym_0, \tag{3.20}$$

Table 2

The predictions for the leptonic CP phases, light neutrino masses m_i ($i = 1, 2, 3$) and the effective mass $|m_{\beta\beta}|$ of the neutrinoless doublet-beta decay, where the unit of mass is meV.

(x, y)	δ_{CP}	$\theta_{23}/^\circ$	$\theta_{12}/^\circ$	α_{21}	α_{31}	m_1	m_2	m_3	$ m_{\beta\beta} $	Mass order
$(-1.898, -0.316)$	π	32.496	34.309	0	π	128.020	128.311	137.136	122.038	NO
$(-1.898, 0.316)$	0	57.504								
$(0.139, -0.612)$	π	32.496		π	0	24.233	25.724	54.747	9.423	NO
$(0.139, 0.612)$	0	57.504								
$(0.101, 0.340)$	π	32.496		0	π	49.669	50.414	11.159	48.507	IO
$(0.101, -0.340)$	0	57.504								
$(-0.120, 0.535)$	π	32.496		π	0	54.866	55.541	25.977	19.931	IO
$(-0.120, -0.535)$	0	57.504								
$(-0.050, 0.233i)$	$\pi/2$	45		0	0	57.284	57.930	75.488	57.901	NO
$(-0.050, -0.233i)$	$-\pi/2$									

where $m_0 = \frac{y_1^2 v_u^2 v_\xi^2}{M \Lambda^2}$ is the overall scale of the light neutrino masses. We see that α , β and γ are real parameters, δ is real or imaginary for v_S being real or imaginary, respectively. Furthermore, the effective mass parameter $|m_{\beta\beta}|$ for the neutrinoless double-beta decay is given by

$$|m_{\beta\beta}| = m_0 |2\alpha + \beta|. \quad (3.21)$$

As shown in Eq. (3.11), if the combination $g_3 g_4 (g_3 g_5 + g_2 g_6) > 0$, which leads to real v_S and δ parameters, the vacuum alignments of the flavons φ_S , η and ξ in Eq. (3.10) are invariant under the action of both $\rho_r(1)$ and $\rho_r(SU)$ elements of H_{CP} . Therefore the generalised CP symmetry is broken to $H_{CP}^v = \{\rho_r(1), \rho_r(SU)\}$ in the neutrino sector. This case is identical to case (I) of the general analysis inspired by symmetry arguments. The corresponding light neutrino mass matrix of Eq. (3.19) is real, the lepton mixing is exactly the TM_1 pattern with conserved CP, and the predictions for light neutrino masses and mixing angles are given in Eq. (2.21) and Eq. (2.23). Notice that the light neutrino mass matrix of Eq. (3.19) depends on three real parameters x , y and m_0 , their values can be fixed by the measured values of the mass squared differences Δm_{sol}^2 and Δm_{atm}^2 and the reactor neutrino mixing angle θ_{13} . As a result, both the absolute scale of the neutrino masses and the lepton mixing angles are fixed. For the best fitting values of $\Delta m_{sol}^2 = 7.45 \times 10^{-5} \text{ eV}^2$, $\Delta m_{atm}^2 = 2.417(2.410) \times 10^{-3} \text{ eV}^2$ and $\sin^2 \theta_{13} = 0.0229$ from Ref. [13], we find there are 8 solutions to the values of x and y in the case that both x and y are real. The corresponding predictions for the light neutrino masses and the lepton mixing parameters are summarized in Table 2. It is obvious that the former 4 solutions correspond to a normal ordering (NO) neutrino mass spectrum, and the latter 4 correspond to inverted ordering (IO) spectrum. Moreover, we see that the predicted values for the atmospheric mixing angle θ_{23} (32.496° and 57.504°) are slightly beyond the 3σ range of the current global data fitting [11–13]. We note that the NLO corrections and the renormalization group evolution effects could bring the model to agree with the experimental data. However, in these scenarios a value of θ_{23} very close to the maximal mixing value of 45° would be unnatural. The next generation neutrino oscillation experiments, in particular those exploiting a high intensity neutrino beam, will reduce the experimental error on θ_{23} to few degrees. If no significant deviations from maximal atmospheric mixing will be detected, these 8 solutions will be ruled out.

Another possibility of $g_3 g_4 (g_3 g_5 + g_2 g_6) < 0$ gives rise to an imaginary v_S such that the parameter δ in the neutrino mass matrix of Eq. (3.19) is purely imaginary as well. The remnant CP symmetry in the neutrino sector is $H_{CP}^v = \{\rho_r(S), \rho_r(U)\}$. This corresponds to the case (II) discussed in the general analysis of Section 2.2. The predictions for the mixing parameters and

Table 3

The transformation properties of the fields under the family symmetry $S_4 \times Z_4 \times Z_5$ and $U(1)_R$, where $\omega_5 = e^{\frac{2\pi i}{5}}$.

Field	l	ν^c	e^c	μ^c	τ^c	$h_{u,d}$	φ_T	ϕ	φ_S	η	χ	ξ	φ_T^0	ζ^0	φ_S^0	ξ^0	η^0	ρ^0	σ^0	
S_4	$\mathbf{3}'$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{2}$	$\mathbf{3}'$	$\mathbf{2}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{3}$	$\mathbf{1}$	$\mathbf{2}$	$\mathbf{1}$	$\mathbf{1}$	
Z_4	1	1	i	-1	$-i$	1	i	i	1	1	1	1	-1	-1	1	1	1	1	1	1
Z_5	ω_5^3	ω_5^2	ω_5^2	ω_5^2	ω_5^2	1	1	1	ω_5	ω_5	ω_5^3	ω_5^3	1	1	ω_5^3	ω_5^3	ω_5^4	ω_5^4	ω_5	ω_5
$U(1)_R$	1	1	1	1	1	0	0	0	0	0	0	0	2	2	2	2	2	2	2	2

the light neutrino masses are given in Eq. (2.29) and Eq. (2.30). The lepton mixing is of the TM_1 form, and maximal Dirac CP violation $|\delta_{CP}| = \pi/2$ and maximal atmospheric mixing $\theta_{23} = 45^\circ$ are produced in this case. Analogously, the light neutrino sector is also controlled by three real parameters, and hence the model is quite predictive, as shown in the last two lines of Table 2. The neutrino mass spectrum can only be normal ordering in this case.

Generally the LO results are modified by the subleading terms invariant under the imposed symmetry. Because of the auxiliary Z_7 symmetry in the present model, all the subleading corrections can be obtained by inserting the combination $\Phi_l \Phi_\nu$ into the LO terms of w_d , w_l and w_ν in Eqs. (3.1), (3.14), (3.16),¹ where $\Phi_l = \{\varphi_T, \phi\}$ and $\Phi_\nu = \{\varphi_S, \eta, \xi\}$ denote the flavons in the charged lepton and neutrino sectors respectively. As a result, the corresponding corrections to the lepton masses and mixing angles are suppressed by $\langle \Phi_l \rangle \langle \Phi_\nu \rangle / \Lambda^2 \sim \lambda^4$ with respect to the LO contributions and therefore can be negligible.

4. Model 2

In this section, we shall try to improve the previous model by generating the reactor mixing angle at the next-to-leading order (NLO) such that the correct order of magnitude of θ_{13} is produced. In this model, the LO lepton mixing is the well-known tri-bimaximal mixing pattern which is broken to TM_1 mixing by NLO contributions. Analogous to the previous model, the present model is based on the symmetry $S_4 \times H_{CP}$ with the extra symmetry $Z_4 \times Z_5 \times U(1)_R$ in order to eliminate unwanted operators. The matter fields, flavon fields, driving fields and their transformation rules under the family symmetry are summarized in Table 3. As previous model of Section 3, the remnant symmetry of the hermitian combination $m_l m_l^\dagger$ is $Z_3^T \times H_{CP}^l$ with $H_{CP}^l = \{\rho_r(1), \rho_r(T), \rho_r(T^2)\}$, and the original symmetry $S_4 \times H_{CP}$ is broken down to $G_{CP}^{nu} = Z_2^{SU} \times H_{CP}^\nu$. As a consequence, the model-independent analysis results of Section 2.2 are realized within one model, and the Dirac CP phase δ_{CP} is predicted to be trivial or maximal. In the following, we firstly discuss the vacuum alignment of the model, then specify the structure of the model at LO and NLO.

4.1. Vacuum alignment

The most general driving superpotential w_d^l associated with the charged lepton sector, which is invariant under the family symmetry $S_4 \times Z_4 \times Z_5$, can be written as

$$w_d^l = f_1(\varphi_T^0(\varphi_T \varphi_T)_3)_1 + f_2(\varphi_T^0(\phi \varphi_T)_3)_1 + f_3 \zeta^0(\varphi_T \varphi_T)_1 + f_4 \zeta^0(\phi \phi)_1. \tag{4.1}$$

¹ All possible S_4 contractions should be considered here, and only the correction to the electron mass terms is an exception with the form $(l \Phi_\nu^4) e^c h_d / \Lambda^4$.

It is exactly the same as Eq. (3.2). Hence the vacuum of the flavon fields φ_T and ϕ is of the same form as shown in Eq. (3.6), i.e.

$$\langle \varphi_T \rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} v_T, \quad \langle \phi \rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} v_\phi, \quad \text{with } v_T = -\frac{f_2}{2f_1} v_\phi. \quad (4.2)$$

We see that v_ϕ and v_T carry the same phase up to π . Since the phase of v_ϕ can be absorbed by leptons, we can take v_ϕ and v_T to be real without loss of generality. From the following predictions for charged lepton masses in Eq. (4.10), we note that the mass hierarchies between the charged leptons can be produced for

$$\frac{v_\phi}{\Lambda} \sim \frac{v_T}{\Lambda} \sim \mathcal{O}(\lambda^2). \quad (4.3)$$

The driving superpotential w_d^v involving the flavons of the neutrino sector reads

$$w_d^v = g_1(\varphi_S^0(\varphi_S\varphi_S)_3)_1 + g_2(\varphi_S^0(\eta\varphi_S)_3)_1 + g_3\xi^0(\varphi_S\varphi_S)_1 + g_4\xi^0(\eta\eta)_1 + M_\eta(\eta^0\eta)_1 \\ + g_5(\eta^0(\chi\chi)_2)_1 + g_6\rho^0(\chi\chi)_1 + g_7\rho^0\xi^2 + g_8\sigma^0(\chi\varphi_S)_1, \quad (4.4)$$

where all the coupling g_i and mass parameter M_η are real because of the imposed generalised CP symmetry. Since the contraction $(\varphi_S\varphi_S)_3$ vanishes due to the antisymmetry of the associated S_4 Clebsch–Gordan coefficients, the first term proportional to g_1 gives null contribution. In the SUSY limit, the vacuum configuration is determined by the vanishing of the derivative of the driving superpotential w_d^v with respect to each component of the driving fields. The minimization equations for the vacuum take the following form:

$$\begin{aligned} \frac{\partial w_d^v}{\partial \varphi_{S_1}^0} &= g_2(\eta_1\varphi_{S_2} - \eta_2\varphi_{S_3}) = 0, \\ \frac{\partial w_d^v}{\partial \varphi_{S_2}^0} &= g_2(\eta_1\varphi_{S_1} - \eta_2\varphi_{S_2}) = 0, \\ \frac{\partial w_d^v}{\partial \varphi_{S_3}^0} &= g_2(\eta_1\varphi_{S_3} - \eta_2\varphi_{S_1}) = 0, \\ \frac{\partial w_d^v}{\partial \xi^0} &= g_3(\varphi_{S_1}^2 + 2\varphi_{S_2}\varphi_{S_3}) + 2g_4\eta_1\eta_2 = 0, \\ \frac{\partial w_d}{\partial \eta_1^0} &= M_\eta\eta_2 + g_5(\chi_3^2 + 2\chi_1\chi_2) = 0, \\ \frac{\partial w_d}{\partial \eta_2^0} &= M_\eta\eta_1 + g_5(\chi_2^2 + 2\chi_1\chi_3) = 0, \\ \frac{\partial w_d}{\partial \rho^0} &= g_6(\chi_1^2 + 2\chi_2\chi_3) + g_7\xi^2 = 0, \\ \frac{\partial w_d}{\partial \sigma^0} &= g_8(\chi_1\varphi_{S_1} + \chi_2\varphi_{S_3} + \chi_3\varphi_{S_2}) = 0. \end{aligned} \quad (4.5)$$

The solution to these equation are

$$\langle \varphi_S \rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} v_S, \quad \langle \eta \rangle = \begin{pmatrix} 1 \\ 1 \end{pmatrix} v_\eta, \quad \langle \chi \rangle = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} v_\chi, \quad \langle \xi \rangle = v_\xi. \quad (4.6)$$

The VEVs v_S , v_η , v_χ and v_ξ are related by

$$v_S^2 = -\frac{84g_5^2g_7^2}{6g_3g_6^2M_\eta^2}v_\xi^4, \quad v_\eta = -\frac{g_5g_7}{2g_6M_\eta}v_\xi^2, \quad v_\chi^2 = \frac{g_7}{2g_6}v_\xi^2, \quad (4.7)$$

where v_ξ parameterizes a flat direction in the driving superpotential w_d^v , and it is in general complex. It is straightforward to check that the VEVs of the flavon fields φ_S , η and ξ preserve the remnant K_4 subgroup generated by Z_2^S and Z_2^{SU} , while the VEV of χ is invariant only under the action of Z_2^{SU} . In our model presented below, φ_S and η couple with the right-handed neutrino at LO, as shown in Eq. (4.12). The resulting lepton mixing is of the tri-bimaximal form. The flavons χ and ξ enter into the neutrino sector at NLO, and the LO residual K_4 symmetry is further broken down to Z_2^{SU} . As a result, the NLO contributions modify the LO tri-bimaximal mixing into TM_1 pattern. In order to achieve the measured size of $\theta_{13} \simeq \lambda/\sqrt{2}$ [38,39], we could choose

$$\frac{v_S}{\Lambda} \sim \frac{v_\eta}{\Lambda} \sim \frac{v_\chi}{\Lambda} \sim \frac{v_\xi}{\Lambda} \sim \mathcal{O}(\lambda). \quad (4.8)$$

Consequently the NLO corrections are suppressed by a factor λ with respect to the LO contributions, and therefore the reactor angle is of the correct order λ . Note that the VEVs of the flavon fields in the neutrino and the charged lepton sectors are chosen to be of different order of magnitude: $\lambda\Lambda$ v.s. $\lambda^2\Lambda$, please see Eq. (4.3) and Eq. (4.8). This mild hierarchy can be accommodated because these two sets of VEVs depend on different model parameters.

4.2. Leading order results

The superpotential for the charged lepton masses, which is allowed by the symmetry, is given by

$$\begin{aligned} w_l = & \frac{y_\tau}{\Lambda}(l\varphi_T)_1\tau^c h_d + \frac{y_{\mu 1}}{\Lambda^2}(l(\varphi_T\varphi_T)\mathbf{3})_1\mu^c h_d + \frac{y_{\mu 2}}{\Lambda^2}(l(\phi\varphi_T)\mathbf{3})_1\mu^c h_d \\ & + \frac{y_{e1}}{\Lambda^3}(l\varphi_T)_1(\varphi_T\varphi_T)_1e^c h_d + \frac{y_{e2}}{\Lambda^3}((l\varphi_T)_2(\varphi_T\varphi_T)_2)_1e^c h_d \\ & + \frac{y_{e3}}{\Lambda^3}((l\varphi_T)\mathbf{3}'(\varphi_T\varphi_T)\mathbf{3}')_1e^c h_d \\ & + \frac{y_{e4}}{\Lambda^3}((l\varphi_T)\mathbf{3}(\varphi_T\varphi_T)\mathbf{3})_1e^c h_d + \frac{y_{e5}}{\Lambda^3}((l\phi)\mathbf{3}'(\varphi_T\varphi_T)\mathbf{3}')_1e^c h_d \\ & + \frac{y_{e6}}{\Lambda^3}((l\phi)\mathbf{3}(\varphi_T\varphi_T)\mathbf{3})_1e^c h_d \\ & + \frac{y_{e7}}{\Lambda^3}((l\varphi_T)_2(\phi\phi)_2)_1e^c h_d + \frac{y_{e8}}{\Lambda^3}(l\varphi_T)_1(\phi\phi)_1e^c h_d + \dots, \end{aligned} \quad (4.9)$$

which is identical to the corresponding superpotential of Model 1 shown in Eq. (3.14). After electroweak and flavor symmetry breaking in the way of Eq. (4.2), we obtain a diagonal charged lepton mass matrix:

$$m_l = \begin{pmatrix} y_e \frac{v_T^2}{\Lambda^2} & 0 & 0 \\ 0 & y_\mu \frac{v_T}{\Lambda} & 0 \\ 0 & 0 & y_\tau \end{pmatrix} \frac{v_T}{\Lambda} v_d, \quad (4.10)$$

where y_e and y_μ are the results of the different contributions of the y_{e_i} and y_{μ_i} respectively with

$$y_\mu = 2y_{\mu 1} + y_{\mu 2} \frac{v_\phi}{v_T}, \quad y_e = y_{e 2} - 2y_{e 3} + 2y_{e 5} \frac{v_\phi}{v_T} + y_{e 7} \frac{v_\phi^2}{v_T^2}. \quad (4.11)$$

Now we turn to the neutrino sector, The LO superpotential relevant to the neutrino masses is of the form

$$w_\nu = y(l\nu^c)_1 h_u + y_1((\nu^c\nu^c)_3 \varphi_S)_1 + y_2((\nu^c\nu^c)_2 \eta)_1, \quad (4.12)$$

where all the three couplings y , y_1 and y_2 are real because of the generalised CP symmetry. We can easily read out the Dirac neutrino mass matrix as

$$m_D = yv_u \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (4.13)$$

Given the vacuum of the flavons φ_S and η shown in Eq. (4.6), the mass matrix for the right-handed neutrino takes the form

$$m_M = a \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + b \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}, \quad (4.14)$$

where $a = y_1 v_S$ and $b = y_2 v_\eta$. The light neutrino mass matrix is given by the see-saw formula, yielding

$$m_\nu = -m_D m_M^{-1} m_D^T = U_{TB} \text{diag}(m_1, m_2, m_3) U_{TB}^T. \quad (4.15)$$

That is to say the LO lepton flavor mixing is the tri-bimaximal pattern. The reason is that the VEVs of φ_S and η break the S_4 family symmetry into a residual $K_4 \cong Z_2^S \times Z_2^{SU}$ subgroup, i.e. the vacuum of φ_S and η in Eq. (4.6) is invariant under both Z_2^S and Z_2^{SU} . Furthermore, the light neutrino masses $m_{1,2,3}$ in Eq. (4.15) are given by

$$m_1 = \frac{y^2 v_u^2}{-3y_1 v_S + y_2 v_\eta}, \quad m_2 = -\frac{y^2 v_u^2}{2y_2 v_\eta}, \quad m_3 = -\frac{y^2 v_u^2}{3y_1 v_S + y_2 v_\eta}. \quad (4.16)$$

It is interesting to note that the following sum rule is satisfied

$$\frac{1}{m_3} - \frac{1}{m_1} = \frac{1}{m_2}. \quad (4.17)$$

Since the VEVs v_S and v_η are related through Eq. (4.7), the phase different between v_S and v_η is fixed to discrete values $0, \pi$ or $\pm\pi/2$ for the product $g_3 g_4 < 0$ or $g_3 g_4 > 0$, respectively. Moreover, the phase of v_ξ can be absorbed by redefining the right-handed neutrino fields, therefore we can set v_ξ to be real, and then another VEV v_S would be real or purely imaginary. For the case of v_S being imaginary, Eq. (4.16) implies that the light neutrino masses are degenerate, i.e. $|m_1| = |m_3|$. Therefore this case is not phenomenologically viable, and we shall choose v_S to be real (or v_S and v_η have the same phase up to relative sign) in the following. Then the neutrino mass-squared differences are predicted to be

$$\Delta m_{sol}^2 \equiv |m_2|^2 - |m_1|^2 = \frac{3(3x+1)(x-1)}{4(3x-1)^2} \left(\frac{y^2 v_u^2}{y_2 v_\eta} \right)^2,$$

$$\Delta m_{atm}^2 \equiv |m_3|^2 - |m_1|^2 = \frac{-12x}{(9x^2-1)^2} \left(\frac{y^2 v_u^2}{y_2 v_\eta} \right)^2, \quad \text{for NO},$$

Table 4

The predictions for the Majorana phases, the light neutrino masses $|m_i|$ ($i = 1, 2, 3$) and the effective mass $|m_{\beta\beta}|$ of the neutrinoless double-beta decay at LO.

x	α_{21}	α_{31}	$ m_1 $ (meV)	$ m_2 $ (meV)	$ m_3 $ (meV)	$ m_{\beta\beta} $ (meV)	Mass order
-0.5173	π	0	10.891	13.897	50.355	2.628	NO
1.0079	0	0	55.913	56.576	28.121	56.134	IO

$$\Delta m_{atm}^2 \equiv |m_2|^2 - |m_3|^2 = \frac{3(3x - 1)(x + 1)}{4(3x + 1)^2} \left(\frac{y^2 v_u^2}{y_2 v_\eta} \right)^2, \quad \text{for IO,} \tag{4.18}$$

where $x = \frac{y_1 v_S}{y_2 v_\eta}$ is a real parameter. Furthermore, the effective mass parameter $|m_{\beta\beta}|$ for the neutrinoless doublet beta is given by

$$|m_{\beta\beta}| = \left| \frac{x + 1}{2(3x - 1)} \right| \left| \frac{y^2 v_u^2}{y_2 v_\eta} \right|. \tag{4.19}$$

Since the solar neutrino mass squared difference Δm_{sol}^2 is positive, we have $x > 1$ or $x < -\frac{1}{3}$ from Eq. (4.18). By further inspecting the atmospheric neutrino mass squared difference Δm_{atm}^2 , we find that neutrino spectrum is normal ordering (NO) for $x < -\frac{1}{3}$ and inverted order (IO) for $x > 1$. Taking the best fit values $\Delta m_{sol}^2 = 7.45 \times 10^{-5} \text{ eV}^2$ and $\Delta m_{atm}^2 = 2.417(2.410) \times 10^{-3} \text{ eV}^2$ for NO (IO) spectrum from Ref. [13], we get two solutions for the ratio x (one for normal ordering and another for inverted ordering):

$$x = -0.5173, \quad 1.0079. \tag{4.20}$$

The corresponding predictions for the Majorana phases, the light neutrino masses and $|m_{\beta\beta}|$ are presented in Table 4.

4.3. Next-to-leading-order corrections

Since the LO tri-bimaximal mixing pattern leads to a vanishing reactor angle θ_{13} which has been definitely excluded by the experimental measurements, NLO corrections are needed to achieve agreement with the present data. In this section, we shall address the NLO corrections indicated by higher dimensional operators compatible with all the symmetries of the model. As we shall show, the NLO contributions break the remnant family $K_4 \cong Z_2^S \times Z_2^{SU}$ in the neutrino sector down to Z_2^{SU} . As a result, a non-zero θ_{13} is generated and it is naturally smaller than θ_{12} and θ_{23} which arise at LO.

In the following, we first discuss the NLO corrections to the charged lepton sector. For the driving superpotential w_d^l , the most relevant subleading operators can be written as

$$\delta w_d^l = (\varphi_T^0 \Psi_l^2 \Psi_\nu^2 \Psi_\nu')_1 / \Lambda^3 + (\zeta^0 \Psi_l^2 \Psi_\nu^2 \Psi_\nu')_1 / \Lambda^3, \tag{4.21}$$

where we have suppressed all dimensionless coupling constants, and all the possible S_4 contractions should be considered with $\Psi_l = \{\varphi_T, \phi\}$, $\Psi_\nu = \{\varphi_S, \eta\}$ and $\Psi_\nu' = \{\chi, \xi\}$. These operators are suppressed by $\langle \Psi_\nu \rangle^2 \langle \Psi_\nu' \rangle / \Lambda^3 \sim \lambda^3$ compared to LO terms in w_d^l of Eq. (4.1). Hence the sub-leading corrections to the VEVs of the φ_T and ϕ appear at the relative order λ^3 such that their vacuum configurations at NLO can be parameterized as

$$\langle \varphi_T \rangle = v_T \begin{pmatrix} \epsilon_1 \lambda^3 \\ 1 + \epsilon_2 \lambda^3 \\ \epsilon_3 \lambda^3 \end{pmatrix}, \quad \langle \phi \rangle = v_\phi \begin{pmatrix} \epsilon_4 \lambda^3 \\ 1 \end{pmatrix} \quad (4.22)$$

where the coefficients ϵ_i ($i = 1, 2, 3, 4$) have absolute value of order one and are generally complex due to the undetermined phase of v_ξ . Note that the shift of the second component of ϕ has been absorbed into the redefinition of the undetermined parameters v_ϕ . The subleading corrections to the charged lepton superpotential w_l take the form

$$\delta w_l = (l\Psi_l\Psi_\nu^2\Psi_\nu')_1 h_d \tau^c / \Lambda^4 + (l\Psi_l^2\Psi_\nu^2\Psi_\nu')_1 h_d \mu^c / \Lambda^5 + (l\Psi_l^3\Psi_\nu^2\Psi_\nu')_1 h_d e^c / \Lambda^6, \quad (4.23)$$

where the dimensionless coupling constants are omitted. The charged lepton mass matrix is obtained by adding the contributions of this set of high dimensional operators evaluated with the insertion of the LO VEVs of Eqs. (4.2), (4.6), to those of the LO superpotential in Eq. (4.9) evaluated with the NLO vacuum configuration in Eq. (4.22). We find that each element of the charged lepton mass matrix receives corrections from both the subleading operators δw_l in Eq. (4.23) and the shifted vacuum alignment of Eq. (4.22). As a consequence, its off-diagonal elements become non-zero and are all suppressed by λ^3 with respect to the diagonal ones. Therefore the charged lepton mass matrix including subleading corrections can be written as

$$m_l^{NLO} = \begin{pmatrix} m_e & \lambda^3 m_\mu & \lambda^3 m_\tau \\ \lambda^3 m_e & m_\mu & \lambda^3 m_\tau \\ \lambda^3 m_e & \lambda^3 m_\mu & m_\tau \end{pmatrix}. \quad (4.24)$$

Its contribution to the lepton mixing angles is of order λ^3 and can be safely neglected. Since the off-diagonal elements are quite small in particular the (2, 1) and (3, 1) entries, perturbatively diagonalizing the above NLO charged lepton mass matrix m_l^{NLO} reveals that the NLO corrections to the charged lepton masses are of relative order λ^6 , and hence they are negligible as well.

Next, we turn to discuss the NLO corrections in the neutrino sector. The NLO contributions to the driving superpotential w_d^v is suppressed by one power of $1/\Lambda$ with respect to the LO terms in Eq. (4.4), and it takes the form²

$$\begin{aligned} \delta w_d^v = & \frac{h_1}{\Lambda} ((\varphi_S^0 \varphi_S)_2 (\chi \chi)_2)_1 + \frac{h_2}{\Lambda} ((\varphi_S^0 \varphi_S)_3 (\chi \chi)_3)_1 + \frac{h_3}{\Lambda} ((\varphi_S^0 \varphi_S)_{3'} (\chi \chi)_{3'})_1 \\ & + \frac{h_4}{\Lambda} \xi ((\varphi_S^0 \varphi_S)_{3'} \chi)_1 + \frac{h_5}{\Lambda} ((\varphi_S^0 \eta)_3 (\chi \chi)_3)_1 + \frac{h_6}{\Lambda} ((\varphi_S^0 \eta)_{3'} (\chi \chi)_{3'})_1 \\ & + \frac{h_7}{\Lambda} ((\varphi_S^0 \eta)_{3'} \chi)_1 \xi + \frac{h_8}{\Lambda} \xi^0 (\varphi_S (\chi \chi)_{3'})_1 + \frac{h_9}{\Lambda} \xi^0 \xi (\varphi_S \chi)_1 \\ & + \frac{h_{10}}{\Lambda} \xi^0 (\eta (\chi \chi)_2)_1 + \frac{h_{11}}{\Lambda} \sigma^0 (\chi (\chi \chi)_{3'})_1 + \frac{h_{12}}{\Lambda} \sigma^0 \xi (\chi \chi)_1 + \frac{h_{13}}{\Lambda} \sigma^0 \xi^3, \end{aligned} \quad (4.25)$$

where all the couplings h_i are again real because of the generalised CP symmetry. Repeating the minimization procedure of Section 4.1, we find that the LO vacuum configuration is modified into

$$\langle \varphi_S \rangle = v'_S \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \delta v_S \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}, \quad \langle \chi \rangle = v_\chi \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + \delta v_\chi \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad (4.26)$$

² The subleading corrections to the terms proportional to η^0 and ρ^0 are of the form $(\eta^0 \Psi_\nu^3 \Psi_\nu')_1 / \Lambda^2$ and $(\rho^0 \Psi_\nu^3 \Psi_\nu')_1 / \Lambda^2$, which are suppressed by $1/\Lambda^2$ instead of $1/\Lambda$.

with

$$\begin{aligned}
 v'_S - v_S &= -\left(\frac{h_8}{g_3} + \frac{h_{10} v_\eta}{3g_3 v_S}\right) \frac{v_\chi^2}{\Lambda}, \\
 \delta v_S &= \left(\frac{h_4 v_S}{g_2 v_\eta} - \frac{h_7}{g_2}\right) \frac{v_\chi v_\xi}{\Lambda}, \\
 \delta v_\chi &= -\frac{h_{13} v_\xi}{3g_8 v_S} \frac{v_\xi^2}{\Lambda} + \frac{2v_\xi}{3v_S} \left(\frac{h_{12}}{g_8} - \frac{h_7}{g_2} + \frac{h_4 v_S}{g_2 v_\eta}\right) \frac{v_\chi^2}{\Lambda},
 \end{aligned} \tag{4.27}$$

and the vacuum of η doesn't acquires non-trivial shifts at this order. Obviously the shifts $v'_S - v_S$, δv_S and δv_χ are suppressed with respect to the LO VEVs v_S and v_χ by a factor λ . Notice that the shifted vacuum of φ_S and χ in Eq. (4.26) is the most general form of VEV invariant under the Z_2^{SU} subgroup. The reason is that the NLO terms δw_d^y of Eq. (4.25) only involve the neutrino flavons φ_S , η , χ and ξ whose LO VEVs leave Z_2^{SU} invariant.

From Section 4.2, we know that the VEVs v_S , v_η and v_ξ^2 have to share the same phase, i.e. the product $g_3 g_4 < 0$ is needed otherwise the light neutrino mass spectrum would be partially degenerate. Furthermore, Eq. (4.6) implies that the phase different between v_χ and v_ξ is $0, \pi$ or $\pm \frac{\pi}{2}$ for $g_6 g_7 > 0$ or $g_6 g_7 < 0$, respectively. As a result, v'_S and v_S carry the same phase. Since it is always possible to absorb the phase of v_ξ by a redefinition of the matter fields, we can take v_ξ to be real without loss of generality. Then v'_S , v_η and v_χ^2 would be real, while v_χ and δv_S can be real or purely imaginary depending on $g_6 g_7 > 0$ or $g_6 g_7 < 0$.

Now we come to the NLO corrections to the LO neutrino superpotential w_v in Eq. (4.12). The higher order corrections to the neutrino Dirac mass are of the form

$$(lv^c \Psi_v^2 \Psi'_v)_1 h_u / \Lambda^3. \tag{4.28}$$

The corresponding contributions are suppressed by λ^3 compared to the LO term $y(lv^c)_1 h_u$. Such small corrections have a tiny impact for the neutrino mass matrix and lepton mixing parameters, and therefore can be neglected. The NLO corrections to the RH neutrino Majorana mass terms are

$$\begin{aligned}
 \delta w_v &= s_1 (v^c v^c)_1 (\chi \chi)_1 / \Lambda + s_2 ((v^c v^c)_2 (\chi \chi)_2)_1 / \Lambda + s_3 ((v^c v^c)_3 (\chi \chi)_3)_1 / \Lambda \\
 &\quad + s_4 ((v^c v^c)_3 (\chi \chi)_3')_1 / \Lambda + s_5 \xi ((v^c v^c)_3 \chi)_1 / \Lambda + s_6 \xi^2 (v^c v^c)_1 / \Lambda.
 \end{aligned} \tag{4.29}$$

The resulting corrections to the RH neutrino mass matrix m_M can be obtained by inserting the LO vacuum of χ and ξ in Eq. (4.6) into these operators. Another source of corrections to m_M arises from the LO superpotential w_v in Eq. (4.12) evaluated with the NLO VEVs of Eq. (4.26). Adding the two contributions, we obtained the corrected RH neutrino mass matrix as

$$m_M^{NLO} = a \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + b \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + c \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + d \begin{pmatrix} 0 & 1 & -1 \\ 1 & 2 & 0 \\ -1 & 0 & -2 \end{pmatrix}, \tag{4.30}$$

with

$$\begin{aligned}
 a &= y_1 v'_S + 2s_4 v_\chi^2 / \Lambda, & b &= y_2 v_\eta + s_2 v_\chi^2 / \Lambda, & c &= s_6 v_\xi^2 / \Lambda - 2s_1 v_\chi^2 / \Lambda, \\
 d &= y_1 \delta v_S + s_5 v_\chi v_\xi / \Lambda = \left[y_1 \left(\frac{h_4 v_S}{g_2 v_\eta} - \frac{h_7}{g_2} \right) + s_5 \right] \frac{v_\chi v_\xi}{\Lambda},
 \end{aligned} \tag{4.31}$$

where parameters a and b have been redefined to include the NLO contributions. Note that c and d arise from the NLO contributions, and they are suppressed by a factor λ with respect to a and b , i.e.

$$a, b \sim \lambda A, \quad c, d \sim \lambda^2 A. \quad (4.32)$$

Applying the see-saw relation, the light neutrino mass matrix at NLO takes the form

$$\begin{aligned} m_\nu^{NLO} &= -m_D(m_M^{NLO})^{-1}m_D^T, \\ &= \alpha \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix} + \beta \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \\ &\quad + \gamma \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + \delta \begin{pmatrix} 0 & 1 & -1 \\ 1 & 2 & 0 \\ -1 & 0 & -2 \end{pmatrix}. \end{aligned} \quad (4.33)$$

It is the most general neutrino mass matrix invariant under residual family symmetry $G_\nu = Z_2^{SU} = \{1, SU\}$, as shown in Eq. (2.15). The parameters α , β , γ and δ are given by

$$\begin{aligned} \alpha &= \frac{-a(2b+c) + d^2}{(3a-b+c)[(3a+b-c)(2b+c) - 6d^2]}, \\ \beta &= \frac{-3a^2 - b^2 + c^2 + 2d^2}{(3a-b+c)[(3a+b-c)(2b+c) - 6d^2]}, \\ \gamma &= -\frac{3a^2 + b(c-b) + d^2}{(3a-b+c)[(3a+b-c)(2b+c) - 6d^2]}, \\ \delta &= -\frac{d}{(3a+b-c)(2b+c) - 6d^2}, \end{aligned} \quad (4.34)$$

where the overall factor $y^2 v_u^2$ is omitted here. Because the theory is required to be invariant under the generalised CP transformations, the phases of the model parameters are strongly constrained. The vacuum alignment of Eq. (4.7) implies that the phase different between v_χ and v_ξ is 0, π or $\pi/2$ for $g_6 g_7 > 0$ and $g_6 g_7 < 0$ respectively. Further recalling that v_s and v_ξ^2 should have a common phase (up to relative sign) to avoid degenerate light neutrino masses at LO. Therefore, a , b and c are real while d is real or imaginary after the unphysical phase of v_ξ is extracted. As a result, α , β and γ in Eq. (4.30) are real parameters whereas δ can be real or purely imaginary. In the following, we discuss the two cases one after another.

Firstly, we consider the case that v_χ is real, which corresponds to the parameter domain of $g_6 g_7 > 0$. We can check that the remnant CP symmetry in the neutrino sector is $H_{CP}^\nu = \{\rho_{\mathbf{r}}(1), \rho_{\mathbf{r}}(SU)\}$ in this case. All the four parameters α , β , γ and δ are real. This is exactly the case (I) of model-independent analysis in Section 2. Remembering that the subleading operators in the charged lepton sector induce corrections to the lepton mixing angles as small as λ^3 . Hence, the lepton flavor mixing is determined by the neutrino sector. From Section 2, we know that the resulting lepton mixing matrix is

$$U_{PMNS} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{\cos \theta}{\sqrt{3}} & \frac{\sin \theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos \theta}{\sqrt{3}} + \frac{\sin \theta}{\sqrt{2}} & -\frac{\cos \theta}{\sqrt{2}} + \frac{\sin \theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos \theta}{\sqrt{3}} - \frac{\sin \theta}{\sqrt{2}} & \frac{\cos \theta}{\sqrt{2}} + \frac{\sin \theta}{\sqrt{3}} \end{pmatrix}, \quad (4.35)$$

with

$$\tan 2\theta = \frac{-2\sqrt{6}\delta}{3\alpha - 2\beta - \gamma} = \frac{2\sqrt{6}d}{3a - b - 2c} \sim \mathcal{O}(\lambda). \quad (4.36)$$

The lepton mixing angles are given by

$$\begin{aligned} \sin \theta_{13} &= \left| \frac{\sin \theta}{\sqrt{3}} \right| \simeq \left| \frac{\sqrt{2}d}{3a - b - 2c} \right| \sim \mathcal{O}(\lambda), \\ \sin^2 \theta_{12} &\simeq \frac{1}{3} + \mathcal{O}(\lambda^2), \quad \sin^2 \theta_{23} \simeq \frac{1}{2} \pm \frac{2d}{3a - b - 2c}. \end{aligned} \quad (4.37)$$

We see that the reactor angle θ_{13} is predicted to be of the correct order of λ , and thus experimentally preferred value could be achieved. The solar mixing angle retains its tri-bimaximal value to the first order of λ , and the atmospheric angle can deviate from its maximal mixing value of 45° . As a consequence, the deviation of the atmospheric angle from maximal mixing, indicated by the latest global fits, can be produced. In addition, we find a simple sum rule $\sin^2 \theta_{23} \simeq 0.5 \pm \sqrt{2} \sin \theta_{13}$. This relation might be testable in the near future as soon as the experimental uncertainties for θ_{23} are reduced. Furthermore, since the light neutrino mass matrix is real, there is no CP violation in this case, both the Dirac CP phase and the Majorana CP phases are 0 or π .

Then we consider the remaining case of v_χ being purely imaginary, i.e. the phase different between v_χ and v_ξ is $\pm \frac{\pi}{2}$. This scenario can be realized in the parameter domain $g_6 g_7 < 0$. The generalised CP symmetry is broken down to $H_{CP}^\nu = \{\rho_r(S), \rho_r(U)\}$ in the neutrino sector. This corresponds to the case (II) of Section 2. The resulting parameters α, β, γ are real and δ is imaginary. The lepton mixing matrix is of the form

$$U_{PMNS} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{\cos \theta}{\sqrt{3}} & \frac{\sin \theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos \theta}{\sqrt{3}} + \frac{i \sin \theta}{\sqrt{2}} & -\frac{i \cos \theta}{\sqrt{2}} + \frac{\sin \theta}{\sqrt{3}} \\ -\frac{1}{\sqrt{6}} & \frac{\cos \theta}{\sqrt{3}} - \frac{i \sin \theta}{\sqrt{2}} & \frac{i \cos \theta}{\sqrt{2}} + \frac{\sin \theta}{\sqrt{3}} \end{pmatrix}, \quad (4.38)$$

with

$$\tan 2\theta = \frac{2i\sqrt{6}\delta}{3(\alpha + \gamma)} = \frac{2i\sqrt{6}d}{3(a + b)} \sim \mathcal{O}(\lambda). \quad (4.39)$$

Consequently the three mixing angles θ_{13}, θ_{12} and θ_{23} are modified to

$$\sin \theta_{13} \simeq \left| \frac{\sqrt{2}d}{3(a + b)} \right| \sim \mathcal{O}(\lambda), \quad \sin^2 \theta_{12} = \frac{1}{3} + \mathcal{O}(\lambda^2), \quad \sin^2 \theta_{23} = \frac{1}{2}. \quad (4.40)$$

It is noteworthy that we obtain maximal Dirac CP violation $\delta_{CP} = \pm \pi/2$ in this case while the Majorana CP phases are still trivial with $\sin \alpha_{21} = \sin \alpha_{13} = 0$. In short summary, our model produces the tri-bimaximal mixing at LO, which is further broken down to trimaximal TM_1 mixing by NLO contributions. Depending on the coupling product $g_6 g_7$ being positive or negative, the two cases arising from the model independent analysis can be realized.

5. Conclusions

The measurement of sizable reactor mixing angle θ_{13} has opened up the possibility of measuring leptonic CP violations. In particular, the measurement of Dirac CP phase is one of the

primary goals of next generation neutrino oscillation experiments. On the theoretical side, the origin of CP violation remains a mystery. Extending family symmetry to include generalised CP symmetry together with its spontaneous breaking is a promising framework to predict both mixing angles and CP phases.

In this work, we analyse the interplay of generalised CP symmetry and the S_4 family symmetry. Firstly we perform a model independent analysis of the possible lepton mixing matrices and the corresponding lepton mixing parameters, which arise from the symmetry breaking of $S_4 \times H_{CP}$ into $Z_3^T \times H_{CP}^I$ in the charged lepton sector and $Z_2^{SU} \times H_{CP}^V$ in the neutrino sector. We find that the lepton flavor mixing is of the TM_1 form and the Dirac CP can be vanishing or maximally broken while the Majorana CP is trivial with $\sin \alpha_{21} = \sin \alpha_{31} = 0$.

Furthermore, we construct two models to realize the above model independent results based on S_4 family symmetry and the generalised CP symmetry. The two models differ in the neutrino sectors. In the first model, the flavon fields enter in the neutrino Dirac mass term instead of the Majorana mass term for right-handed neutrinos at LO. The resulting light neutrino mass matrix is predicted to depend on three real parameters, and therefore the absolute neutrino masses and the effective mass $|m_{\beta\beta}|$ for neutrinoless double beta decay are completely fixed after considering the constraints from the measured values of the neutrino mass squared differences Δm_{sol}^2 and Δm_{atm}^2 and the reactor angle θ_{13} . The lepton mixing matrix is the TM_1 pattern, and the subleading corrections are small enough to be negligible. In the case of $g_3 g_4 (g_3 g_5 + g_2 g_6) > 0$, the Dirac CP phase δ_{CP} is 0 or π , and neutrino mass spectrum can be normal ordering or inverted ordering. For the case of $g_3 g_4 (g_3 g_5 + g_2 g_6) < 0$, the Dirac CP is maximal $\delta_{CP} = \pm\pi/2$, and the neutrino mass spectrum can only be normal ordering.

In the second model, the S_4 family symmetry is broken down to $Z_2^S \times Z_2^{SU}$ in the neutrino sector at LO, and therefore the LO lepton mixing is of the tri-bimaximal form. NLO correction terms break the remnant symmetry $Z_2^S \times Z_2^{SU}$ into Z_2^{SU} , as a result, the TM_1 mixing is produced and the relative smallness of θ_{13} with respect to θ_{12} and θ_{23} is explained. Depending on the product $g_6 g_7$ being positive or negative, the Dirac CP is predicted to be conserved or maximally broken. Moreover, we have shown that the desired vacuum alignment together with their phase structure can be achieved.

Acknowledgements

One of the author (G.J.D.) is grateful to Stephen F. King, Christoph Luhn and Alexander J. Stuart for stimulating discussions on generalised CP symmetry. G.J.D. would also like to thank Stephen F. King and the School of Physics and Astronomy at the University of Southampton for hospitality during his visit, where part of this work was done. The research was partially supported by the National Natural Science Foundation of China under Grant Nos. 11275188 and 11179007.

Appendix A. Group Theory of S_4

S_4 is the permutation group of order 4 with 24 elements, and it has been widely used as a family symmetry. In this work, we shall follow the conventions and notations of Refs. [19,40], where S_4 is expressed in terms of three generators S , T and U . These three generators satisfy the multiplication rules:

$$S^2 = T^3 = U^2 = (ST)^3 = (SU)^2 = (TU)^2 = (STU)^4 = 1. \quad (\text{A.1})$$

Table 5

The representation matrices of the generators S , T and U for the five irreducible representations of S_4 in the chosen basis, where $\omega = e^{2\pi i/3}$.

	S	T	U
$\mathbf{1}, \mathbf{1}'$	1	1	± 1
$\mathbf{2}$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
$\mathbf{3}, \mathbf{3}'$	$\frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}$	$\mp \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$

Note that the generators S and T alone generate the group A_4 , while the generated group by T and U is S_3 . The S_4 group elements can be divided into 5 conjugacy classes

$$\begin{aligned}
 1C_1 &= \{1\}, \\
 3C_2 &= \{S, TST^2, T^2ST\}, \\
 6C'_2 &= \{U, TU, SU, T^2U, STSU, ST^2SU\}, \\
 8C_3 &= \{T, ST, TS, STS, T^2, ST^2, T^2S, ST^2S\}, \\
 6C_4 &= \{STU, TSU, T^2SU, ST^2U, TST^2U, T^2STU\},
 \end{aligned} \tag{A.2}$$

where the conjugacy class is denoted by kC_n , k is the number of elements belonging to it, and the subscript n is the order of the elements contained in it. As a result of these conjugacy classes and the theorems that prove that the number of inequivalent irreducible representations is equal to the number of conjugacy classes and the sum of the squares of the dimensions of the inequivalent irreducible representations must be equal to the order of the group, it is easy to see that S_4 has two singlet irreducible representations $\mathbf{1}$ and $\mathbf{1}'$, one two-dimensional representation $\mathbf{2}$ and two three-dimensional irreducible representations $\mathbf{3}$ and $\mathbf{3}'$. In this work, we shall work in the basis where the representation matrix of the generator T is diagonal. As a result, the charged lepton mass matrix would be diagonal if the remnant subgroup $Z_3^T \equiv \{1, T, T^2\}$ is preserved in the charged lepton sector. The explicit forms of the representation matrix for the three generators are listed in Table 5, and hence the chosen basis coincides with that of Ref. [19]. The character table of S_4 group follows immediately, as shown in Table 6. Moreover, the Kronecker products between different irreducible representations are as follows

$$\begin{aligned}
 \mathbf{1} \otimes \mathbf{R} &= \mathbf{R}, & \mathbf{1}' \otimes \mathbf{1}' &= \mathbf{1}, & \mathbf{1}' \otimes \mathbf{2} &= \mathbf{2}, & \mathbf{1}' \otimes \mathbf{3} &= \mathbf{3}', & \mathbf{1}' \otimes \mathbf{3}' &= \mathbf{3}, \\
 \mathbf{2} \otimes \mathbf{2} &= \mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{2}, & \mathbf{2} \otimes \mathbf{3} &= \mathbf{2} \otimes \mathbf{3}' = \mathbf{3} \otimes \mathbf{3}', \\
 \mathbf{3} \otimes \mathbf{3} &= \mathbf{3}' \otimes \mathbf{3}' = \mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3} \oplus \mathbf{3}', & \mathbf{3} \otimes \mathbf{3}' &= \mathbf{1}' \oplus \mathbf{2} \oplus \mathbf{3} \oplus \mathbf{3}'
 \end{aligned} \tag{A.3}$$

where \mathbf{R} stands for any irreducible representation of S_4 .

In the end, we present the Clebsch–Gordan (CG) coefficients in the chosen basis. All the CG coefficients can be reported in the form of $\alpha \otimes \beta$, α_i denotes the element of the left base vectors α , and β_i is the element of the right base vectors β . For the product of the singlet $\mathbf{1}'$ with a doublet or a triplet, we have

$$\mathbf{1}' \otimes \mathbf{2} = \mathbf{2} = \alpha \begin{pmatrix} \beta_1 \\ -\beta_2 \end{pmatrix}, \quad \mathbf{1}' \otimes \mathbf{3} = \mathbf{3}' = \alpha \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix},$$

Table 6

Character table of the group S_4 , where G denotes the representative element of each conjugacy class.

Classes	$1C_1$	$3C_2$	$6C'_2$	$8C_3$	$6C_4$
G	1	S	U	T	STU
1	1	1	1	1	1
1'	1	1	-1	1	-1
2	2	2	0	-1	0
3	3	-1	-1	0	1
3'	3	-1	1	0	-1

$$\mathbf{1}' \otimes \mathbf{3}' = \mathbf{3} = \alpha \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}. \tag{A.4}$$

The CG coefficients for the products involving the doublet representation **2** are found to be

$$\mathbf{2} \otimes \mathbf{2} = \mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{2}, \quad \text{with} \quad \begin{cases} \mathbf{1} = \alpha_1\beta_2 + \alpha_2\beta_1 \\ \mathbf{1}' = \alpha_1\beta_2 - \alpha_2\beta_1 \\ \mathbf{2} = \begin{pmatrix} \alpha_2\beta_2 \\ \alpha_1\beta_1 \end{pmatrix} \end{cases}$$

$$\mathbf{2} \otimes \mathbf{3} = \mathbf{3} \oplus \mathbf{3}', \quad \text{with} \quad \begin{cases} \mathbf{3} = \begin{pmatrix} \alpha_1\beta_2 + \alpha_2\beta_3 \\ \alpha_1\beta_3 + \alpha_2\beta_1 \\ \alpha_1\beta_1 + \alpha_2\beta_2 \end{pmatrix} \\ \mathbf{3}' = \begin{pmatrix} \alpha_1\beta_2 - \alpha_2\beta_3 \\ \alpha_1\beta_3 - \alpha_2\beta_1 \\ \alpha_1\beta_1 - \alpha_2\beta_2 \end{pmatrix} \end{cases}$$

$$\mathbf{2} \otimes \mathbf{3}' = \mathbf{3} \oplus \mathbf{3}', \quad \text{with} \quad \begin{cases} \mathbf{3} = \begin{pmatrix} \alpha_1\beta_2 - \alpha_2\beta_3 \\ \alpha_1\beta_3 - \alpha_2\beta_1 \\ \alpha_1\beta_1 - \alpha_2\beta_2 \end{pmatrix} \\ \mathbf{3}' = \begin{pmatrix} \alpha_1\beta_2 + \alpha_2\beta_3 \\ \alpha_1\beta_3 + \alpha_2\beta_1 \\ \alpha_1\beta_1 + \alpha_2\beta_2 \end{pmatrix} \end{cases}$$

Finally, for the products of the triplet representations **3** and **3'**, we find

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{3}' \otimes \mathbf{3}' = \mathbf{1} \oplus \mathbf{2} \oplus \mathbf{3} \oplus \mathbf{3}', \quad \text{with} \quad \begin{cases} \mathbf{1} = \alpha_1\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_2 \\ \mathbf{2} = \begin{pmatrix} \alpha_2\beta_2 + \alpha_1\beta_3 + \alpha_3\beta_1 \\ \alpha_3\beta_3 + \alpha_1\beta_2 + \alpha_2\beta_1 \end{pmatrix} \\ \mathbf{3} = \begin{pmatrix} \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_1\beta_2 - \alpha_2\beta_1 \\ \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix} \\ \mathbf{3}' = \begin{pmatrix} 2\alpha_1\beta_1 - \alpha_2\beta_3 - \alpha_3\beta_2 \\ 2\alpha_3\beta_3 - \alpha_1\beta_2 - \alpha_2\beta_1 \\ 2\alpha_2\beta_2 - \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix} \end{cases}$$

$$\mathbf{3} \otimes \mathbf{3}' = \mathbf{1}' \oplus \mathbf{2} \oplus \mathbf{3} \oplus \mathbf{3}', \quad \text{with} \quad \left\{ \begin{array}{l} \mathbf{1} = \alpha_1\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_2 \\ \mathbf{2} = \begin{pmatrix} \alpha_2\beta_2 + \alpha_1\beta_3 + \alpha_3\beta_1 \\ -(\alpha_3\beta_3 + \alpha_1\beta_2 + \alpha_2\beta_1) \end{pmatrix} \\ \mathbf{3} = \begin{pmatrix} 2\alpha_1\beta_1 - \alpha_2\beta_3 - \alpha_3\beta_2 \\ 2\alpha_3\beta_3 - \alpha_1\beta_2 - \alpha_2\beta_1 \\ 2\alpha_2\beta_2 - \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix} \\ \mathbf{3}' = \begin{pmatrix} \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_1\beta_2 - \alpha_2\beta_1 \\ \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix} \end{array} \right.$$

We note that the CG coefficients presented above are in accordance with the results of Refs. [19, 40,41].

References

- [1] F.P. An, et al., DAYA-BAY Collaboration, Phys. Rev. Lett. 108 (2012) 171803, arXiv:1203.1669 [hep-ex]; F.P. An, et al., Chin. Phys. C 37 (2013) 011001, arXiv:1210.6327 [hep-ex].
- [2] J.K. Ahn, et al., RENO Collaboration, Phys. Rev. Lett. 108 (2012) 191802, arXiv:1204.0626 [hep-ex].
- [3] Y. Abe, et al., DOUBLE-CHOOZ Collaboration, Phys. Rev. Lett. 108 (2012) 131801, arXiv:1112.6353 [hep-ex]; Y. Abe, et al., Phys. Rev. D 86 (2012) 052008, arXiv:1207.6632 [hep-ex].
- [4] K. Abe, et al., T2K Collaboration, Phys. Rev. Lett. 107 (2011) 041801, arXiv:1106.2822 [hep-ex].
- [5] P. Adamson, et al., MINOS Collaboration, Phys. Rev. Lett. 107 (2011) 181802, arXiv:1108.0015 [hep-ex].
- [6] F.P. An, et al., Daya Bay Collaboration, arXiv:1310.6732 [hep-ex].
- [7] R.d.A. Toorop, F. Feruglio, C. Hagedorn, Phys. Lett. B 703 (2011) 447, arXiv:1107.3486 [hep-ph]; R. de Adelhart Toorop, F. Feruglio, C. Hagedorn, Nucl. Phys. B 858 (2012) 437, arXiv:1112.1340 [hep-ph].
- [8] G.-J. Ding, Nucl. Phys. B 862 (2012) 1, arXiv:1201.3279 [hep-ph].
- [9] S.F. King, C. Luhn, A.J. Stuart, Nucl. Phys. B 867 (2013) 203, arXiv:1207.5741 [hep-ph].
- [10] G.-J. Ding, S. Morisi, J.W.F. Valle, Phys. Rev. D 87 (2013) 053013, arXiv:1211.6506 [hep-ph].
- [11] D.V. Forero, M. Tortola, J.W.F. Valle, Phys. Rev. D 86 (2012) 073012, arXiv:1205.4018 [hep-ph].
- [12] G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, A.M. Rotunno, Phys. Rev. D 86 (2012) 013012, arXiv:1205.5254 [hep-ph]; F. Capozzi, G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, arXiv:1312.2878 [hep-ph].
- [13] M.C. Gonzalez-Garcia, M. Maltoni, J. Salvado, T. Schwetz, J. High Energy Phys. 1212 (2012) 123, arXiv:1209.3023 [hep-ph].
- [14] M. Bass, et al., LBNE Collaboration, arXiv:1311.0212 [hep-ex].
- [15] G. Altarelli, F. Feruglio, Rev. Mod. Phys. 82 (2010) 2701, arXiv:1002.0211 [hep-ph]; H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, M. Tanimoto, Prog. Theor. Phys. Suppl. 183 (2010) 1, arXiv:1003.3552 [hep-th]; S.F. King, C. Luhn, Rep. Prog. Phys. 76 (2013) 056201, arXiv:1301.1340 [hep-ph].
- [16] G. Ecker, W. Grimus, W. Konetschny, Nucl. Phys. B 191 (1981) 465; G. Ecker, W. Grimus, H. Neufeld, Nucl. Phys. B 247 (1984) 70; G. Ecker, W. Grimus, H. Neufeld, J. Phys. A 20 (1987) L807; H. Neufeld, W. Grimus, G. Ecker, Int. J. Mod. Phys. A 3 (1988) 603.
- [17] F. Feruglio, C. Hagedorn, R. Ziegler, J. High Energy Phys. 1307 (2013) 027, arXiv:1211.5560 [hep-ph].
- [18] M. Holthausen, M. Lindner, M.A. Schmidt, J. High Energy Phys. 1304 (2013) 122, arXiv:1211.6953 [hep-ph].
- [19] G.-J. Ding, S.F. King, C. Luhn, A.J. Stuart, J. High Energy Phys. 1305 (2013) 084, arXiv:1303.6180 [hep-ph].
- [20] F. Feruglio, C. Hagedorn, R. Ziegler, arXiv:1303.7178 [hep-ph].
- [21] C. Luhn, Nucl. Phys. B 875 (2013) 80, arXiv:1306.2358 [hep-ph].
- [22] G.-J. Ding, S.F. King, A.J. Stuart, J. High Energy Phys. 1312 (2013) 006, arXiv:1307.4212 [hep-ph].
- [23] P.F. Harrison, W.G. Scott, Phys. Lett. B 535 (2002) 163, arXiv:hep-ph/0203209; P.F. Harrison, W.G. Scott, Phys. Lett. B 547 (2002) 219, arXiv:hep-ph/0210197; P.F. Harrison, W.G. Scott, Phys. Lett. B 594 (2004) 324, arXiv:hep-ph/0403278; W. Grimus, L. Lavoura, Phys. Lett. B 579 (2004) 113, arXiv:hep-ph/0305309; Y. Farzan, A.Y. Smirnov, J. High Energy Phys. 0701 (2007) 059, arXiv:hep-ph/0610337;

- S. Gupta, A.S. Josphipura, K.M. Patel, Phys. Rev. D 85 (2012) 031903, arXiv:1112.6113 [hep-ph];
W. Grimus, L. Lavoura, Fortschr. Phys. 61 (2013) 535, arXiv:1207.1678 [hep-ph];
P.M. Ferreira, W. Grimus, L. Lavoura, P.O. Ludl, J. High Energy Phys. 1209 (2012) 128, arXiv:1206.7072.
- [24] R. Krishnan, P.F. Harrison, W.G. Scott, J. High Energy Phys. 1304 (2013) 087, arXiv:1211.2000 [hep-ph].
- [25] R.N. Mohapatra, C.C. Nishi, Phys. Rev. D 86 (2012) 073007, arXiv:1208.2875 [hep-ph].
- [26] I. Girardi, A. Meroni, S.T. Petcov, M. Spinrath, arXiv:1312.1966 [hep-ph].
- [27] G.C. Branco, J.M. Gerard, W. Grimus, Phys. Lett. B 136 (1984) 383;
I. de Medeiros Varzielas, D. Emmanuel-Costa, Phys. Rev. D 84 (2011) 117901, arXiv:1106.5477 [hep-ph];
I. de Medeiros Varzielas, D. Emmanuel-Costa, P. Leser, Phys. Lett. B 716 (2012) 193, arXiv:1204.3633 [hep-ph];
I. de Medeiros Varzielas, J. High Energy Phys. 1208 (2012) 055, arXiv:1205.3780 [hep-ph];
G. Bhattacharyya, I. de Medeiros Varzielas, P. Leser, Phys. Rev. Lett. 109 (2012) 241603, arXiv:1210.0545 [hep-ph];
I.P. Ivanov, L. Lavoura, Eur. Phys. J. C 73 (2013) 2416, arXiv:1302.3656 [hep-ph];
I.d.M. Varzielas, D. Pidt, arXiv:1307.0711 [hep-ph].
- [28] M.-C. Chen, K.T. Mahanthappa, Phys. Lett. B 681 (2009) 444, arXiv:0904.1721 [hep-ph];
A. Meroni, S.T. Petcov, M. Spinrath, Phys. Rev. D 86 (2012) 113003, arXiv:1205.5241 [hep-ph];
S. Antusch, S.F. King, M. Spinrath, Phys. Rev. D 87 (2013) 096018, arXiv:1301.6764 [hep-ph].
- [29] S. Antusch, S.F. King, C. Luhn, M. Spinrath, Nucl. Phys. B 850 (2011) 477, arXiv:1103.5930 [hep-ph];
S. Antusch, M. Holthausen, M.A. Schmidt, M. Spinrath, arXiv:1307.0710 [hep-ph];
S.F. King, arXiv:1311.3295 [hep-ph].
- [30] W. Grimus, M.N. Rebelo, Phys. Rep. 281 (1997) 239, arXiv:hep-ph/9506272.
- [31] C.H. Albright, A. Dueck, W. Rodejohann, Eur. Phys. J. C 70 (2010) 1099, arXiv:1004.2798 [hep-ph].
- [32] X.-G. He, A. Zee, Phys. Rev. D 84 (2011) 053004, arXiv:1106.4359 [hep-ph].
- [33] W. Rodejohann, H. Zhang, Phys. Rev. D 86 (2012) 093008, arXiv:1207.1225 [hep-ph].
- [34] I. de Medeiros Varzielas, L. Lavoura, J. Phys. G 40 (2013) 085002, arXiv:1212.3247.
- [35] W. Grimus, J. Phys. G 40 (2013) 075008, arXiv:1301.0495 [hep-ph].
- [36] J. Beringer, et al., Particle Data Group Collaboration, Phys. Rev. D 86 (2012) 010001.
- [37] G. Altarelli, F. Feruglio, Nucl. Phys. B 741 (2006) 215, arXiv:hep-ph/0512103.
- [38] S.F. King, Phys. Lett. B 718 (2012) 136, arXiv:1205.0506 [hep-ph];
S. Antusch, C. Gross, V. Maurer, C. Sluka, Nucl. Phys. B 866 (2013) 255, arXiv:1205.1051 [hep-ph].
- [39] H. Minakata, A.Y. Smirnov, Phys. Rev. D 70 (2004) 073009, arXiv:hep-ph/0405088;
N. Li, B.-Q. Ma, Eur. Phys. J. C 42 (2005) 17, arXiv:hep-ph/0504161;
N. Qin, B.Q. Ma, Phys. Lett. B 702 (2011) 143, arXiv:1106.3284 [hep-ph];
Y.H. Ahn, H.-Y. Cheng, S. Oh, Phys. Rev. D 83 (2011) 076012, arXiv:1102.0879 [hep-ph].
- [40] C. Hagedorn, S.F. King, C. Luhn, J. High Energy Phys. 1006 (2010) 048, arXiv:1003.4249 [hep-ph].
- [41] G.-J. Ding, Nucl. Phys. B 827 (2010) 82, arXiv:0909.2210 [hep-ph].