

Research Article

Perfect Fluid Dark Matter Influence on Thermodynamics and Phase Transition for a Reissner-Nordstrom-Anti-de Sitter Black Hole

Zhaoyi Xu ^{1,2,3,4} Xian Hou ^{1,3,4} Jiancheng Wang,^{1,2,3,4} and Yi Liao ^{5,6}

¹Yunnan Observatories, Chinese Academy of Sciences, 396 Yangfangwang, Guandu District, Kunming, 650216, China

²University of Chinese Academy of Sciences, Beijing, 100049, China

³Key Laboratory for the Structure and Evolution of Celestial Objects, Chinese Academy of Sciences, 396 Yangfangwang, Guandu District, Kunming, 650216, China

⁴Center for Astronomical Mega-Science, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing, 100012, China

⁵Department of Physics, National University of Defense Technology, Changsha, 410073, China

⁶Interdisciplinary Center for Quantum Information, National University of Defense Technology, Changsha, 410073, China

Correspondence should be addressed to Zhaoyi Xu; zyxu88@ynao.ac.cn, Xian Hou; xianhou.astro@gmail.com, and Yi Liao; liaoyitianyi@gmail.com

Received 5 June 2018; Accepted 23 December 2018; Published 23 January 2019

Guest Editor: Farook Rahaman

Copyright © 2019 Zhaoyi Xu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

Based on Reissner-Nordstrom-anti-de Sitter(RN-AdS) black hole surrounded by perfect fluid dark matter, we study the thermodynamics and phase transition by extending the phase space defined by the charge square Q^2 and the conjugate quantity ψ , where ψ is a function of horizon radius. The first law of thermodynamics and the equation of state are derived in the form $Q^2 = Q^2(T, \psi)$. By investigating the critical behaviour of perfect fluid dark matter around Reissner-Nordstrom-anti-de Sitter black hole, we find that these thermodynamics system are similar to Van der Waals system and can be explained by mean field theory. We also explore the Ruppeiner thermodynamic geometry feature and their connection with microscopic structure. We find that in extended phase space there are existence singularity points of Ruppeiner curvature and they could explained as phase transitions.

1. Introduction

Black hole thermodynamics are one of the most important topics in modern physics research and have been widely studied in recent years. The laws of black hole dynamics and thermodynamics were analyzed by Bekenstein and Hawking [1–5]. The four laws of black hole thermodynamics have been discussed [6]. Since the Hawking-Page phase transition was discovered, phase transitions have become an important topic in the black hole area. There are lots of work on the phase transition of different black holes, such as Reissner-Nordstrom black hole, Kerr black hole, and Kerr-Newman black hole [7–10]. These works have also been generalized to other black holes or applied to general situations [11–16]. Recently, several studies have considered the cosmological

constant as a dynamical variable which is similar to thermodynamical pressure [17]. Utilizing this method, some works have obtained the phase transition in AdS-black hole, in which the analogy between the critical behaviours of the Van der Waals gas and the RN-AdS black hole have been found [18–22].

From recent observations, we know that our universe is dominated by Dark Energy and Dark Matter [23, 24]. The dark energy makes the universe to be in accelerated expansion, and its state of equation is very close to the cosmological constant or the vacuum energy [25]. But the dynamics of dark energy are more like quintessence or other dynamical dark energy in behaviour [26]. The dark energy with quintessence could affect the black hole spacetime [27]. For the Schwarzschild black hole in quintessence field, the

modified black hole metric has been obtained by Kiselev [28, 29]. Recently the rotational quintessence black hole and Kerr-Newman-AdS black hole solutions have also been obtained [30, 31]. For the Schwarzschild black hole surrounded by quintessence matter, the thermodynamics and phase transition have been discussed in Tharanath et al. [32]; Ghaderi & Malakolkalami [33, 34]. For rotational black hole surrounded by quintessence case, the thermodynamics and phase transition have been studied recently by Xu & Wang [35]. The Reissner-Nordstrom black hole and Reissner-Nordstrom-dS black hole have been studied by Ghaderi & Malakolkalami [34]; Wei & Chu [36]; Wei & Ren [37]; Thomas et al. [38]; Mandal & Biswas [39]; Ma et al. [40]; the thermodynamics [41, 42] and phase transition through holography framework for Reissner-Nordstrom-AdS black hole have been investigated [43]. Following these works, the cold dark matter around black hole in phantom field background have been obtained by Li & Yang [44], and references therein. In this paper, we study the thermodynamics and phase transition of Reissner-Nordstrom-AdS black hole surround by perfect fluid dark matter.

This paper focuses on the study of the influence of perfect fluid dark matters on thermodynamics and phase transition for Reissner-Nordstrom-anti-de Sitter black hole. In Section 2, we introduce the Reissner-Nordstrom-anti-de Sitter black hole with perfect fluid dark matter background. In Section 3, we study the thermodynamical features, the equation of state in (Q^2, ψ) space, and the critical behaviour. In Section 4, we study the Ruppeiner thermodynamic geometry and their connection with microscopic structure for these black hole systems. We summarize our results in Section 5.

2. The Spacetime of Perfect Fluid Dark Matter around Reissner-Nordstrom-AdS Black Hole

We consider the dark matter field minimally coupled to gravity, electromagnetic field, and cosmological constant [28, 29, 44, 45]:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{16\pi G} R - \frac{1}{8\pi G} \Lambda + \frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \mathcal{L}_{DM} \right), \quad (1)$$

where G is the Newton gravity constant, Λ is the cosmological constant, $F_{\mu\nu}$ is the Faraday tensor of electromagnetic field, \mathcal{L}_{DM} is the dark matter Lagrangian density, and this dark matter can be any dark matter model. By variation, we obtain the field equation from action principle as

$$\begin{aligned} R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \Lambda g_{\mu\nu} &= -8\pi G (\overline{T}_{\mu\nu} + T_{\mu\nu}(DM)) \\ &= -8\pi G T_{\mu\nu}, \\ F^{\mu\nu}{}_{;\nu} &= 0, \\ F^{\mu\nu;\alpha} + F^{\nu\alpha;\mu} + F^{\alpha\mu;\nu} &= 0 \end{aligned} \quad (2)$$

where $\overline{T}_{\mu\nu}$ is the energy-momentum tensors of ordinary matter and $T_{\mu\nu}(DM)$ is the energy-momentum tensors of

dark matter. In the case of black holes surrounded by dark matter, we assume that dark matter is perfect fluid. The energy-momentum tensors can be written as $T^t_t = -\rho$, $T^r_r = T^\theta_\theta = T^\phi_\phi = p$ (where $T^\mu_\mu = g^{\mu\nu} T_{\mu\nu}$). In addition, for the simplest case, we assume $T^r_r = T^\theta_\theta = T^\phi_\phi = T^t_t(1 - \delta)$, where δ is a constant. We refer to such dark matter as the ‘‘perfect fluid dark matter’’ in this work.

For the Reissner-Nordstrom-AdS spacetime metric in perfect fluid DM, the black hole solution is as follows [28, 29, 44]:

$$ds^2 = -f(r) dt^2 + f^{-1}(r) dr^2 + r^2 (d\theta^2 + \sin^2\theta d\phi^2), \quad (3)$$

where

$$f(r) = 1 - \frac{2M}{r} + \frac{Q^2}{r^2} + \frac{1}{3}\Lambda r^2 + \frac{\alpha}{r} \ln\left(\frac{r}{|\alpha|}\right), \quad (4)$$

where α is a parameter describing the intensity of the perfect fluid DM, M is the black hole mass, and Q is the charge of black hole. This solution corresponds to a specific case of the general solution in Kiselev [28, 29] and Li & Yang [44]. It is interesting to note that this black hole solution implies that the rotational velocity is asymptotically flat in the equatorial plane, which could explain the observed rotation curves in spiral galaxies [28, 29, 44].

3. Thermodynamics of Dark Matter around Reissner-Nordstrom-AdS Black Hole

In Section 2, we have obtained the metric of spherically symmetric Reissner-Nordstrom-AdS black hole in perfect fluid dark matter. These black holes have three horizons which are Cauchy horizon r_- , event horizon r_+ , and cosmological horizon r_Λ . In this work, we always use event horizon r_+ . The black hole mass M can be expressed in event horizon r_+ as

$$M = \frac{r_+}{2} + \frac{Q^2}{2r_+} + \frac{1}{6}\Lambda r_+^3 + \frac{\alpha}{2} \ln\left(\frac{r_+}{|\alpha|}\right). \quad (5)$$

The semi-hawking temperature and the entropy are given by

$$\begin{aligned} T &= \frac{1}{4\pi} \left. \frac{df(r)}{dr} \right|_{r=r_+} \\ &= \frac{r_+ (2M + \alpha (1 - \ln(r_+/|\alpha|)) + (2/3)\Lambda r_+^3) - 2Q^2}{4\pi r_+^3} \\ &= \frac{r_+ (\alpha + r_+ + \Lambda r_+^3) - Q^2}{4\pi r_+^3}, \end{aligned} \quad (6)$$

$$S = \int_0^{r_+} \frac{1}{T} \left(\frac{\partial M}{\partial r_+} \right) dr_+ = \pi r_+^2. \quad (7)$$

Now we study the thermodynamical properties of the black hole with perfect fluid dark matter by extending to new phase space. This phase space is constructed by the entropy S , the perfect fluid dark matter density α , the charge square Q^2 , and the pressure $P = -\Lambda/8\pi$ corresponding to the

cosmological constant Λ . Therefore the black hole mass can be expressed as

$$M(S, Q^2, P, \alpha) = \frac{1}{2} \sqrt{\frac{S}{\pi}} + \frac{Q^2}{2} \sqrt{\frac{\pi}{S}} - \frac{4}{3} PS \sqrt{\frac{S}{\pi}} + \frac{\alpha}{2} \ln \left(\frac{1}{|\alpha|} \sqrt{\frac{S}{\pi}} \right). \quad (8)$$

The intensive parameters are defined by

$$\begin{aligned} T &= \left. \frac{\partial M}{\partial S} \right|_{P, Q^2, \alpha}, \\ \psi &= \left. \frac{\partial M}{\partial Q^2} \right|_{S, P, \alpha}, \\ V &= \left. \frac{\partial M}{\partial P} \right|_{S, Q^2, \alpha}, \\ \Pi &= \left. \frac{\partial M}{\partial \alpha} \right|_{S, Q^2, P}, \end{aligned} \quad (9)$$

where T denotes the temperature and the new physical quantity ψ is related to the specific volume as $\psi = 1/\nu$, where $\nu = 2r_+$. In thermodynamical space, the volume is $V = 4\pi r_+^3/3$ and the quantity $\Pi = (1/2) \ln(r_+/|\alpha|)$. The generalized first law of black hole thermodynamics in this extended phase space is expressed by

$$dM = TdS + \psi dQ^2 + VdP + \Pi d\alpha, \quad (10)$$

and the generalized Smarr formula is given by

$$M = 2TS + \psi Q^2 - 2VP + \Pi\alpha. \quad (11)$$

For the first law of black hole thermodynamics, the term ψdQ becomes ψdQ^2 in this phase space, where ψ represents the electric potential. This change leads to the interesting behaviour with $dM = SdT + \psi dQ$ in formal phase space. When the perfect fluid dark matter is around black hole, the phase transition of black hole occurs in (P, ν) plane. We can discuss this phase transition in (Q^2, ψ) plane, including the critical point, Gibbs free energy, and critical exponents under the effect of perfect fluid dark matter. Through calculations, we obtain the state equation $Q^2(T, \psi)$ as

$$\begin{aligned} Q^2 &= r_+ \left(\alpha + r_+ + \Lambda r_+^3 - 4\pi r_+^2 T \right) \\ &= \frac{1}{2\psi} \left[\alpha + \frac{1}{2\psi} + \frac{3}{8\psi^3 l^2} - \frac{\pi}{\psi^2} T \right], \end{aligned} \quad (12)$$

where $l^2 = 3/\Lambda$. The above equation describes the behaviours of Q^2 for different T, ψ and α .

Q^2 and ψ also satisfy the Maxwell equal area theorem given by Spallucci & Smailagic [46]:

$$\oint \psi dQ^2 = 0. \quad (13)$$

For $T > T_c$, where T_c critical temperature, there is an inflection point which is similar to the Van der Waals system. From general method, the coordinates of the critical point are determined by the following:

$$\begin{aligned} \left. \frac{\partial^2 Q^2}{\partial \psi^2} \right|_{T_c} &= 0, \\ \left. \frac{\partial Q^2}{\partial \psi} \right|_{T_c} &= 0. \end{aligned} \quad (14)$$

We then obtain the following equations:

$$\begin{aligned} \frac{15}{4\psi_c^3 l^2} + \alpha + \frac{3}{2\psi_c} - \frac{6\pi}{\psi_c^2} T_c &= 0, \\ -\frac{3}{2\psi_c^3 l^2} - \alpha - \frac{1}{\psi_c} + \frac{3\pi}{\psi_c^2} T_c &= 0. \end{aligned} \quad (15)$$

We then get the equation of critical ψ_c and T_c as

$$\begin{aligned} \alpha \psi_c^3 + \frac{1}{2} \psi_c^2 - \frac{3}{4l^2} &= 0, \\ T_c &= \frac{\psi_c^2}{3\pi} \left(\frac{3}{2\psi_c^3 l^2} + \alpha + \frac{1}{\psi_c} \right). \end{aligned} \quad (16)$$

The universal number is

$$\begin{aligned} \rho_c &= Q_c^2 T_c \psi_c \\ &= \frac{\psi_c^2}{3\pi} \left(\alpha + \frac{1}{4} + \frac{3}{2\psi_c^3 l^2} \right) \left(\frac{\alpha}{3} + \frac{1}{12\psi_c} - \frac{1}{16\psi_c^3 l^2} \right), \end{aligned} \quad (17)$$

where $0 < \alpha < 2$ for perfect fluid dark matter. When $\alpha = 0$, we have the following values:

$$\begin{aligned} T_c &= \frac{\sqrt{6}}{3\pi l}, \\ Q_c^2 &= \frac{l^2}{36}, \\ \psi_c &= \sqrt{\frac{3}{2l^2}}, \end{aligned} \quad (18)$$

which have been obtained by Dehyadegari et al. [47].

Now we study the critical behaviour near the phase transition with perfect fluid dark matter in new phase space. We first define $\psi_r = \psi/\psi_c$, $Q_r^2 = Q^2/Q_c^2$, and $T_r = T/T_c$; we then get $\psi_r = 1 + \chi$, $Q_r^2 = 1 + \varrho$, and $T_r = 1 + t$, where χ, ϱ and t represent the deviations away from critical points. The thermodynamics quantities are defined as

$$\begin{aligned} C_\psi &= |t|^{-a}, \\ \eta &= |t|^b, \\ \kappa_T &= |t|^{-c} \end{aligned} \quad (19)$$

$$|Q^2 - Q_c^2| = |\psi - \psi_c|^d$$

where a, b, c and d are the critical exponents. The critical exponent a is derived by fixing potential ψ for heat capacity as

$$C_\psi = T \left. \frac{\partial S}{\partial T} \right|_\psi = 0, \quad (20)$$

$$a = 0.$$

The critical exponent b is derived from Q^2 as

$$Q_r^2 = \frac{1}{2Q_c^2 \psi_c \psi_r} \left(\alpha + \frac{1}{2\psi_r \psi_c} + \frac{3}{8l^2 \psi_r^3 \psi_c^3} - \frac{\pi T_c}{\psi_r^2 \psi_c^2} T_r \right) \quad (21)$$

$$= \frac{F_1(\psi_c)}{\psi_r} + \frac{F_2(\psi_c)}{\psi_r^2} + \frac{F_3(\psi_c)}{\psi_r^3} + \frac{F_4(\psi_c)}{\psi_r^4} T_r,$$

where $F_1(\psi_c) = \alpha/(2Q_c^2 \psi_c)$, $F_2(\psi_c) = 1/(4Q_c^2 \psi_c^2)$, $F_3(\psi_c) = 3/(16Q_c^2 \psi_c^4 l^2)$, $F_4(\psi_c) = -\pi T_c/(2Q_c^2 \psi_c^3)$, and the critical points ψ_c, Q_c^2, T_c are functions of α, l^2 . In order to obtain critical exponent, we expand the equation near the critical point using $\psi_r = 1 + \chi$, $T_r = 1 + t$ and $Q_r^2 = 1 + \varrho$, and obtain

$$\varrho = F_4(\psi_c)t - 4F_4(\psi_c)t\chi$$

$$- (F_1(\psi_c) + 4F_2(\psi_c) + 4F_3(\psi_c) + 50F_4(\psi_c))\chi^3 \quad (22)$$

+ high order term.

Through differentiating (22) with respect to χ and χ , and using (13), we obtain $\chi_s = -\chi_l = \sqrt{-4F_4(\psi_c)t/(F_1(\psi_c) + 4F_2(\psi_c) + 4F_3(\psi_c) + 50F_4(\psi_c))}$, where l and s represent large and small black hole phase, respectively. We then find that

$$|\chi_s - \chi_l| = 2\chi_s$$

$$= \sqrt{\frac{-16F_4(\psi_c)}{F_1(\psi_c) + 4F_2(\psi_c) + 4F_3(\psi_c) + 50F_4(\psi_c)}} t^{1/2}, \quad (23)$$

$$b = \frac{1}{2}.$$

The critical exponent c is derived from isothermal compressibility coefficient κ_T as

$$\kappa_T = \left. \frac{\partial \psi}{\partial Q^2} \right|_T \propto \frac{\psi_c}{-4F_4(\psi_c) Q_c^2 t}, \quad c = 1, \quad (24)$$

The critical exponent d is obtained from (22) as

$$\varrho|_{t=0}$$

$$= - (F_1(\psi_c) + 4F_2(\psi_c) + 4F_3(\psi_c) + 50F_4(\psi_c)) \chi^3, \quad (25)$$

$$d = 3.$$

From the above analysis, we find that these critical exponents resemble those in Van der Waals system, implying that the critical phenomenon can be explained by mean

field theory [17]. These critical exponents satisfy the scale symmetry given by

$$a + 2b + c = 2,$$

$$a + b(d + 1) = 2,$$

$$c(d + 1) = (2 - a)(d - 1), \quad (26)$$

$$c = b(d - 1).$$

The perfect fluid dark matters around black holes could explain the formation of supermassive black holes in approximate stationary situation. In the past decades, some high redshift quasars have been discovered and the centre supermassive black hole with the mass beyond 10 billion of solar masses. Usually such black hole is difficult to form in the Universe less than one billion years old. Because perfect fluid dark matter black holes satisfy the first law of thermodynamics, the perfect fluid dark matter could accelerate the formation of supermassive black holes.

4. Geothermodynamics of Dark Matter around Reissner-Nordstrom-AdS Black Hole

In the black hole thermodynamics, the thermodynamic geometry method is a usual tool to understand the property of black hole thermodynamics system. In the work, we use the Ruppeiner metric to study the thermodynamical effect of the perfect fluid dark matter on the microscopical structure of Reissner-Nordstrom-AdS black hole [48]. We define the metric on (M, Q^2) space given by

$$g_{\mu\nu}(R) = \frac{1}{T} \frac{\partial^2 M}{\partial X^\mu \partial X^\nu} = \begin{pmatrix} \frac{1}{T} \frac{\partial^2 M}{\partial S^2} & \frac{1}{T} \frac{\partial^2 M}{\partial S \partial Q^2} \\ \frac{1}{T} \frac{\partial^2 M}{\partial S \partial Q^2} & \frac{1}{T} \frac{\partial^2 M}{\partial Q^4} \end{pmatrix} \quad (27)$$

where $X^\mu = (S, Q^2)$.

From (5) and (6), we find that $M(S, Q^2)$ and $T(S, Q^2)$ in (M, Q^2) space are given by

$$M(S, Q^2) = \frac{1}{2} \sqrt{\frac{S}{\pi}} + \frac{Q^2}{2} \sqrt{\frac{\pi}{S}} + \frac{1}{2l^2 \pi} S \sqrt{\frac{S}{\pi}}$$

$$+ \frac{\alpha}{2} \ln \left(\frac{1}{|\alpha|} \sqrt{\frac{S}{\pi}} \right), \quad (28)$$

$$T(S, Q^2) = \frac{\alpha}{4S} + \frac{3}{4\pi l^2} \sqrt{\frac{S}{\pi}} + \frac{1}{4\pi} \sqrt{\frac{\pi}{S}} - \frac{Q^2}{4S} \sqrt{\frac{\pi}{S}}. \quad (29)$$

From geometry, the Ricci scalar can be calculated for perfect fluid dark matter around Reissner-Nordstrom-AdS black holes by

$$\begin{aligned}
R(RN - AdSDM) &= R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} \\
&= \frac{H^2}{(-l^2\pi^2Q^2 - 2\alpha l^2\sqrt{\pi^3S} + l^2\pi S + 3S^2)^2} \\
&= \frac{H^2}{\left(-l^2\pi^2Q^2 - \frac{\alpha l^2\pi^2}{\psi} + \frac{l^2\pi^3}{4\psi^2} + \frac{3\pi^2}{16\psi^4}\right)^2},
\end{aligned} \tag{30}$$

where H are function of α , l^2 , Q^2 and S . From above equation, we find that the phase transition occurs when the following condition is satisfied:

$$-l^2\pi^2Q^2 - \frac{\alpha l^2\pi^2}{\psi} + \frac{l^2\pi^3}{4\psi^2} + \frac{3\pi^2}{16\psi^4} = 0. \tag{31}$$

We know that the sign of the Ricci scalar R can be explained by intermolecular interaction in thermodynamical system. The positive sign refers to the repulsive interaction between the constituents of the thermodynamical system, while the negative sign refers to the attractive interaction between the constituents of the thermodynamical system ([42] and references therein). For perfect fluid dark matter around Reissner-Nordstrom-AdS black hole, the interactions are absent in the thermodynamical system for a null Ricci scalar R , which is similar to that in classical ideal gas [49].

5. Summary

In the paper, we study the perfect fluid dark matter influence on thermodynamics and phase transition of Reissner-Nordstrom-AdS black hole by extending phase space defined by the charge square Q^2 and conjugate quantity ψ . The first law of thermodynamics and the equation of state are derived in the form of $Q^2 = Q^2(T, \psi)$. We analyze the critical behaviour of dark matter around Reissner-Nordstrom-AdS black hole and find that these thermodynamics system resemble the Van der Waals system which can be explained by mean field theory. We also find that the critical exponents satisfy the scale law of thermodynamical system. Using Ruppeiner thermodynamic geometry, we study the geometric property of perfect fluid dark matter around black holes. We find that, in extended phase space, some singular points appear on the Ruppeiner curvature, which can be explained as the critical points of phase transitions.

The Reissner-Nordstrom-AdS black hole surrounded by dark matters could appear in the Universe. In the future work we plan to study the observed effects of perfect fluid dark matter on black holes and the influence of perfect fluid dark matter on gravitational lensing and the evolution of dark matter in the universe.

Data Availability

No data were used to support this study.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The authors acknowledge the financial support from the National Natural Science Foundation of China, 11573060 and 11661161010.

References

- [1] J. D. Bekenstein, "Black holes and the second law," *Lettere Al Nuovo Cimento Series 2*, vol. 4, no. 15, pp. 737–740, 1972.
- [2] J. D. Bekenstein, "Generalized second law of thermodynamics in black-hole physics," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 9, no. 12, pp. 3292–3300, 1974.
- [3] J. D. Bekenstein, "Black holes and entropy," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 7, pp. 2333–2346, 1973.
- [4] S. W. Hawking, "Gravitational radiation from colliding black holes," *Physical Review Letters*, vol. 26, no. 21, pp. 1344–1346, 1971.
- [5] S. W. Hawking, "Black hole explosions?" *Nature*, vol. 284, no. 5443, pp. 30–31, 1974.
- [6] J. M. Bardeen, B. Carter, and S. W. Hawking, "The four laws of black hole mechanics," *Communications in Mathematical Physics*, vol. 31, pp. 161–170, 1973.
- [7] M. M. Caldarelli, G. Cognola, and D. Klemm, "Thermodynamics of Kerr-Newman-AdS black holes and conformal field theories," *Classical and Quantum Gravity*, vol. 17, no. 2, pp. 399–420, 2000.
- [8] P. C. Davies, "Thermodynamic phase transitions of Kerr-Newman black holes in de Sitter space," *Classical and Quantum Gravity*, vol. 6, no. 12, pp. 1909–1914, 1989.
- [9] P. C. W. Davies, "Thermodynamics of black holes," *Reports on Progress in Physics*, vol. 41, no. 8, pp. 1313–1355, 1978.
- [10] P. C. W. Davies, "The Thermodynamic Theory of Black Holes," *Proceedings of the Royal Society of London Series A*, vol. 353, pp. 499–521, 1977.
- [11] S. W. Hawking and D. N. Page, "Thermodynamics of black holes in anti-de Sitter space," *Communications in Mathematical Physics*, vol. 87, no. 4, pp. 577–588, 1983.
- [12] P. Hut, "Charged black holes and phase transitions," *Monthly Notices of the Royal Astronomical Society*, vol. 180, pp. 379–389, 1977.
- [13] X.-B. Gong, Y. Liao, and Z.-Y. Xu, "A mathematical form of force-free magnetosphere equation around Kerr black holes and its application to Meissner effect," *Physics Letters B*, vol. 760, pp. 112–116, 2016.
- [14] Y. Liao, X.-B. Gong, and J.-S. Wu, "Phase-transition Theory of Kerr Black Holes in the Electromagnetic Field," *The Astrophysical Journal*, vol. 835, no. 2, p. 247, 2017.
- [15] A. Mandal, S. Samanta, and B. Ranjan Majhi, "Phase transition and critical phenomena of black holes: A general approach," *Physical Review D*, vol. 94, Article ID 064069, 2016.
- [16] B. Ranjan Majhi and S. Samanta, "P-V criticality of AdS black holes in a general framework," *Physics Letters B*, vol. 773, pp. 203–207, 2017.
- [17] D. Kubiznák and R. B. Mann, "P – V criticality of charged AdS black holes," *Journal of High Energy Physics*, vol. 7, p. 33, 2012.
- [18] A. Belhaj, M. Chabab, H. El Moumni, K. Masmar, and M. B. Sedra, "On thermodynamics of AdS black holes in M-theory," *The European Physical Journal C*, vol. 76, no. 73, 2016.

- [19] A. Belhaj, M. Chabab, H. El Moumni, K. Masmr, and M. B. Sedra, "Maxwell's equal-area law for Gauss–Bonnet–Anti-de Sitter black holes," *The European Physical Journal C*, vol. 75, article 71, 2015.
- [20] A. Belhaj, M. Chabab, H. E. Moumni, and M. B. Sedra, "On thermodynamics of AdS black holes in arbitrary dimensions," *Chinese Physics Letters*, vol. 29, no. 10, Article ID 100401, 2012.
- [21] A. Belhaj, M. Chabab, H. El Moumni, L. Medari, and M. B. Sedra, "The Thermodynamical Behaviors of Kerr–Newman AdS Black Holes," *Chinese Physics Letters*, vol. 30, Article ID 090402, 2013.
- [22] M. Chabab, H. El Moumni, and K. Masmr, "On thermodynamics of charged AdS black holes in extended phases space via M2-branes background," *The European Physical Journal C*, vol. 76, p. 304, 2016.
- [23] E. Komatsu, K. M. Smith, J. Dunkley et al., "Seven-year wilkinson microwave anisotropy probe (WMAP*) observations: cosmological interpretation," *The Astrophysical Journal*, vol. 192, no. 2, p. 18, 2011.
- [24] A. G. Riess, L.-G. Strolger, J. Tonry et al., "Type Ia Supernova Discoveries at $z > 1$ from the Hubble Space Telescope: Evidence for Past Deceleration and Constraints on Dark Energy Evolution," *The Astrophysical Journal*, vol. 607, p. 665, 2004.
- [25] P. J. E. Peebles and B. Ratra, "The cosmological constant and dark energy," *Reviews of Modern Physics*, vol. 75, no. 2, pp. 559–606, 2003.
- [26] E. J. Copeland, M. Sami, and S. Tsujikawa, "Dynamics of dark energy," *International Journal of Modern Physics D*, vol. 15, no. 11, pp. 1753–1935, 2006.
- [27] Z. Stuchlík, "Influence of the RELICT Cosmological Constant on Accretion Discs," *Modern Physics Letters A*, vol. 20, no. 8, pp. 561–575, 2005.
- [28] V. V. Kiselev, "Quintessence and black holes," *Classical and Quantum Gravity*, vol. 20, no. 6, pp. 1187–1197, 2003.
- [29] V. V. Kiselev, "Quintessential solution of dark matter rotation curves and its simulation by extra dimensions," *High Energy Physics*, 11 pages, 2003.
- [30] B. Toshmatov, Z. Stuchlík, and B. Ahmedov, "Rotating black hole solutions with quintessential energy," 2015, <https://arxiv.org/abs/1512.01498v1>.
- [31] Z. Xu and J. Wang, "Kerr–Newman–AdS black hole in quintessential dark energy," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 95, Article ID 06415, 2017.
- [32] R. Tharanath, N. Varghese, and V. C. Kuriakose, "Phase transition, quasinormal modes and Hawking radiation of Schwarzschild black hole in quintessence field," *Modern Physics Letters A*, vol. 29, Article ID 1450057, 2014.
- [33] K. Ghaderi and B. Malakolkalami, "Effects of quintessence on thermodynamics of the black holes," *Astrophysics and Space Science*, vol. 361, p. 161, 2016.
- [34] K. Ghaderi and B. Malakolkalami, "Thermodynamics of the Schwarzschild and the Reissner–Nordström black holes with quintessence," *Nuclear Physics B*, vol. 903, pp. 10–18, 2016.
- [35] Z. Xu and J. Wang, "Thermodynamics and Phase Transition in Rotational Kiselev Black Hole," 2016, <https://arxiv.org/abs/1610.00376>.
- [36] Y.-H. Wei and Z.-H. Chu, "Thermodynamic Properties of a Reissner–Nordström Quintessence Black Hole," *Chinese Physics Letters*, vol. 28, Article ID 100403, 2011.
- [37] Y.-H. Wei and J. Ren, "Thermodynamic properties of Reissner–Nordström–de Sitter quintessence black holes," *Chinese Physics B*, vol. 22, Article ID 030402, 2013.
- [38] B. B. Thomas, M. Saleh, and T. C. Kofane, "Thermodynamics and phase transition of the Reissner–Nordström black hole surrounded by quintessence," *General Relativity and Gravitation*, vol. 44, no. 9, pp. 2181–2189, 2012.
- [39] A. Mandal and R. Biswas, "Effects of cosmic acceleration on black hole thermodynamics," *Astrophysics and Space Science*, vol. 357, p. 8, 2015.
- [40] M.-S. Ma, R. Zhao, and Y.-Q. Ma, "Thermodynamic stability of black holes surrounded by quintessence," *General Relativity and Gravitation*, vol. 49, p. 79, 2017.
- [41] P. Pradhan, "Thermodynamic Products in Extended Phase Space," *International Journal of Modern Physics D*, vol. 26, no. 2, 2017.
- [42] M. Chabab, H. El Moumni, S. Iraoui, K. Masmr, and S. Zhizeh, "More insight into microscopic properties of RN–AdS black hole surrounded by quintessence via an alternative extended phase space," *International Journal of Geometric Methods in Modern Physics*, vol. 15, no. 10, 2018.
- [43] X.-X. Zeng and L.-F. Li, "Van der Waals phase transition in the framework of holography," *Physics Letters B*, vol. 764, pp. 100–108, 2017.
- [44] M.-H. Li and K.-C. Yang, "Galactic dark matter in the phantom field," *Physical Review D: Particles, Fields, Gravitation and Cosmology*, vol. 86, Article ID 123015, 2012.
- [45] V. V. Kiselev, "Vector field and rotational curves in dark galactic halos," *Classical and Quantum Gravity*, vol. 22, no. 3, pp. 541–557, 2005.
- [46] E. Spallucci and A. Smailagic, "Maxwell's equal area law and the Hawking–Page phase transition," 2013, <https://arxiv.org/abs/1310.2186v1>.
- [47] A. Dehyadegari, A. Sheykhi, and A. Montakhab, "Critical behavior and microscopic structure of charged AdS black holes via an alternative phase space," *Physics Letters B*, vol. 768, pp. 235–240, 2017.
- [48] A. Sahay, T. Sarkar, and G. Sengupta, "Thermodynamic geometry and phase transitions in Kerr–Newman–AdS black holes," *Journal of High Energy Physics*, vol. 7, p. 82, 2010.
- [49] G. Ruppeiner, "Thermodynamics: a Riemannian geometric model," *Physical Review A: Atomic, Molecular and Optical Physics*, vol. 20, no. 4, pp. 1608–1613, 1979.